



# Existence of multiple solutions for a class of nonhomogeneous problems with critical growth

Mohammed Massar 

Mohamed First University, Faculty of Sciences & Techniques, Al Hoceima, Morocco

Received 27 April 2016, appeared 6 April 2017

Communicated by Gabriele Bonanno

**Abstract.** In this paper, we study the existence and multiplicity of solutions for  $(p_1(x), p_2(x))$ -equation with critical growth. The technical approach is mainly based on the variational method combined with the genus theory.

**Keywords:** nonhomogeneous, multiple solutions, critical growth.

**2010 Mathematics Subject Classification:** 35J62, 35J66, 35D30.

## 1 Introduction

In this article, we are concerned with the following problem

$$\begin{cases} -\Delta_{p_1(x)}u - \Delta_{p_2(x)}u - a(x)|u|^{m(x)-2}u = \lambda|u|^{q(x)-2}u + f(x, u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (P_\lambda)$$

where  $\Omega \subset \mathbb{R}^N (N \geq 1)$  is a bounded smooth domain,  $\lambda$  is a positive parameter and  $f : \overline{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function which satisfies some assumptions provided later. Moreover,  $p_1, p_2, q \in C(\overline{\Omega})$  and  $m(x) = \max(p_1(x), p_2(x))$  for all  $x \in \overline{\Omega}$ , such that


$$\begin{aligned} 1 < p_i^- = \min_{x \in \overline{\Omega}} p_i(x) \leq p_i^+ = \max_{x \in \overline{\Omega}} p_i(x) < N, \quad i = 1, 2, \\ m^+ = \max_{x \in \overline{\Omega}} m(x) < q^- = \min_{x \in \overline{\Omega}} q(x) \leq q(x) \leq m^*(x) \quad \forall x \in \overline{\Omega}, \end{aligned} \quad (mq_1)$$

where  $m^*(x) = \frac{Nm(x)}{N-m(x)}$  for all  $x \in \overline{\Omega}$  and the set

$$B = \{x \in \Omega : q(x) = m^*(x)\} \text{ is nonempty.} \quad (mq_2)$$

In recent years, the study of various mathematical problems with variable exponent growth condition has been received considerable attention in recent years. A more and more important number of surveys and books dealing with this type of problems and their corresponding functional spaces setting have been published (see [1–4, 12, 16–20]). We also have to mention

---

 Email: massarmed@hotmail.com

the books [13] and [21] as important references in this field. This great interest may be justified by their various physical applications. In fact, there are applications concerning elastic mechanics [25], electrorheological fluids [23, 24], image restoration [9], dielectric breakdown, electrical resistivity and polycrystal plasticity [6, 7] and continuum mechanics [5].

It is well known that although most of the materials can be accurately modeled with the help of the classical Lebesgue and Sobolev spaces  $L^p$  and  $W^{1,p}$ , where  $p$  is a fixed constant, but there are some nonhomogeneous materials, for which this is not adequate, e.g. the rheological fluids mentioned above, which are characterized by their ability to drastically change their mechanical properties under the influence of an exterior electromagnetic field. Thus it is necessary for the exponent  $p$  to be variable, hence the need for spaces with variable exponents. This leads, on the one hand, to many interesting applications, and, on the other hand, to the study of much more mathematically complicated problems.

In [19], Mihăilescu considered the problem

$$\begin{cases} -\Delta_{p_1(x)}u - \Delta_{p_2(x)}u = \pm(-\lambda|u|^{m(x)-2}u + |u|^{q(x)-2}u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $m(x) = \max\{p_1(x), p_2(x)\}$  for any  $x \in \bar{\Omega}$  or  $m(x) < q(x) < \frac{Nm(x)}{N-m(x)}$  for any  $x \in \bar{\Omega}$ . In the first case, using mountain pass theorem, he established the existence of infinity many solutions. In the second case, by simple variational arguments, he proved that the problem has a solution for  $\lambda$  large enough. The novelty of this paper lies in the fact we consider problem  $(P_\lambda)$ , with growth  $q(x)$  which is critical in a set with positive measure. The difficulty in this case, is due to the lack of compactness of the imbedding  $W_0^{1,m(x)}(\Omega) \hookrightarrow L^{m^*(x)}(\Omega)$  and the Palais–Smale condition for the corresponding energy functional could not be checked directly. To deal with this difficulty, we use a version of the concentration compactness lemma due to Lions for variable exponents [8].

Here, we are interested in the existence and multiplicity of weak solutions under the following hypotheses on  $a(x)$  and  $f$ .

(a<sub>1</sub>)  $a(x) \in L^\infty(\Omega)$  and there exists  $\alpha > 0$  such that

$$\int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} - a(x) \frac{|u|^{m(x)}}{m(x)} \right) dx \geq \alpha \int_{\Omega} \frac{|u|^{m(x)}}{m(x)} dx, \quad \forall u \in W_0^{1,m(x)}(\Omega);$$

(a<sub>2</sub>)  $m(x) = m^+$  for all  $x \in \Gamma^+ := \{x \in \Omega : a(x) > 0\}$ ;

(f<sub>1</sub>)  $f \in C(\bar{\Omega} \times \mathbb{R}, \mathbb{R})$ , odd with respect to  $t$  such that

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{f(x, t)}{|t|^{m(x)-1}} &= 0, \quad \text{uniformly in } x \in \bar{\Omega}, \\ \lim_{|t| \rightarrow +\infty} \frac{f(x, t)}{|t|^{q(x)-1}} &= 0, \quad \text{uniformly in } x \in \bar{\Omega}; \end{aligned}$$

(f<sub>2</sub>)  $F(x, t) \leq \frac{1}{m^+} f(x, t)t, \forall t \in \mathbb{R}$  and a.e.  $x \in \Omega$ , where  $F(x, t) = \int_0^t f(x, s)ds$ .

**Example 1.1.** In this example we just exhibit a function  $a(x)$  satisfying assumption (a<sub>1</sub>). Let  $\Omega = B(0, 2) := \{x \in \mathbb{R}^N : |x| < 2\}$  ( $N \geq 3$ ),  $p_1(x) = 2 - \frac{1}{4}x_1 - \frac{1}{16}|x|^2$  and  $p_2(x) = 2 - \frac{1}{4}x_2 - \frac{1}{16}|x|^2$  for all  $x = (x_1, x_2, \dots, x_N) \in \bar{\Omega}$ , where  $|x|^2 = \sum_{i=1}^N x_i^2$ . Then  $p_i \in C(\bar{\Omega})$

and  $1 < p_i^- \leq p_i^+ < N$ ,  $i = 1, 2$ . Let  $e_1 = (1, 0, \dots, 0)$ . Then for any  $x \in \Omega$ , the function  $h_1 : t \mapsto p_1(x + te_1)$  is monotone in  $I_x = \{t : x + te_1 \in \Omega\}$ . In fact, we have

$$h_1(t) = 2 - \frac{1}{4}(x_1 + t) - \frac{1}{16}(x_1 + t)^2 - \frac{1}{16} \sum_{i=2}^N x_i^2,$$

thus  $h_1'(t) = -\frac{1}{8}(2 + x_1 + t)$ . Since  $|x_1 + t| \leq |x + te_1| < 2$ ,  $h_1'(t) < 0$  for all  $t \in I_x$ , and hence  $h_1$  is decreasing in  $I_x$ . Similarly, for  $e_2 = (0, 1, \dots, 0)$ , the function  $h_2 : t \mapsto p_2(x + te_2)$  is monotone in  $I_x = \{t : x + te_2 \in \Omega\}$ . Therefore  $p_i$ ,  $i = 1, 2$ , satisfying conditions of [15, Theorem 3.3], thus the modular Poincaré inequality holds:

$$\int_{\Omega} |\nabla u|^{p_i(x)} dx \geq C_i \int_{\Omega} |u|^{p_i(x)} dx \quad (C_i > 0, i = 1, 2),$$

which is equivalent to

$$\int_{\Omega} \frac{|\nabla u|^{p_i(x)}}{p_i(x)} dx \geq C'_i \int_{\Omega} \frac{|u|^{p_i(x)}}{p_i(x)} dx \quad (C'_i > 0, i = 1, 2),$$

since  $1 < p_i^- \leq p_i(x) \leq p_i^+ < N$ .

Now, let  $V \in L^\infty(\Omega)$  such that  $1 + V(x) > c > 0$ . Note that  $p_i$ ,  $i = 1, 2$ , are log-Hölder continuous functions. Indeed, let  $x, y \in \Omega$  such that  $|x - y| \leq \frac{1}{2}$ , then

$$\begin{aligned} |p_1(x) - p_1(y)| &\leq \frac{1}{4}|x_1 - y_1| + \frac{1}{16}||x|^2 - |y|^2| \\ &\leq \frac{1}{4}|x - y| + \frac{1}{16}||x| - |y||(|x| + |y|) \\ &\leq \frac{1}{4}|x - y| + \frac{1}{16}|x - y|(|x| + |y|) \\ &\leq \frac{1}{2}|x - y| \\ &\leq \frac{1}{2 \log\left(\frac{1}{|x-y|}\right)}. \end{aligned}$$

In the same way, we get  $|p_2(x) - p_2(y)| \leq \frac{1}{2 \log\left(\frac{1}{|x-y|}\right)}$ . Applying [11, Theorem 2.2], we can find  $\alpha_1, \alpha_2 > 0$  such that

$$\int_{\Omega} \frac{|\nabla u|^{p_1(x)}}{p_1(x)} dx \geq \alpha_1 \int_{\Omega} (1 + V(x)) \frac{|u|^{p_1(x)}}{p_1(x)} dx$$

and

$$\int_{\Omega} \frac{|\nabla u|^{p_2(x)}}{p_2(x)} dx \geq \alpha_2 \int_{\Omega} (1 + V(x)) \frac{|u|^{p_2(x)}}{p_2(x)} dx.$$

Observe that  $|u|^{p_1(x)} + |u|^{p_2(x)} \geq |u|^{m(x)}$ , thus  $\frac{|u|^{p_1(x)}}{p_1(x)} + \frac{|u|^{p_2(x)}}{p_2(x)} \geq \frac{|u|^{m(x)}}{m(x)}$ . It follows that

$$\begin{aligned} \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} \right) dx &\geq \min(\alpha_1, \alpha_2) \int_{\Omega} (1 + V(x)) \left( \frac{|u|^{p_1(x)}}{p_1(x)} + \frac{|u|^{p_2(x)}}{p_2(x)} \right) dx \\ &\geq \min(\alpha_1, \alpha_2) \int_{\Omega} (1 + V(x)) \frac{|u|^{m(x)}}{m(x)} dx. \end{aligned}$$

Therefore

$$\int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} \right) dx - \int_{\Omega} \min(\alpha_1, \alpha_2) V(x) \frac{|u|^{m(x)}}{m(x)} dx \geq \min(\alpha_1, \alpha_2) \int \frac{|u|^{m(x)}}{m(x)} dx,$$

and hence the function  $a(x) := \min(\alpha_1, \alpha_2)V(x)$  satisfies condition  $(a_1)$ .

## 2 Preliminaries

Here, we state some interesting properties of the variable exponent Lebesgue and Sobolev spaces that will be useful to discuss problem  $(P_\lambda)$ . Every where below we consider  $\Omega \subset \mathbb{R}^N$  to be a bounded domain with smooth boundary and  $p(x) \in C_+(\overline{\Omega})$ , where

$$C_+(\Omega) = \{h \in C(\overline{\Omega}) : h(x) > 1 \text{ for all } x \in \overline{\Omega}\}.$$

Define the variable exponent Lebesgue space by

$$L^{p(x)}(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} \text{ measurable} : \int_{\Omega} |u(x)|^{p(x)} dx < \infty \right\}.$$

This space endowed with the Luxemburg norm,

$$\|u\|_{L^{p(x)}(\Omega)} = \inf \left\{ \tau > 0 : \int_{\Omega} \left| \frac{u(x)}{\tau} \right|^{p(x)} dx \leq 1 \right\}$$

is a separable and reflexive Banach space. Denoting by  $L^{p'(x)}(\Omega)$  the conjugate space of  $L^{p(x)}(\Omega)$  where  $\frac{1}{p(x)} + \frac{1}{p'(x)} = 1$ ; for any  $u \in L^{p(x)}(\Omega)$  and  $v \in L^{p'(x)}(\Omega)$  we have the following Hölder type inequality

$$\int_{\Omega} |uv| dx \leq \left( \frac{1}{p^-} + \frac{1}{p'^-} \right) \|u\|_{L^{p(x)}(\Omega)} \|v\|_{L^{p'(x)}(\Omega)}.$$

Now, we introduce the modular of the Lebesgue–Sobolev space  $L^{p(x)}(\Omega)$  as the mapping  $\rho_{p(x)} : L^{p(x)}(\Omega) \rightarrow \mathbb{R}$ , defined by

$$\rho_{p(x)}(u) = \int_{\Omega} |u|^{p(x)} dx, \quad \forall u \in L^{p(x)}(\Omega).$$

In the following proposition, we give some relations between the Luxemburg norm and the modular.

**Proposition 2.1** ([14]). *If  $u, u_n \in L^{p(x)}(\Omega)$ , then following properties hold true:*

- (1)  $\|u\|_{L^{p(x)}(\Omega)} \leq 1 \Rightarrow \|u\|_{L^{p(x)}(\Omega)}^{p^+} \leq \rho_{p(x)}(u) \leq \|u\|_{L^{p(x)}(\Omega)}^{p^-}$ ;
- (2)  $\|u\|_{L^{p(x)}(\Omega)} \geq 1 \Rightarrow \|u\|_{L^{p(x)}(\Omega)}^{p^-} \leq \rho_{p(x)}(u) \leq \|u\|_{L^{p(x)}(\Omega)}^{p^+}$ ;
- (3)  $\lim_{n \rightarrow \infty} \|u_n\|_{L^{p(x)}(\Omega)} = 0 \Leftrightarrow \lim_{n \rightarrow \infty} \rho_{p(x)}(u_n) = 0$ ;
- (4)  $\lim_{n \rightarrow \infty} \|u_n\|_{L^{p(x)}(\Omega)} = \infty \Leftrightarrow \lim_{n \rightarrow \infty} \rho_{p(x)}(u_n) = \infty$ .

Next, we define the variable exponent Sobolev space  $W^{1,p(x)}(\Omega)$  by

$$W^{1,p(x)}(\Omega) = \left\{ u \in L^{p(x)}(\Omega) : |\nabla u| \in L^{p(x)}(\Omega) \right\}.$$

endowed with the norm

$$\|u\|_{W^{1,p(x)}(\Omega)} = \|u\|_{L^{p(x)}(\Omega)} + \|\nabla u\|_{L^{p(x)}(\Omega)},$$

where by  $\|\nabla u\|_{L^{p(x)}(\Omega)}$  we understand  $\| |\nabla u| \|_{L^{p(x)}(\Omega)}$ . We denote by  $W_0^{1,p(x)}(\Omega)$  the closure of  $C_0^\infty(\Omega)$  in  $W^{1,p(x)}(\Omega)$  with respect to the above norm. The space  $W^{1,p(x)}(\Omega)$  and  $W_0^{1,p(x)}(\Omega)$  are separable and reflexive Banach spaces.

**Proposition 2.2** ([14]).

(1) If  $r \in C_+(\overline{\Omega})$  and  $r(x) < p^*(x)$  for all  $x \in \overline{\Omega}$ , then the embedding

$$W^{1,p(x)}(\Omega) \hookrightarrow L^{r(x)}(\Omega)$$

is compact and continuous.

(2) There is a constant  $C > 0$  such that

$$\|u\|_{L^{p(x)}(\Omega)} \leq C \|\nabla u\|_{L^{p(x)}(\Omega)} \quad \text{for all } u \in W_0^{1,p(x)}(\Omega).$$

### 3 Main result

From now on, we consider  $m(x) = \max(p_1(x), p_2(x))$  for all  $x \in \overline{\Omega}$ . By (2) of Proposition 2.2, we know that  $\|u\| := \|\nabla u\|_{L^{m(x)}(\Omega)}$  and  $\|u\|_{W^{1,m(x)}(\Omega)}$  are equivalent norms in  $W_0^{1,m(x)}(\Omega)$ . In the following, we will use  $\|u\|$  instead of  $\|u\|_{W^{1,m(x)}(\Omega)}$  on  $W_0^{1,m(x)}(\Omega)$ .

**Definition 3.1.** We say that  $u \in W_0^{1,m(x)}(\Omega)$  is a weak solution of problem  $(P_\lambda)$  if

$$\begin{aligned} & \int_{\Omega} \left( |\nabla u|^{p_1(x)-2} + |\nabla u|^{p_2(x)-2} \right) \nabla u \nabla v dx - \int_{\Omega} a(x) |u|^{m(x)-2} u v dx \\ & = \lambda \int_{\Omega} |u|^{q(x)-2} u v dx + \int_{\Omega} f(x, u) v dx, \end{aligned}$$

for all  $v \in W_0^{1,m(x)}(\Omega)$ .

Our main result is the following theorem.

**Theorem 3.2.** Assume that  $(a_1)$ – $(a_2)$ ,  $(f_1)$ – $(f_2)$  and  $(mq_1)$ – $(mq_2)$  hold. Then, there exists a sequence  $\{\lambda_k\} \subset (0, +\infty)$  with  $\lambda_k > \lambda_{k+1}$ , such that for any  $\lambda \in (\lambda_{k+1}, \lambda_k]$ , problem  $(P_\lambda)$  has at least  $k$  pairs of nontrivial solutions.

### 4 Proof of main result

We will start by recalling an important abstract theorem involving genus theory, which will be used in the proof of Theorem 3.2.

**Theorem 4.1** ([22]). *Let  $E$  be an infinite dimensional Banach space with  $E = V \oplus X$ , where  $V$  is finite dimensional and let  $I \in C^1(E, \mathbb{R})$  be a even function with  $I(0) = 0$  and satisfying*

- (i) *There are constants  $\beta, \rho > 0$  such that  $I(u) \geq \beta$  for all  $u \in \partial B_\rho \cap X$ ;*
- (ii) *There is  $\tau > 0$  such that  $I$  satisfies the  $(PS)_c$  condition, for  $0 < c < \tau$ ;*
- (iii) *For each finite dimensional subspace  $\tilde{E} \subset E$ , there is  $R = R(\tilde{E}) > 0$  such that  $I(u) \leq 0$  for all  $u \in \tilde{E} \setminus B_R(0)$ .*

*Suppose that  $V$  is  $k$  dimensional and  $V = \text{span}\{e_1, \dots, e_k\}$ . For  $n \geq k$ , inductively choose  $e_{n+1} \notin E_n := \text{span}\{e_1, \dots, e_n\}$ . Let  $R_n = R(E_n)$  and  $D_n = B_{R_n} \cap E_n$ . Define*

$$G_n = \{h \in C(D_n, E) : h \text{ is odd and } h(u) = u, \forall u \in \partial B_{R_n} \cap E_n\}$$

and

$$\Gamma_j = \left\{ h \left( \overline{D_n \setminus Y} \right) : h \in G_n, n \geq j, Y \in \Sigma, \text{ and } \gamma(Y) \leq n - j \right\},$$

where

$$\Sigma = \{Y \subset E \setminus \{0\} : Y \text{ is closed in } E \text{ and } Y = -Y\}$$

and  $\gamma(Y)$  is the genus of  $Y \in \Sigma$ . For each  $j \in \mathbb{N}$ , let

$$c_j = \inf_{K \in \Gamma_j} \max_{u \in K} I(u).$$

Then  $0 < \beta \leq c_j \leq c_{j+1}$  for  $j > k$ , and if  $j > k$  and  $c_j < \tau$ , we have that  $c_j$  is the critical value of  $I$ . Moreover, if  $c_j = c_{j+1} = \dots = c_{j+l} = c < \tau$  for  $j > k$ , then  $\gamma(K_c) \geq l + 1$ , where

$$K_c = \{u \in E : I(u) = c \text{ and } I'(u) = 0\}.$$

In the sequel, we derive some results related to the above theorem and the Palais–Smale compactness condition.

Since we will rely on the critical point theory, we define the energy functional corresponding to problem  $(P_\lambda)$  as  $I_\lambda : W_0^{1, m(x)}(\Omega) \mapsto \mathbb{R}$ ,

$$I_\lambda(u) = \int_\Omega \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} \right) dx - \int_\Omega a(x) \frac{|u|^{m(x)}}{m(x)} - \lambda \int_\Omega \frac{|u|^{q(x)}}{q(x)} dx - \int_\Omega F(x, u) dx.$$

Clearly,  $I_\lambda$  is  $C^1$  functional and the critical points of it are weak solutions of problem  $(P_\lambda)$ .

**Lemma 4.2.** *Assume that  $(a_1)$ ,  $(f_1)$  and  $(mq_1)$  hold. Then for each  $\lambda > 0$ ,  $I_\lambda$  satisfies condition (i) given in Theorem 4.1.*

*Proof.* Let  $\delta > 0$ . By  $(a_1)$ , we have

$$\begin{aligned}
& \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} - a(x) \frac{|u|^{m(x)}}{m(x)} \right) dx \\
&= \frac{1}{1+\delta} \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} - a(x) \frac{|u|^{m(x)}}{m(x)} \right) dx \\
&\quad + \frac{\delta}{1+\delta} \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} - a(x) \frac{|u|^{m(x)}}{m(x)} \right) dx \\
&= \frac{1}{1+\delta} \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} - a(x) \frac{|u|^{m(x)}}{m(x)} \right) dx \\
&\quad + \frac{\delta}{1+\delta} \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} \right) dx - \frac{\delta}{1+\delta} \int_{\Omega} a(x) \frac{|u|^{m(x)}}{m(x)} dx \\
&\geq \frac{\alpha}{1+\delta} \int_{\Omega} \frac{|u|^{m(x)}}{m(x)} dx + \frac{\delta}{1+\delta} \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} \right) dx - \frac{\delta \|a\|_{\infty}}{(1+\delta)m^-} \int_{\Omega} |u|^{m(x)} dx \\
&\geq \frac{\delta}{1+\delta} \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} \right) dx + \frac{1}{1+\delta} \left( \frac{\alpha}{m^+} - \frac{\delta \|a\|_{\infty}}{m^-} \right) \int_{\Omega} |u|^{m(x)} dx.
\end{aligned}$$

We can choose  $\delta > 0$  such that  $C_0 := \frac{1}{1+\delta} \left( \frac{\alpha}{m^+} - \frac{\delta \|a\|_{\infty}}{m^-} \right) > 0$ . So

$$\begin{aligned}
\int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} - a(x) \frac{|u|^{m(x)}}{m(x)} \right) dx &\geq \frac{\delta}{1+\delta} \int_{\Omega} \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} \right) dx \\
&\quad + C_0 \int_{\Omega} |u|^{m(x)} dx.
\end{aligned} \tag{4.1}$$

By the seconde part of  $(f_1)$ , there is  $\eta_{\varepsilon} > 0$  such that

$$|f(x, t)| \leq \varepsilon |t|^{q(x)-1} \text{ for all } |t| \geq \eta_{\varepsilon} \text{ and for all } x \in \bar{\Omega}.$$

Thanks to the continuity of  $f$ , there is  $A_{\varepsilon} > 0$ , such that

$$|f(x, t)| \leq A_{\varepsilon} \text{ for all } |t| \leq \eta_{\varepsilon} \text{ and for all } x \in \bar{\Omega}.$$

Therefore

$$|f(x, t)| \leq \varepsilon |t|^{q(x)-1} + A_{\varepsilon} \text{ for all } (x, t) \in \bar{\Omega} \times \mathbb{R}. \tag{4.2}$$

On the other hand, by the first part of  $(f_1)$ , for each  $\varepsilon > 0$  there exists  $0 < \delta_{\varepsilon} < 1$  such that

$$|f(x, t)| \leq \varepsilon |t|^{m(x)-1} \text{ for all } |t| \leq \delta_{\varepsilon} \text{ and for all } x \in \bar{\Omega}. \tag{4.3}$$

For  $|t| \geq \delta_{\varepsilon}$ , it follows from (4.2) that

$$\frac{|f(x, t)|}{|t|^{q(x)-1}} \leq \varepsilon + \frac{A_{\varepsilon}}{\delta_{\varepsilon}^{q(x)-1}} \leq \varepsilon + \frac{A_{\varepsilon}}{\delta_{\varepsilon}^{q^+-1}} = C_{\varepsilon},$$

that is

$$|f(x, t)| \leq C_{\varepsilon} |t|^{q(x)-1} \text{ for all } |t| \geq \delta_{\varepsilon} \text{ and for all } x \in \bar{\Omega}. \tag{4.4}$$

Combining (4.3)–(4.4), we obtain

$$|f(x, t)| \leq \varepsilon |t|^{m(x)-1} + C_\varepsilon |t|^{q(x)-1} \quad \text{for all } (x, t) \in \overline{\Omega} \times \mathbb{R}.$$

By integrating this last inequality, we get

$$|F(x, t)| \leq \frac{\varepsilon}{m(x)} |t|^{m(x)} + \frac{C_\varepsilon}{q(x)} |t|^{q(x)} \quad \text{for all } (x, t) \in \overline{\Omega} \times \mathbb{R}. \quad (4.5)$$

Therefore

$$\begin{aligned} I_\lambda(u) &\geq \frac{\delta}{1+\delta} \int_\Omega \left( \frac{|\nabla u|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u|^{p_2(x)}}{p_2(x)} \right) dx + \left( C_0 - \frac{\varepsilon}{m^-} \right) \int_\Omega |u|^{m(x)} dx - \frac{\lambda + C_\varepsilon}{q^-} \int_\Omega |u|^{q(x)} dx \\ &\geq \frac{\delta}{(1+\delta) \max(p_1^+, p_2^+)} \int_\Omega \left( |\nabla u|^{p_1(x)} + |\nabla u|^{p_2(x)} \right) dx + \left( C_0 - \frac{\varepsilon}{m^-} \right) \int_\Omega |u|^{m(x)} dx \\ &\quad - \frac{\lambda + C_\varepsilon}{q^-} \int_\Omega |u|^{q(x)} dx. \end{aligned}$$

Hence for  $\varepsilon$  sufficiently small,

$$I_\lambda(u) \geq \frac{\delta}{(1+\delta) \max(p_1^+, p_2^+)} \int_\Omega \left( |\nabla u|^{p_1(x)} + |\nabla u|^{p_2(x)} \right) dx - \frac{\lambda + C_\varepsilon}{q^-} \int_\Omega |u|^{q(x)} dx.$$

Using the fact that

$$|\nabla u|^{p_1(x)} + |\nabla u(x)|^{p_2(x)} \geq |\nabla u(x)|^{m(x)} \quad \text{for all } x \in \Omega, \quad (4.6)$$

it follows that

$$I_\lambda(u) \geq \frac{\delta}{(1+\delta) \max(p_1^+, p_2^+)} \int_\Omega |\nabla u|^{m(x)} dx - \frac{\lambda + C_\varepsilon}{q^-} \int_\Omega |u|^{q(x)} dx.$$

By the continuous embedding  $W_0^{1,m(x)}(\Omega) \hookrightarrow L^{q(x)}(\Omega)$ , there exists  $C_1 > 0$  such that

$$\|u\|_{L^{q(x)}(\Omega)} \leq C_1 \|u\|.$$

Consequently, by Proposition 2.1, for  $\|u\| = \varrho$ , with  $0 < \varrho < 1$ ,

$$I_\lambda(u) \geq \frac{\delta}{(1+\delta) \max(p_1^+, p_2^+)} \|u\|^{m^+} - \frac{(\lambda + C_\varepsilon)C_2}{q^-} \|u\|^{q^-}.$$

Since  $m^+ < q^-$ , there exists  $\beta > 0$  such that  $I_\lambda(u) \geq \beta$  for  $\|u\| = \varrho$ , where  $\varrho$  is chosen sufficiently small.  $\square$

**Lemma 4.3.** *Assume that  $(a_1)$ ,  $(f_1)$  and  $(mq_1)$  hold. Then  $I_\lambda$  satisfies condition (iii) given in Theorem 4.1.*

*Proof.* Let  $E$  to be a finite dimensional subspace of  $W_0^{1,m(x)}(\Omega)$ . By (4.2),

$$|f(x, t)| \leq \varepsilon |t|^{q(x)-1} + A_\varepsilon \quad \text{for all } (x, t) \in \overline{\Omega} \times \mathbb{R}.$$



Integrating this inequality, we entail

$$\begin{aligned} \frac{|F(x, t)|}{|t|^{q(x)}} &\leq \frac{\varepsilon}{q(x)} + \frac{A_\varepsilon}{|t|^{q(x)-1}} \\ &\leq \frac{\varepsilon}{q^-} + \frac{A_\varepsilon}{|t|^{q^- - 1}} \quad \text{as } |t| \rightarrow +\infty, \end{aligned}$$

hence for  $\varepsilon > 0$ , there is  $\delta_\varepsilon > 0$ , such that

$$|F(x, t)| \leq \varepsilon |t|^{q(x)} \quad \text{for all } |t| \geq \delta_\varepsilon \text{ and for all } x \in \overline{\Omega}. \quad (4.7)$$

By continuity, there is  $M_\varepsilon > 0$  such that

$$|F(x, t)| \leq M_\varepsilon \quad \text{for all } |t| \leq \delta_\varepsilon \text{ and for all } x \in \overline{\Omega}. \quad (4.8)$$

This and (4.7) imply that

$$F(x, t) \geq -M_\varepsilon - \varepsilon |t|^{q(x)} \quad \text{for all } (x, t) \in \overline{\Omega} \times \mathbb{R}. \quad (4.9)$$

Thus

$$\begin{aligned} I_\lambda(u) &\leq \frac{1}{\min(p_1^-, p_2^-)} \int_\Omega (|\nabla u|^{p_1(x)} dx + |\nabla u|^{p_2(x)}) dx + \frac{\|a\|_\infty}{m^-} \int_\Omega |u|^{m(x)} dx \\ &\quad + \left( \varepsilon - \frac{\lambda}{q^+} \right) \int_\Omega |u|^{q(x)} dx + M_\varepsilon |\Omega|. \end{aligned}$$

By choosing  $\varepsilon = \frac{\lambda}{2q^+}$ , we obtain

$$\begin{aligned} I_\lambda(u) &\leq \frac{1}{\min(p_1^-, p_2^-)} \int_\Omega (|\nabla u|^{p_1(x)} dx + |\nabla u|^{p_2(x)}) dx + \frac{\|a\|_\infty}{m^-} \int_\Omega |u|^{m(x)} dx \\ &\quad - \frac{\lambda}{2q^+} \int_\Omega |u|^{q(x)} dx + M_\varepsilon |\Omega| \\ &\leq \frac{1}{\min(p_1^-, p_2^-)} \left( 2|\Omega| + 2 \int_\Omega |\nabla u|^{m(x)} dx + \|a\|_\infty \int_\Omega |u|^{m(x)} dx \right) \\ &\quad - \frac{\lambda}{2q^+} \int_\Omega |u|^{q(x)} dx + M_\varepsilon |\Omega|. \end{aligned}$$

Since  $\dim E < \infty$ , the norms  $\|\cdot\|$  and  $\|\cdot\|_{L^{q(x)}(\Omega)}$  are equivalent in  $E$ . According to Proposition 2.1, for  $\min(\|u\|, \|u\|_{L^{m(x)}(\Omega)}, \|u\|_{L^{q(x)}(\Omega)}) > 1$ ,

$$\begin{aligned} I_\lambda(u) &\leq \frac{1}{\min(p_1^-, p_2^-)} \left( 2|\Omega| + 2\|u\|^{m^+} + \|a\|_\infty \|u\|_{L^{m(x)}(\Omega)}^{m^+} \right) - \frac{\lambda}{2q^+} \|u\|_{L^{q(x)}(\Omega)}^{q^-} + M_\varepsilon |\Omega| \\ &\leq \frac{1}{\min(p_1^-, p_2^-)} \left( 2|\Omega| + 2\|u\|^{m^+} + C' \|a\|_\infty \|u\|^{m^+} \right) - \frac{\lambda C_3}{2q^+} \|u\|^{q^-} + M_\varepsilon |\Omega|. \end{aligned}$$

Using the fact that  $m^+ < q^-$ , we conclude that  $I_\lambda(u) < 0$  for  $\|u\| \geq R > 1$ , where  $R$  is chosen large enough.  $\square$

#### 4.1 Palais–Smale condition

**Lemma 4.4.** *Assume that  $(a_1)$ ,  $(f_1)$ – $(f_2)$  and  $(mq_1)$  hold. Then any (PS) sequence of  $I_\lambda$  is bounded in  $W_0^{1,m(x)}(\Omega)$ .*

*Proof.* Let  $\{u_n\} \subset W_0^{1,m(x)}(\Omega)$  such that  $I_\lambda(u_n) \rightarrow c$  and  $I'_\lambda(u_n) \rightarrow 0$ . By  $(f_2)$ , for  $n$  large enough,

$$\begin{aligned}
c + 1 + \|u_n\| &\geq I_\lambda(u_n) - \frac{1}{m^+} \langle I'_\lambda(u_n), u_n \rangle \\
&= \int_\Omega \left( \frac{1}{p_1(x)} - \frac{1}{m^+} \right) |\nabla u_n|^{p_1(x)} dx + \int_\Omega \left( \frac{1}{p_2(x)} - \frac{1}{m^+} \right) |\nabla u_n|^{p_2(x)} dx \\
&\quad + \int_\Omega \left( \frac{1}{m^+} - \frac{1}{m(x)} \right) a(x) |u_n|^{m(x)} dx + \lambda \int_\Omega \left( \frac{1}{m^+} - \frac{1}{q(x)} \right) |u_n|^{q(x)} dx \\
&\quad + \int_\Omega \left( \frac{1}{m^+} f(x, u_n) u_n - F(x, u_n) \right) dx \\
&\geq \int_\Omega \left( \frac{1}{p_1(x)} - \frac{1}{m^+} \right) |\nabla u_n|^{p_1(x)} dx + \int_\Omega \left( \frac{1}{p_2(x)} - \frac{1}{m^+} \right) |\nabla u_n|^{p_2(x)} dx \\
&\quad + \int_\Omega \left( \frac{1}{m^+} - \frac{1}{m(x)} \right) a(x) |u_n|^{m(x)} dx + \lambda \int_\Omega \left( \frac{1}{m^+} - \frac{1}{q(x)} \right) |u_n|^{q(x)} dx.
\end{aligned}$$

Therefore

$$\begin{aligned}
\lambda \left( \frac{1}{m^+} - \frac{1}{q^-} \right) \int_\Omega |u_n|^{q(x)} dx &\leq \lambda \int_\Omega \left( \frac{1}{m^+} - \frac{1}{q(x)} \right) |u_n|^{q(x)} dx \\
&\leq c + 1 + \|u_n\| + \int_\Omega \left( \frac{1}{m(x)} - \frac{1}{m^+} \right) a(x) |u_n|^{m(x)} dx \\
&\leq c + 1 + \|u_n\| + \|a\|_\infty \left( \frac{1}{m^-} - \frac{1}{m^+} \right) \int_\Omega |u_n|^{m(x)} dx.
\end{aligned}$$

On the other hand, by  $(mq_1)$ , for any  $\varepsilon > 0$ , there exists  $C_\varepsilon > 0$  such that

$$|t|^{m(x)} \leq \varepsilon |t|^{q(x)} + C_\varepsilon \quad \text{for all } (x, t) \in \Omega \times \mathbb{R}.$$

It follows that

$$\begin{aligned}
\lambda \left( \frac{1}{m^+} - \frac{1}{q^-} \right) \int_\Omega |u_n|^{q(x)} dx &\leq c + 1 + \|u_n\| + \varepsilon \|a\|_\infty \left( \frac{1}{m^-} - \frac{1}{m^+} \right) \int_\Omega |u_n|^{q(x)} dx \\
&\quad + \|a\|_\infty \left( \frac{1}{m^-} - \frac{1}{m^+} \right) C_\varepsilon |\Omega|,
\end{aligned}$$

hence

$$\begin{aligned}
&\left( \lambda \left( \frac{1}{m^+} - \frac{1}{q^-} \right) - \varepsilon \|a\|_\infty \left( \frac{1}{m^-} - \frac{1}{m^+} \right) \right) \int_\Omega |u_n|^{q(x)} dx \\
&\leq c + 1 + \|u_n\| + \|a\|_\infty \left( \frac{1}{m^-} - \frac{1}{m^+} \right) C_\varepsilon |\Omega|.
\end{aligned}$$

Choosing  $\varepsilon = \frac{\lambda}{2\|a\|_\infty} \left( \frac{1}{m^+} - \frac{1}{q^-} \right) / \left( \frac{1}{m^-} - \frac{1}{m^+} \right)$ , we obtain

$$\frac{\lambda}{2} \left( \frac{1}{m^+} - \frac{1}{q^-} \right) \int_\Omega |u_n|^{q(x)} dx \leq c + 1 + \|u_n\| + \|a\|_\infty \left( \frac{1}{m^-} - \frac{1}{m^+} \right) C_\varepsilon |\Omega|.$$

Thus,

$$\int_{\Omega} |u_n|^{q(x)} dx \leq C_4(1 + \|u_n\|). \quad (4.10)$$

By (4.7) and (4.8), for  $\varepsilon > 0$ , there exists  $C_\varepsilon > 0$  such that

$$|F(x, t)| \leq \varepsilon |t|^{q(x)} + C_\varepsilon \quad \text{for all } (x, t) \in \bar{\Omega} \times \mathbb{R}. \quad (4.11)$$

(4.1), (4.6), (4.10) and (4.11) imply that for  $n$  large enough

$$\begin{aligned} \frac{\delta}{(1 + \delta) \max(p_1^+, p_2^+)} \int_{\Omega} |\nabla u_n|^{m(x)} dx &\leq \frac{\delta}{1 + \delta} \int_{\Omega} \left( \frac{|\nabla u_n|^{p_1(x)}}{p_1(x)} + \frac{|\nabla u_n|^{p_2(x)}}{p_2(x)} \right) dx \\ &\leq I_\lambda(u_n) + \frac{\lambda}{q^-} \int_{\Omega} |u_n|^{q(x)} dx + \int_{\Omega} F(x, u_n) dx \\ &\leq C_5 + \left( \frac{\lambda}{q^-} + \varepsilon \right) \int_{\Omega} |u_n|^{q(x)} dx + C_\varepsilon |\Omega| \\ &\leq \left( \frac{\lambda}{q^-} + \varepsilon \right) C_4(1 + \|u_n\|) + C_5 + C_\varepsilon |\Omega|. \end{aligned}$$

Hence

$$\int_{\Omega} |\nabla u_n|^{m(x)} dx \leq C_6(1 + \|u_n\|)$$

and so

$$\min(\|u_n\|^{m^-}, \|u_n\|^{m^+}) \leq C(1 + \|u_n\|).$$

Consequently  $\{u_n\}$  is bounded in  $W_0^{1, m(x)}(\Omega)$ .  $\square$

In view of Lemma 4.4,  $\{u_n\}$  is bounded in  $W_0^{1, m(x)}(\Omega)$ . So, up to subsequence, we may assume that

$$\begin{aligned} u_n &\rightharpoonup u \quad \text{in } W_0^{1, m(x)}(\Omega), \\ u_n &\rightharpoonup u \quad \text{in } L^{q(x)}(\Omega), \\ u_n &\rightarrow u \quad \text{in } L^{r(x)}(\Omega), \quad r \in C_+(\bar{\Omega}), \quad r(x) < m^*(x) \quad \forall x \in \bar{\Omega}. \end{aligned}$$

Taking in to account  $(mq_2)$ , from the concentration compactness lemma of Lions [8], there exist two nonnegative measures  $\mu, \nu \in \mathcal{M}(\Omega)$ , a countable set  $\mathcal{J}$ , points  $\{x_j\}_{j \in \mathcal{J}}$  in  $\Omega$  and sequences  $\{\mu_j\}_{j \in \mathcal{J}}, \{\nu_j\}_{j \in \mathcal{J}} \subset [0, +\infty)$ , such that

$$\begin{aligned} |\nabla u_n|^{m(x)} &\rightharpoonup \mu \geq |\nabla u|^{m(x)} + \sum_{j \in \mathcal{J}} \mu_j \delta_{x_j} \quad \text{in } \Omega \\ |u_n|^{q(x)} &\rightharpoonup \nu \geq |u|^{q(x)} + \sum_{j \in \mathcal{J}} \nu_j \delta_{x_j} \quad \text{in } \Omega \\ S \nu_j^{\frac{1}{m^*(x_j)}} &\leq \mu_j^{\frac{1}{m(x_j)}} \quad \text{for all } j \in \mathcal{J}, \end{aligned} \quad (4.12)$$

where

$$S = \inf_{\phi \in C_0^\infty(\Omega)} \frac{\|\nabla \phi\|_{L^{m(x)}(\Omega)}}{\|\phi\|_{L^{q(x)}(\Omega)}}.$$

Let  $\phi \in C_0^\infty(\mathbb{R}^N)$  such that

$$0 \leq \phi \leq 1, \quad \phi \equiv 1 \quad \text{in } B_1(0), \quad \phi = 0 \quad \text{in } \mathbb{R}^N \setminus B_2(0).$$

For  $\varepsilon > 0$  and  $j \in \mathcal{J}$  denote

$$\phi_\varepsilon^j(x) = \phi\left(\frac{x - x_j}{\varepsilon}\right), \quad \text{for all } x \in \mathbb{R}^N.$$

We claim that

$$\begin{aligned} & \int_{\Omega} \left( |u \nabla \phi_\varepsilon^j|^{p_1(x)} + |u \nabla \phi_\varepsilon^j|^{p_2(x)} \right) dx \\ & \leq C \left( |B_{2\varepsilon}(x_j)| + \|u\|_{L^{m^*(x)}(B_{2\varepsilon}(x_j))}^{m^+} + \|u\|_{L^{m^*(x)}(B_{2\varepsilon}(x_j))}^{m^-} \right), \end{aligned} \quad (4.13)$$

for all  $u \in W_0^{1,m(x)}(\Omega)$ , where  $C > 0$  is independent of  $\varepsilon$  and  $j \in \mathcal{J}$ . Indeed,

$$\begin{aligned} & \int_{\Omega} \left( |u \nabla \phi_\varepsilon^j|^{p_1(x)} + |u \nabla \phi_\varepsilon^j|^{p_2(x)} \right) dx \\ & \leq 2|B_{2\varepsilon}(x_j)| + 2 \int_{|u \nabla \phi_\varepsilon^j| \geq 1} \left( |u \nabla \phi_\varepsilon^j|^{m(x)} \right) dx \\ & \leq 2|B_{2\varepsilon}(x_j)| + 2 \int_{\Omega} \left( |u \nabla \phi_\varepsilon^j|^{m(x)} \right) dx \\ & \leq C_7 \left( |B_{2\varepsilon}(x_j)| + \int_{\Omega} |u \nabla \phi_\varepsilon^j|^{m(x)} dx \right) \\ & = C_7 \left( |B_{2\varepsilon}(x_j)| + \int_{B_{2\varepsilon}(x_j)} |u|^{m(x)} \left| \frac{1}{\varepsilon} \nabla \phi \left( \frac{x - x_j}{\varepsilon} \right) \right|^{m(x)} dx \right). \end{aligned} \quad (4.14)$$

Using Hölder's inequality,

$$\begin{aligned} & \int_{B_{2\varepsilon}(x_j)} |u|^{m(x)} \left| \frac{1}{\varepsilon} \nabla \phi \left( \frac{x - x_j}{\varepsilon} \right) \right|^{m(x)} dx \\ & \leq C_8 \left\| |u|^{m(x)} \right\|_{L^{\frac{m^*(x)}{m(x)}}(B_{2\varepsilon}(x_j))} \left\| \left| \frac{1}{\varepsilon} \nabla \phi \left( \frac{x - x_j}{\varepsilon} \right) \right|^{m(x)} \right\|_{L^{\frac{m^*(x)}{m^*(x)-m(x)}}(B_{2\varepsilon}(x_j))}. \end{aligned} \quad (4.15)$$

Furthermore,

$$\begin{aligned} & \int_{B_{2\varepsilon}(x_j)} \left| \frac{1}{\varepsilon} \nabla \phi \left( \frac{x - x_j}{\varepsilon} \right) \right|^{\frac{m(x)m^*(x)}{m^*(x)-m(x)}} dx = \int_{B_{2\varepsilon}(x_j)} \left| \frac{1}{\varepsilon} \nabla \phi \left( \frac{x - x_j}{\varepsilon} \right) \right|^N dx \\ & = \int_{B_2(0)} |\nabla \phi(y)|^N dy = C_9. \end{aligned} \quad (4.16)$$

Gathering (4.14)–(4.16) and taking into account Proposition 2.1, we deduce

$$\begin{aligned} & \int_{\Omega} \left( |u \nabla \phi_\varepsilon^j|^{p_1(x)} + |u \nabla \phi_\varepsilon^j|^{p_2(x)} \right) dx \leq C_7 \left( |B_{2\varepsilon}(x_j)| + C_{10} \left\| |u|^{m(x)} \right\|_{L^{\frac{m^*(x)}{m(x)}}(B_{2\varepsilon}(x_j))} \right) \\ & \leq C \left( |B_{2\varepsilon}(x_j)| + \|u\|_{L^{m^*(x)}(B_{2\varepsilon}(x_j))}^{m^+} + \|u\|_{L^{m^*(x)}(B_{2\varepsilon}(x_j))}^{m^-} \right), \end{aligned}$$

and the claim follows.

**Lemma 4.5.** *Assume that  $(a_1)$ ,  $(f_1)$ – $(f_2)$  and  $mq_1$  hold. If  $\{u_n\}$  is a (PS) sequence of  $I_\lambda$ , then for each  $j \in \mathcal{J}$ ,*

$$v_j = 0 \quad \text{or} \quad v_j \geq \frac{S^N}{\lambda^{\frac{N}{m(x_j)}}}.$$

*Proof.* Let  $\phi_\varepsilon^j$  as above. By Lemma 4.4, we see that for each  $j \in \mathcal{J}$ ,  $\{u_n \phi_\varepsilon^j\}$  is bounded in  $W_0^{1,m(x)}(\Omega)$ . Since  $I'_\lambda(u_n) \rightarrow 0$ ,  $\langle I'_\lambda(u_n), u_n \phi_\varepsilon^j \rangle = o_n(1)$ . From (4.6),

$$\begin{aligned} \int_\Omega |\nabla u_n|^{m(x)} \phi_\varepsilon^j dx &\leq \int_\Omega \left( |\nabla u_n|^{p_1(x)} + |\nabla u_n|^{p_2(x)} \right) \phi_\varepsilon^j dx \\ &= - \int_\Omega |\nabla u_n|^{p_1(x)-2} u_n \nabla u_n \nabla \phi_\varepsilon^j dx - \int_\Omega |\nabla u_n|^{p_2(x)-2} u_n \nabla u_n \nabla \phi_\varepsilon^j dx \\ &\quad + \lambda \int_\Omega |u_n|^{q(x)} \phi_\varepsilon^j dx + \int_\Omega a(x) |u_n|^{m(x)} \phi_\varepsilon^j dx \\ &\quad + \int_\Omega f(x, u_n) u_n \phi_\varepsilon^j dx + o_n(1). \end{aligned} \quad (4.17)$$

For any  $\tau > 0$ , by Young's inequality, there exist  $C_\tau, C'_\tau > 0$  such that

$$\begin{aligned} \int_\Omega \left| |\nabla u_n|^{p_1(x)-2} u_n \nabla u_n \nabla \phi_\varepsilon^j \right| dx &\leq \tau \int_\Omega |\nabla u_n|^{p_1(x)} dx + C_\tau \int_\Omega |u_n \nabla \phi_\varepsilon^j|^{p_1(x)} dx, \\ &\leq \tau \left( |B_{2\varepsilon}(x_j)| + \int_\Omega |\nabla u_n|^{m(x)} dx \right) + C_\tau \int_\Omega |u_n \nabla \phi_\varepsilon^j|^{p_1(x)} dx \end{aligned}$$

and

$$\begin{aligned} \int_\Omega \left| |\nabla u_n|^{p_2(x)-2} u_n \nabla u_n \nabla \phi_\varepsilon^j \right| dx &\leq \tau \int_\Omega |\nabla u_n|^{p_2(x)} dx + C'_\tau \int_\Omega |u_n \nabla \phi_\varepsilon^j|^{p_2(x)} dx \\ &\leq \tau \left( |B_{2\varepsilon}(x_j)| + \int_\Omega |\nabla u_n|^{m(x)} dx \right) + C'_\tau \int_\Omega |u_n \nabla \phi_\varepsilon^j|^{p_2(x)} dx. \end{aligned}$$

Using the fact  $\{u_n\}$  is bounded in  $W_0^{1,m(x)}(\Omega)$ , it follows that

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \int_\Omega \left| |\nabla u_n|^{p_1(x)-2} u_n \nabla u_n \nabla \phi_\varepsilon^j \right| dx &\leq \tau C_{11} + C_\tau \int_\Omega |u \nabla \phi_\varepsilon^j|^{p_1(x)} dx \\ \limsup_{n \rightarrow +\infty} \int_\Omega \left| |\nabla u_n|^{p_2(x)-2} u_n \nabla u_n \nabla \phi_\varepsilon^j \right| dx &\leq \tau C_{11} + C'_\tau \int_\Omega |u \nabla \phi_\varepsilon^j|^{p_2(x)} dx. \end{aligned}$$

Now, bearing in mind (4.13), we obtain

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \left[ \int_\Omega \left| |\nabla u_n|^{p_1(x)-2} u_n \nabla u_n \nabla \phi_\varepsilon^j \right| dx + \int_\Omega \left| |\nabla u_n|^{p_2(x)-2} u_n \nabla u_n \nabla \phi_\varepsilon^j \right| dx \right] \\ \leq \tau C_{12} + \max(C_\tau, C'_\tau) \left( \int_\Omega |u \nabla \phi_\varepsilon^j|^{p_1(x)} dx + \int_\Omega |u \nabla \phi_\varepsilon^j|^{p_2(x)} dx \right) \\ \leq \tau C_{12} + \max(C_\tau, C'_\tau) C \left( |B_{2\varepsilon}(x_j)| + \|u\|_{L^{m^*(x)}(B_{2\varepsilon}(x_j))}^{m^+} + \|u\|_{L^{m^*(x)}(B_{2\varepsilon}(x_j))}^{m^-} \right) \end{aligned} \quad (4.18)$$

On the other hand, by the compactness lemma of Strauss [10] and Sobolev embedding,

$$\lim_{n \rightarrow +\infty} \int_\Omega f(x, u_n) u_n \phi_\varepsilon^j dx = \int_\Omega f(x, u) u \phi_\varepsilon^j dx \quad (4.19)$$

and

$$\lim_{n \rightarrow +\infty} \int_\Omega a(x) |u_n|^{m(x)} \phi_\varepsilon^j dx = \int_\Omega a(x) |u|^{m(x)} \phi_\varepsilon^j dx. \quad (4.20)$$

From (4.12) and (4.17)–(4.20),

$$\begin{aligned} \int_\Omega \phi_\varepsilon^j d\mu &\leq \tau C_{12} + \max(C_\tau, C'_\tau) C \left( |B_{2\varepsilon}(x_j)| + \|u\|_{L^{m^*(x)}(B_{2\varepsilon}(x_j))}^{m^+} + \|u\|_{L^{m^*(x)}(B_{2\varepsilon}(x_j))}^{m^-} \right) \\ &\quad + \lambda \int_\Omega \phi_\varepsilon^j dv - \int_{B_{2\varepsilon}(x_j)} a(x) |u|^{m(x)} \phi_\varepsilon^j dx \\ &\quad + \int_{B_{2\varepsilon}(x_j)} f(x, u) u \phi_\varepsilon^j dx. \end{aligned}$$

Letting  $\varepsilon \rightarrow 0$ , we obtain

$$\mu_j \leq \tau C_{12} + \lambda v_j,$$

Now, letting  $\tau \rightarrow 0$ , we derive  $\mu_j \leq \lambda v_j$ . Therefore

$$S v_j^{\frac{1}{m^*(x_j)}} \leq \mu_j^{\frac{1}{m(x_j)}} \leq (\lambda v_j)^{\frac{1}{m(x_j)}}$$

and hence the result follows.  $\square$

**Lemma 4.6.** *Assume that  $(a_1)$ – $(a_2)$ ,  $(f_1)$ – $(f_2)$  and  $(mq_1)$  hold. If  $\lambda < 1$ , then  $I_\lambda$  satisfies  $(PS)_c$  condition for  $c < \lambda^{1-\frac{N}{m^+}} \left( \frac{1}{m^+} - \frac{1}{q^-} \right) S^N$ .*

*Proof.* Let  $\{u_n\} \subset W_0^{1,m(x)}(\Omega)$  such that

$$I_\lambda(u_n) \rightarrow c \quad \text{and} \quad I'_\lambda(u_n) \rightarrow 0.$$

Then

$$\begin{aligned} c &= I_\lambda(u_n) - \frac{1}{m^+} \langle I'_\lambda(u_n), u_n \rangle + o_n(1) \\ &= \int_\Omega \left( \frac{1}{p_1(x)} - \frac{1}{m^+} \right) |\nabla u_n|^{p_1(x)} dx + \int_\Omega \left( \frac{1}{p_2(x)} - \frac{1}{m^+} \right) |\nabla u_n|^{p_2(x)} dx \\ &\quad + \int_\Omega \left( \frac{1}{m^+} - \frac{1}{m(x)} \right) a(x) |u_n|^{m(x)} dx + \lambda \int_\Omega \left( \frac{1}{m^+} - \frac{1}{q(x)} \right) |u_n|^{q(x)} dx \\ &\quad + \int_\Omega \left( \frac{1}{m^+} f(x, u_n) u_n - F(x, u_n) \right) dx + o_n(1). \end{aligned}$$

By  $(a_2)$  and  $(f_2)$ , we obtain

$$c \geq \lambda \int_\Omega \left( \frac{1}{m^+} - \frac{1}{q^-} \right) |u_n|^{q(x)} dx + o_n(1).$$

Using (4.12), it follows that

$$\begin{aligned} c &\geq \lambda \lim_{n \rightarrow +\infty} \int_\Omega \left( \frac{1}{m^+} - \frac{1}{q^-} \right) |u_n|^{q(x)} dx \\ &\geq \lambda \left( \frac{1}{m^+} - \frac{1}{q^-} \right) \left( \int_\Omega |u|^{q(x)} dx + \sum_{j \in \mathcal{J}} v_j \right) \\ &\geq \lambda \left( \frac{1}{m^+} - \frac{1}{q^-} \right) v_j \quad \text{for all } j \in \mathcal{J}. \end{aligned}$$

If  $v_j > 0$  for some  $j \in \mathcal{J}$ , by Lemma 4.5, we get

$$c \geq \lambda \left( \frac{1}{m^+} - \frac{1}{q^-} \right) \frac{S^N}{\lambda^{\frac{N}{m^+}}}.$$

Since  $\lambda < 1$ ,

$$c \geq \lambda \left( \frac{1}{m^+} - \frac{1}{q^-} \right) \frac{S^N}{\lambda^{\frac{N}{m^+}}} = \lambda^{1-\frac{N}{m^+}} \left( \frac{1}{m^+} - \frac{1}{q^-} \right) S^N,$$

which is impossible, and so  $v_j = 0$  for all  $j \in \mathcal{J}$ . Hence

$$\lim_{n \rightarrow +\infty} \int_{\Omega} |u_n|^{q(x)} dx = \int_{\Omega} |u|^{q(x)} dx.$$

This implies  $\lim_{n \rightarrow +\infty} \int_{\Omega} |u_n - u|^{q(x)} dx = 0$ , thanks to Proposition 2.1, we deduce

$$u_n \rightarrow u \text{ in } L^{q(x)}(\Omega).$$

By standard arguments, we see that

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \left( |u_n|^{q(x)-2} u_n - |u|^{q(x)-2} u \right) (u_n - u) dx = 0,$$

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \left( |u_n|^{m(x)-2} u_n - |u|^{m(x)-2} u \right) (u_n - u) dx = 0$$

and

$$\lim_{n \rightarrow +\infty} \int_{\Omega} (f(x, u_n) - f(x, u))(u_n - u) dx = 0.$$

On the other hand, we have

$$\begin{aligned} o_n(1) &= \langle I'_\lambda(u_n) - I'_\lambda(u), u_n - u \rangle \\ &= \int_{\Omega} \left( |\nabla u_n|^{p_1(x)-2} \nabla u_n - |\nabla u|^{p_1(x)-2} \nabla u \right) \nabla (u_n - u) dx \\ &\quad + \int_{\Omega} \left( |\nabla u_n|^{p_1(x)-2} \nabla u_n - |\nabla u|^{p_2(x)-2} \nabla u \right) \nabla (u_n - u) dx \\ &\quad - \int_{\Omega} a(x) \left( |u_n|^{m(x)-2} u_n - |u|^{m(x)-2} u \right) (u_n - u) dx \\ &\quad - \lambda \int_{\Omega} \left( |u_n|^{q(x)-2} u_n - |u|^{q(x)-2} u \right) (u_n - u) dx - \int_{\Omega} (f(x, u_n) - f(x, u))(u_n - u) dx. \end{aligned}$$

It follows that

$$\begin{aligned} \lim_{n \rightarrow +\infty} \left[ \int_{\Omega} \left( |\nabla u_n|^{p_1(x)-2} \nabla u_n - |\nabla u|^{p_1(x)-2} \nabla u \right) \nabla (u_n - u) dx \right. \\ \left. + \int_{\Omega} \left( |\nabla u_n|^{p_1(x)-2} \nabla u_n - |\nabla u|^{p_2(x)-2} \nabla u \right) \nabla (u_n - u) dx \right] = 0. \end{aligned}$$

Since  $(|\nabla u_n|^{p_i(x)-2} \nabla u_n - |\nabla u|^{p_i(x)-2} \nabla u) \nabla (u_n - u) \geq 0$  in  $\Omega$ ,  $i = 1, 2$ ,

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \left( |\nabla u_n|^{p_i(x)-2} \nabla u_n - |\nabla u|^{p_i(x)-2} \nabla u \right) \nabla (u_n - u) dx = 0, \quad i = 1, 2. \quad (4.21)$$

Recalling the following well known inequality in  $\mathbb{R}^N$ ,

$$\left[ (|\xi|^{p-2} \xi - |\eta|^{p-2} \eta)(\xi - \eta) \right]^{\frac{p}{2}} (|\xi|^p + |\eta|^p)^{\frac{2-p}{2}} \geq (p-1) |\xi - \eta|^p \quad \text{if } 1 < p < 2, \quad (4.22)$$

$$(|\xi|^{p-2} \xi - |\eta|^{p-2} \eta)(\xi - \eta) \geq 2^{-p} |\xi - \eta|^p \quad \text{if } p \geq 2. \quad (4.23)$$

Divide  $\Omega$  in two parts as follows:

$$\Omega_i^+ = \{x \in \Omega : p_i(x) \geq 2\} \quad \text{and} \quad \Omega_i^- = \{x \in \Omega : p_i(x) < 2\}, \quad i = 1, 2.$$

By (4.21) and (4.23), it yields

$$\lim_{n \rightarrow +\infty} \int_{\Omega_i^+} |\nabla (u_n - u)|^{p_i(x)} dx = 0, \quad i = 1, 2. \quad (4.24)$$

On the other hand, by (4.22) and Hölder's inequality,

$$\begin{aligned}
\int_{\Omega_i^-} |\nabla(u_n - u)|^{p_i(x)} dx &\leq \frac{1}{p_i^- - 1} \int_{\Omega_i^-} (p_i(x) - 1) |\nabla u_n - u|^{p_i(x)} dx \\
&\leq \frac{1}{p_i^- - 1} \int_{\Omega_i^-} \left[ \left( |\nabla u_n|^{p_i(x)-2} \nabla u_n - |\nabla u|^{p_i(x)-2} \nabla u \right) \nabla(u_n - u) \right]^{\frac{p_i(x)}{2}} \\
&\quad \times \left( |\nabla u_n|^{p_i(x)} + |\nabla u|^{p_i(x)} \right)^{\frac{2-p_i(x)}{2}} dx \\
&\leq C_{p_i} \left\| \left[ \left( |\nabla u_n|^{p_i(x)-2} \nabla u_n - |\nabla u|^{p_i(x)-2} \nabla u \right) \nabla(u_n - u) \right]^{\frac{p_i(x)}{2}} \right\|_{L^{\frac{2}{p_i(x)}}(\Omega_i^-)} \\
&\quad \times \left\| \left( |\nabla u_n|^{p_i(x)} + |\nabla u|^{p_i(x)} \right)^{\frac{2-p_i(x)}{2}} \right\|_{L^{\frac{2}{2-p_i(x)}}(\Omega_i^-)}.
\end{aligned}$$

Since  $\{u_n\}$  is bounded in  $W_0^{1,p_i(x)}(\Omega)$ , Proposition 2.1 implies

$$\int_{\Omega_i^-} |\nabla(u_n - u)|^{p_i(x)} dx \leq C'_{p_i} \left\| \left[ \left( |\nabla u_n|^{p_i(x)-2} \nabla u_n - |\nabla u|^{p_i(x)-2} \nabla u \right) \nabla(u_n - u) \right]^{\frac{p_i(x)}{2}} \right\|_{L^{\frac{2}{p_i(x)}}(\Omega_i^-)}.$$

Using again Proposition 2.1 and (4.21), it follows that

$$\lim_{n \rightarrow +\infty} \int_{\Omega_i^-} |\nabla(u_n - u)|^{p_i(x)} dx = 0, \quad i = 1, 2. \quad (4.25)$$

(4.24) and (4.25) imply

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \left( |\nabla(u_n - u)|^{p_1(x)} + |\nabla(u_n - u)|^{p_2(x)} \right) dx = 0.$$

Using inequality (4.6), we get

$$\int_{\Omega} |\nabla(u_n - u)|^{m(x)} dx = 0,$$

and hence  $u_n \rightarrow u$  in  $W_0^{1,m(x)}(\Omega)$ .  $\square$

**Lemma 4.7.** *Under assumptions of Theorem 3.2, there exists a sequence  $\{M_n\} \subset (0, +\infty)$  independent of  $\lambda$ , with  $M_n \leq M_{n+1}$ , such that for any  $\lambda > 0$ ,*

$$c_n^\lambda = \inf_{K \in \Gamma_n} \max_{u \in K} I_\lambda(u) < M_n.$$

*Proof.* The proof is similar to [26, Lemma 5] and so, we will omit it.  $\square$

## 4.2 Proof of Theorem 3.2

By choosing for each  $k \geq 1$ ,  $\lambda_k$  sufficiently small, we construct a sequence  $(\lambda_k)$ , with  $\lambda_k > \lambda_{k+1}$  such that  $M_k < \lambda_k^{1-\frac{N}{m^+}} \left( \frac{1}{m^+} - \frac{1}{q^-} \right) S^N$ . Therefore, for  $\lambda \in (\lambda_{k+1}, \lambda_k]$ ,

$$0 < c_1^\lambda \leq c_2^\lambda \leq \dots \leq c_k^\lambda < M_k < \lambda^{1-\frac{N}{m^+}} \left( \frac{1}{m^+} - \frac{1}{q^-} \right) S^N.$$



Thanks to Theorem 4.1, the levels  $c_1^\lambda \leq c_2^\lambda \leq \dots \leq c_k^\lambda$  are critical values of  $I_\lambda$ . So, if

$$c_1^\lambda < c_2^\lambda < \dots < c_k^\lambda,$$

$I_\lambda$  has at least  $k$  critical points. Now, if  $c_j^\lambda = c_{j+1}^\lambda$  for some  $j = 1, \dots, k-1$ , again Theorem 4.1 implies that  $K_{c_j^\lambda}$  is an infinite set [22, Chap. 7] and hence in this case, problem  $(P_\lambda)$  has infinitely many solutions. Conclusion, problem  $(P_\lambda)$  has at least  $k$  pair solutions.

## Acknowledgements

The author would like to thank the referee and the editor for their valuable comments and suggestions which have led to an improvement of the presentation of this paper.

## References

- [1] E. ACERBI, G. MINGIONE, Regularity results for a class of functionals with non-standard growth, *Arch. Ration. Mech. Anal.* **156**(2001), No. 2, 121–140. [MR1814973](#)
- [2] C. O. ALVES, Existence of solution for a degenerate  $p(x)$ -Laplacian equation in  $\mathbb{R}^N$ , *J. Math. Anal. Appl.* **345**(2008), No. 2, 731–742. [MR2429173](#)
- [3] C. O. ALVES, S. LIU, On superlinear  $p(x)$ -Laplacian equations in  $\mathbb{R}^N$ , *Nonlinear Anal.* **73**(2010), No. 8, 2566–2579. [MR2674092](#)
- [4] C. O. ALVES, J. P. BARREIRO, Existence and multiplicity of solutions for a  $p(x)$ -Laplacian equation with critical growth, *J. Math. Anal. Appl.* **403**(2013), No. 1, 143–154. [MR3035079](#)
- [5] S. ANTONTSEV, S. SHMAREV, Elliptic equations with anisotropic nonlinearity and nonstandard growth conditions, in: *Handbook of differential equations. Stationary partial differential equations*, Vol. III, Chap. 1, North Holland, 2006. [url](#)
- [6] M. BOCEA, M. MIHĂILESCU,  $\Gamma$ -convergence of power-law functionals with variable exponents, *Nonlinear Anal.* **73**(2010), No. 1, 110–121. [MR2645836](#)
- [7] M. BOCEA, M. MIHĂILESCU, C. POPOVICI, On the asymptotic behavior of variable exponent power-law functionals and applications, *Ric. Mat.* **59**(2010), No. 2, 207–238. [MR2738653](#)
- [8] J. F. BONDER, A. SILVA, Concentration-compactness principle for variable exponent spaces and applications, *Electron. J. Differential Equations* **2010**, No. 141, 1–18. [MR2729462](#)
- [9] Y. CHEN, S. LEVINE, M. RAO, Variable exponent, linear growth functionals in image restoration, *SIAM J. Appl. Math.* **66**(2006), No. 4, 1383–1406. [MR2246061](#)
- [10] J. CHABROWSKI, *Weak convergence methods for semilinear elliptic equations*, World Scientific Publishing Company, 1999. [MR1885519](#); [url](#)
- [11] J. CUADRO, G. LÓPEZ, Unique continuation for solutions of  $p(x)$ -Laplacian equations, *Electron. J. Differential Equations* **2012**, No. 7, 1–12. [MR2889613](#)
- [12] G. DAI, R. MA, Solutions for a  $p(x)$ -Kirchhoff type equation with Neumann boundary data, *Nonlinear Anal. Real World Appl.* **12**(2011), No. 5, 2666–2680. [MR2813212](#)

- [13] L. DIENING, P. HARJULEHTO, P. HÄSTÖ, M. RŮŽIČKA, *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Mathematics, Vol. 2017, Springer-Verlag, Berlin, 2011. [MR2790542](#); [url](#)
- [14] X. FAN, D. ZHAO, On the spaces  $L^{p(x)}(\Omega)$  and  $W^{m,p(x)}(\Omega)$ , *J. Math. Anal. Appl.* **263**(2001), No. 2, 424–446. [MR1866056](#)
- [15] X. FAN, Q. ZHANG, D. ZHAO, Existence of solutions for  $p(x)$ -Laplacian Dirichlet problem, *J. Math. Anal. Appl.* **302**(2005), No. 2, 306–317. [MR2107835](#)
- [16] P. HARJULEHTO, P. HÄSTÖ, ÚT V. LÊ, M. NUORTIO, Overview of differential equations with non-standard growth, *Nonlinear Anal.* **72**(2010), No. 12, 4551–4574. [MR2639204](#)
- [17] E. M. HSSINI, M. MASSAR, N. TSOULI, Existence and multiplicity of solutions for a  $p(x)$ -Kirchhoff type problems, *Bol. Soc. Parana. Mat.* **33**(2015), No. 2, 201–215. [MR3267328](#); [url](#)
- [18] M. MASSAR, M. TALBI, N. TSOULI, Multiple solutions for nonlocal system of  $(p(x), q(x))$ -Kirchhoff type, *Appl. Math. Comput.* **242**(2014), 216–226. [MR3239653](#)
- [19] M. MIHĂILESCU, On a class of nonlinear problems involving a  $p(x)$ -Laplace type operator, *Czechoslovak Math. J.* **58**(2008), No. 1, 155–172. [MR2402532](#)
- [20] M. MIHĂILESCU, V. RĂDULESCU, On a nonhomogeneous quasilinear eigenvalue problem in Sobolev spaces with variable exponent, *Proc. Amer. Math. Soc.* **135**(2007), No. 9, 2929–2937. [MR2317971](#)
- [21] J. MUSIELAK, *Orlicz spaces and modular spaces*, Lecture Notes in Mathematics, Vol. 1034, Springer-Verlag, Berlin, 1983. [MR724434](#); [url](#)
- [22] P. H. RABINOWITZ, *Minimax methods in critical point theory with applications to differential equations*, CBMS Regional Conference Series in Mathematics, Vol. 65, 1986. [MR845785](#); [url](#)
- [23] M. RŮŽIČKA, Flow of shear dependent electrorheological fluids, *C. R. Acad. Sci. Paris Sér. I Math.* **329**(1999), No. 5, 393–398. [MR1710119](#)
- [24] M. RŮŽIČKA, *Electrorheological fluids: modeling and mathematical theory*, Springer-Verlag, Berlin, 2000 [MR1810360](#); [url](#).
- [25] V. V. ZHIKOV, Averaging of functionals in the calculus of variations and elasticity, *Math. USSR IZV.* **29**(1987), 33–66.
- [26] W. ZHIHUI, W. XINMIN, A multiplicity result for quasilinear elliptic equations involving critical Sobolev exponents, *Nonlinear Anal.* **18**(1992), No. 6, 559–567. [MR1154480](#)