

THE COMPLEX OF WORDS AND NAKAOKA STABILITY

MORITZ C. KERZ

(communicated by Charles Weibel)

Abstract

We give a new simple proof of the exactness of the complex of injective words and use it to prove Nakaoka's homology stability for symmetric groups. The methods are generalized to show acyclicity in low degrees for the complex of words in "general position".

Introduction

In this paper we show the vanishing of homology for various complexes of words and give an elementary, self-contained proof of Nakaoka stability (Theorem 2):

$$H_m(\Sigma_{n-1}, \mathbb{Z}) = H_m(\Sigma_n, \mathbb{Z})$$

for $n/2 > m$ where Σ_n denotes the permutation group of n elements. An elementary proof of this fact has not been available in the literature.

In the first section the complex $C_*(m)$ of abelian groups is studied which in degree n is freely generated by injective words of length n . The alphabet consists of m letters. The complex $C_*(m)$ has the only non vanishing homology in degree m (Theorem 1). This is a result of F.D. Farmer [3] who connected it to properties of the associated poset of injective words and its CW-complex.

Then, considering the action of the permutation group on the alphabet, a hyperhomology argument is used to deduce Nakaoka stability.

In the second section – independent from the first – more general complexes of words are shown to have vanishing homology in low degrees. The proof is analogous to Theorem 1.

A very general setting is used but the essential application is to the complex of words consisting of vectors in general position over a finite field. In fact this complex is shown to become exact in some fixed degree and fixed dimension of the vector space if only the base field has enough elements.

This could be of interest for example for Suslin's GL -stability [8] which works up to now only over infinite fields.

1. An elementary proof of Nakaoka stability

In this section we are concerned with the complex of injective words, denoted $C_*(m)$ ($m \in \mathbf{N}$), and its homology. For $n > 0$ define $C_n(m)$ as the abelian group freely generated by the injective words of length n with alphabet $\mathbf{m} = \{1, 2, \dots, m\}$. Injective word means no element of our alphabet may appear twice in it. Set $C_0(m) = \mathbb{Z}$.

In what follows words are written in brackets, e.g. $(1, 2) + (3, 1) \in C_2(3)$.

The differential $d : C_n(m) \rightarrow C_{n-1}(m)$ for $n > 1$ is defined by

$$d(x_1, \dots, x_n) = \sum_{j=1}^n (-1)^{j+1} (x_1, \dots, \hat{x}_j, \dots, x_n),$$

$x_1, \dots, x_n \in \mathbf{m}$. Similarly $d : C_1(m) \rightarrow C_0(m)$ maps (x) to 1.

In the theory of posets and their associated CW-complexes the following theorem has a nice interpretation. The homology of $C_*(m)$ is equal to the simplicial homology of the (shellable) poset of injective words. Shellability reduces the simplicial homology groups to those of a wedge of m -spheres. We refer to [1], [3], [7] for exact definitions and statements.

Our proof of Theorem 1 is new and rather straightforward compared to Farmer's original elementary proof.

Theorem 1. (Farmer [3]) *The homology of $C_*(m)$ vanishes except in degree m .*

We have to introduce some notations which will be used throughout the paper. An element $c \in C_n(m)$ is called a term, if there exists an $N \in \mathbb{Z}$ and $x_1, \dots, x_n \in \{1, \dots, m\}$ with $c = N(x_1, \dots, x_n)$.

As $C_*(m)$ has a canonical basis all our sum decompositions correspond to partitions of the basis. We also speak about the appearance of numbers in an element of $C_n(m)$. For example 2 appears in $(2, 3) + 4(5, 1)$ but 4 does not.

Although our complex $C_*(m)$ has no obvious cup product, we have a partially defined product. If $c \in C_n(m)$, $c' \in C_l(m)$ are terms, $c = N(x_1, \dots, x_n)$, $c' = M(x'_1, \dots, x'_l)$, we define $cc' := NM(x_1, \dots, x_n, x'_1, \dots, x'_l)$ if the numbers $x_1, \dots, x_n, x'_1, \dots, x'_l$ are distinct. This construction extends bilinearly to arbitrary $c \in C_n(m)$, $c' \in C_l(m)$ for which the numbers appearing in both of them are distinct.

There is a Leibniz rule for $c \in C_n(m)$ and $c' \in C_l(m)$ satisfying the latter condition.

$$d(cc') = d(c)c' + (-1)^n c d(c')$$

Proof of Theorem 1. The exactness in degree 0 is clear. For the rest we use induction on m . For the case $m = 2$ we have to check that

$$C_2(2) \longrightarrow C_1(2) \longrightarrow \mathbb{Z}$$

is exact. But $\ker(d : C_1(2) \rightarrow \mathbb{Z})$ is generated by elements of the form

$$(x) - (x') = d(x, x')$$

with $x \neq x'$.

For the induction step we use a straightforward lemma.

Lemma 1. *If we have a number $x \in \{1, \dots, m\}$ which does not appear in a cycle $c \in C_n(m)$, it is a boundary.*

Proof. According to the Leibniz rule $c = c + (x) d(c) = d((x) c)$, since $d(c) = 0$. \square

Given an arbitrary cycle c of degree $n < m$ we have to show that in order to apply the lemma we can eliminate a number from c by adding boundaries. Therefore we will push a fixed number $x \in \{1, \dots, m\}$ to the right until it vanishes from the cycle.

If x appears somewhere in the cycle at the first entry, write

$$c = \sum_j (x) c_j + c'$$

with terms c_j and c' does not have x at the first entry. To each c_j choose a number $x_j \in \{1, \dots, m\}$, $x_j \neq x$ which does not appear in c_j .

$$c - d\left(\sum_j (x_j) (x) c_j\right) = c' + \sum_j (x_j) c_j - (x_j) (x) d c_j$$

Now x does not appear at the first entry anymore.

The next steps until the vanishing of x are similar. Suppose x does not appear at the first $i > 0$ entries of c . Now we can write

$$c = \sum_j s_j (x) c_j + c'$$

with distinct terms c_j , the s_j have length i , x does not appear in s_j and x does not appear at the first $i + 1$ entries of c' . One calculates:

$$0 = d c = \sum_j [(d s_j) (x) c_j + (-1)^i s_j c_j + (-1)^{i+1} s_j (x) d c_j] + d c' .$$

If we forget those terms in the last equation for which x does not appear at the i -th entry, this equation implies $d s_j = 0$ for all j . Since $\text{length}(s_j) = i < m - \text{length}((x) c_j)$, there are by induction s'_j with $d s'_j = s_j$ and such that the following products make sense.

$$z := \sum_j s'_j (x) c_j$$

In the cycle

$$c - d z = c' - \sum_j [(-1)^{i+1} s'_j c_j + (-1)^i s'_j (x) d c_j]$$

x does not appear at the first $i + 1$ entries.

Finally, x vanishes completely from our cycle and we are in the situation where we can apply Lemma 1. \square

Using the corresponding two hyperhomology spectral sequences for the natural action of the symmetric group Σ_m on our complex $C_*(m)$ (cf [2]) one can now obtain

a stability result due to Nakaoka.[†]

Theorem 2. (Nakaoka [5]) $H_m(\Sigma_{n-1}) = H_m(\Sigma_n)$ for $m < n/2$.

Proof. We use induction on n . It is well known for $n = 3$.

$$H_1(\Sigma_2) = H_1(\Sigma_3) = \mathbb{Z}/(2)$$

For $n \geq 4$ define $C'_l(n) := C_{l+1}(n)$ for $l \geq 0$. Then

$$H_m(\Sigma_n, \mathbb{Z}) = H_m(\Sigma_n, C'_*(n))$$

when $m < n - 1$ because of Theorem 1 and a standard spectral sequence argument [2]. The second spectral sequence of the bi-complex gives for $E_{*,*}^1$:

$$\begin{array}{cccccc} & & \cdots & & & \\ H_2(\Sigma_n, C'_0) & H_2(\Sigma_n, C'_1) & \cdots & H_2(\Sigma_n, C'_{n-2}) & H_2(\Sigma_n, C'_{n-1}) & \\ H_1(\Sigma_n, C'_0) & H_1(\Sigma_n, C'_1) & \cdots & H_1(\Sigma_n, C'_{n-2}) & H_1(\Sigma_n, C'_{n-1}) & \\ H_0(\Sigma_n, C'_0) & H_0(\Sigma_n, C'_1) & \cdots & H_0(\Sigma_n, C'_{n-2}) & H_0(\Sigma_n, C'_{n-1}) & \end{array}$$

Using Shapiro's Lemma we get:

$$\begin{array}{cccccc} & & \cdots & & & \\ H_2(\Sigma_{n-1}) & H_2(\Sigma_{n-2}) & \cdots & H_2(\Sigma_1) & 0 & \\ H_1(\Sigma_{n-1}) & H_1(\Sigma_{n-2}) & \cdots & H_1(\Sigma_1) & 0 & \\ H_0(\Sigma_{n-1}) & H_0(\Sigma_{n-2}) & \cdots & H_0(\Sigma_1) & \mathbb{Z} & \end{array}$$

The horizontal arrows can be computed as $0, 1, 0, 1, \dots$, since they are the sums of the signs in d'_1, d'_2, \dots .

We have $E_{i,0}^2 = H_i(\Sigma_{n-1})$ for $i \geq 0$. By our global induction some $d : E_{i,j}^1 \rightarrow E_{i,j-1}^1$ are isomorphism. Especially, $E_{i-1,2}^2 = 0$ for $n/2 > i$ and

$$E_{i,j}^2 = 0 \text{ for } i < \frac{n-j-1}{2}, 0 < j < n-1.$$

For $n/2 > i$ only the first column on the diagonal $(E_{k,l})_{k+l=i}$ survives. In fact $E_{i,0}^\infty = E_{i,0}^2 = H_i(\Sigma_{n-1})$ for $n/2 > i$, because the differentials that could kill these groups come from $E_{i-l,l+1}^{\geq 2} = 0$ for $l > 0$. Finally, $H_i(\Sigma_{n-1}) = H_i(\Sigma_n)$ for $n/2 > i$. □

2. Words in general position

Let $\mathbf{n} = \{i | 1 \leq i \leq n\}$. We associate to every (nonempty) set X a complex $F(X)_*$, the so-called complex of words with alphabet X , as follows:

$$\begin{aligned} (F(X))_n &= \mathbb{Z} \langle \{f : \mathbf{n} \rightarrow X\} \rangle \\ d_n(f) &= \sum_{k=1}^n (-1)^{n+1} f \circ \delta_k. \end{aligned}$$

[†]Hendrik Maazen gives another prove of Nakaoka's stability in his dissertation. I have to thank Wilberd van der Kallen for this remark.

Here $\delta_k : [n] \rightarrow [n + 1]$ are the coface maps

$$\delta_k(j) = \begin{cases} j & \text{if } 1 \leq j < k \\ j + 1 & \text{if } k \leq j \end{cases} .$$

It is immediate that the homology of $F(X)$ vanishes. For if $c \in F(X)_n$ is a cycle, $c = d((x_0)c)$ for arbitrary $x_0 \in X$ (the notational convention is explained below Theorem 1).

For given X certain subcomplexes of $F(X)$ can be used in hyperhomology spectral sequences as above, if their homology vanishes to some extent. These subgroups are determined by conditions which one could call "general position conditions". For applications cf [8], [9].

It could be asked, how to explain the fact that the vanishing of homology is not affected by these conditions. In order to give a general result we translate the proof of Theorem 1 into our more complicated setting.

Examples

- (i) Let X be a finite set. The complex of injective words is $(F_{G^{inj}}(X))_n := \{f \in F(X)_n | f \text{ injective}\}$. According to Theorem 1 the homology of this subcomplex is zero except in degree $m = \text{card}(X)$ where

$$\text{rank}(H_m(F(X)_*)) = (-1)^m \left(1 - \sum_{i=0}^{m-1} (-1)^i m(m-1) \cdots (m-i)\right).$$

This equals the number of fixed point free permutations of X .

- (ii) Let k be a field and V a k -vector space. The complex of vectors in general position is $(F_{G^{vec}}(V))_n := \{f \in F(V)_n | f \text{ in general position}\}$. A sequence of vectors $x_1, \dots, x_n \in V$ is said to be in general position if no nontrivial linear combination of zero

$$a_1 x_1 + \cdots + a_n x_n = 0$$

with $a_i \in k$ and at most $\dim(V)$ many nonvanishing a_i exists.

If k is infinite the complex is exact. A general vanishing result is contained in our main theorem.

We introduce axioms for elements of a given (nonempty) set X to be in general position relative to some other elements.

Definition 1 (General position). Let $G_{l,m}$ be relations in $l + m$ variables from X ; $l > 0, m \geq 0$. The family $(G_{l,m})_{l,m \in \mathbb{N}}$ is called a general position relation if the following properties are satisfied.

Let x, y, z be finite sequences of elements of X of length l, m, n :

- (i) $G_{n,m}$ is symmetric in the first n and last m arguments.
- (ii) If $G_{l+m,n}(x, y; z)$ then $G_{l,m+n}(x; y, z)$. If $G_{l,m+n}(x; y, z)$ then $G_{l,n}(x; z)$.
- (iii) If $G_{l,m+n}(x; y, z)$ and $G_{m,n}(y; z)$ then $G_{l+m,n}(x, y; z)$.

For fixed G we say x is in G -general position to y iff $G_{l,m}(x; y)$.
 Given $x \in F(X)_l$ and $y \in F(X)_m$ we say x is in G -general position to y if every term of x is in G -general position to every term of y (for $l = 0$ we demand nothing).

Definition 2. Given a finite sequence b of elements of X the corresponding complex of words in general position to b is $(F_G(X; b))_n := \{f \in F(X)_n \mid f \text{ is in } G\text{-general position to } b\}$.

Before we can state the main theorem we have to introduce an invariant which determines an upper bound for the vanishing of the homology of $F_G(X)$.

Definition 3. Given a general position relation G we define $|G|$ to be the smallest natural number $n \geq 0$ such that there is a sequence x of elements of X with $\text{length}(x) = n$ such that there is no further element in X which is in general position to x .

Examples

- (i) Example (i) is induced by saying x is in G^{inj} -general position to y if the underlying sets of x and y are disjoint and the entries of x are distinct. We have $|G^{inj}| = \text{card}(X)$.
- (ii) Example (ii) is induced by saying (x_1, \dots, x_i) is in G^{vec} -general position to (y_1, \dots, y_j) if for all $a_l \in k$, $l = 1, \dots, i + j$, and only $\dim(V)$ many of them nonvanishing

$$a_1 x_1 + \dots + a_i x_i + a_{i+1} y_1 + \dots + a_{i+j} y_j = 0$$

implies $a_l = 0$ for all $l \in \{1, \dots, i\}$.

If $\text{card}(k) = \infty$ or $\dim(V) = \infty$ then $|G^{vec}| = \infty$.

Unfortunately the exact value of $|G^{vec}|$ is not known for all finite dimensional vector spaces over finite fields. The following lemma comprises what is known.

Lemma 2. (a) For $\dim(V) \geq \text{card}(k)$ we have $|G^{vec}| = \dim(V) + 1$.
 (b) For $\dim(V) = 2$ we have $|G^{vec}| = \text{card}(k) + 1$.

Proof of (a). Let $n = \dim(V) + 1$ and $(e_i)_{1 \leq i \leq n-1}$ be a basis of V . First we show $|G^{vec}| \geq n$. Otherwise we had a sequence $(x_i)_{1 \leq i \leq n-1}$, $x_i \in V$, such that there does not exist a vector $x \in V$ in G^{vec} -general position to (x_i) . This can only be true, if (x_i) is a basis of V . But if it is a basis, the element $x_1 + x_2 + \dots + x_{n-1}$ would be in general position to it. Contradiction.

Now the sequence $f = (e_1, \dots, e_{n-1}, e_1 + e_2 + \dots + e_{n-1})$ is maximal in the sense of Definition 3. To see this we have to show there is no vector $x = a_1 e_1 + \dots + a_{n-1} e_{n-1}$, $a_i \in k$, in G^{vec} -general position to f . If there is an index j such that $a_j = 0$, x is obviously not in general position to f . But if no coefficient in x vanishes two of them have to be equal since $\dim(V) \geq \text{card}(k)$. In case $a_1 = a_2$ we can write

$$x = a_1 (e_1 + \dots + e_{n-1}) + (a_3 - a_1) e_3 + \dots + (a_{n-1} - a_1) e_{n-1}.$$

This shows f is in fact maximal.

Proof of (b). For $\dim(V) = 2$ we have $x \in V$ in G^{vec} -general position to $y \in V$ iff $x \neq 0 \neq y$ and $[x] \neq [y] \in \mathbb{P}_k(V)$. So $|Gen^{vec}|$ is simply the number of k -rational points in $\mathbb{P}_k(V)$. \square

Question. What is $|G^{vec}|$ for $2 < \dim(V) < \text{card}(k)$?

Now the main theorem reads as follows.

Theorem 3. For a set X with general position condition G and a finite sequence $a = (a_1, \dots, a_l)$ of elements of X the corresponding homology groups $H_m(F_G(X; a))$ vanish for $m \leq (|G| - l - 1)/2$.

In contrast to Theorem 1 one should notice that it is in general not possible to erase the factor $1/2$ from the inequality of the theorem. It is left to the reader to find a counter-example.

Corollary 1. For fixed $m, d \geq 0$ we have $H_m(F_{G^{vec}}(k^d)) = 0$ for almost all finite fields k .

Proof. A simple cardinality argument shows

$$\lim_{\text{card}(k) \rightarrow \infty} |G^{vec}| = \infty$$

for fixed dimension of the underlying vector space and variable base field k . Now Corollary 1 follows from Theorem 3. \square

Proof of Theorem 3. Denote the degree by m . Exactness at $m = 0$ is trivial. We proceed by induction on $m \geq 1$. Let c be a cycle in $F_m(X; a_1, \dots, a_l)$.

Case $m = 1$:

We can suppose $c = (x) - (x')$. Because $1 \leq (|G| - l - 1)/2$ we have $l + 2 < |G|$ so that there exists $y \in X$ in G -general position to (x, x', a_1, \dots, a_l) .

According to Definition 1(iii) (y, x, x') is in G -general position to (a_1, \dots, a_l) and it is allowed to write $d((y)c) = c$.

Induction step:

Fix $x \in X$ in G -general position to (a_1, \dots, a_l) .

The simplest case is c in G -general position to (x, a_1, \dots, a_l) . Here we can apply a construction similar to Lemma 1; we have $c = d((x)c)$, since $(x)c$ is in G -general position to (a_1, \dots, a_l) .

We reduce to this case by changing c by boundaries. To be more precise we introduce a number $I(c) \in \{0, \dots, m\}$ which is m , iff the above applies, that is c in G -general position to (x, a_1, \dots, a_l) .

By adding boundaries we will see that we can increase $I(c)$.

For $g \in (F_G(X; a))_n$ we define $I(g) \in \{0, \dots, n\}$ as the the greatest natural number $i \leq n$ such that $\pi_i(v)$ is in G -general position to $(x, \pi'_i(v), a_1, \dots, a_l)$ for any term v of g (π_i denotes the projection to the first i entries and π'_i the projection to the last $n - i$ entries).

Reduction to $I(c) > 0$:

Suppose $I(c) = 0$. Let $c = \sum_j x_j$ with x_j terms. Choose for every j a $y_j \in X$ with y_j in G -general position to $(x, x_j, a_1, \dots, a_l)$. This is possible, since

$$\text{length}(x, x_j, a_1, \dots, a_l) = 1 + m + l < |G|.$$

Clearly

$$I(c - d(\sum_j (y_j) x_j)) > 0.$$

Now suppose $m > I(c) > 0$:

Write

$$c = \sum_j s_j x_j + x'$$

such that exactly those terms v of c for which $I(v) > I(c)$ are in x' , $\text{length}(s_j) = I(c)$ and all x_j are distinct terms.

Lemma 3. $d(s_j) = 0$ for all j .

Proof. The terms v of $d(c)$ such that $I(v) < I(c)$ are exactly the terms of $d(s_j) x_j$. In order to see this, notice that for a term v of $d(c)$ we have $I(v) = I(c) - 1$ iff the $I(c)$ -th entry of v is not in G -general position to $(x, \pi'_{I(c)}(v), a_1, \dots, a_l)$.

Now projecting the identity $0 = d(c)$ to the terms v with $I(v) < I(c)$ we get $0 = \sum_j d(s_j) x_j$. Lemma 3 is proven since the x_j are distinct terms. \square

According to our assumption s_j is in G -general position to $(x, x_j, a_1, \dots, a_l)$. By induction on m we know there exist s'_j with $d(s'_j) = s_j$ and s'_j in G -general position to $(x, x_j, a_1, \dots, a_l)$, because $2i + [(m - i) + l + 1] \leq 2m + l \leq |G| - 1$.

Now

$$\begin{aligned} I(c - d(\sum_j s'_j x_j)) &= I(x' + (-1)^{I(c)} s'_j d(x_j)) \\ &> I(c). \end{aligned}$$

This finishes the reduction to $I(c) = m$ and therefore the induction step to $\text{length}(c) = m$ is accomplished by applying the standard trick mentioned at the beginning of the proof. \square

Theorem 3 could be useful in the generalization of Suslin's GL -stability [8] to finite fields. A thorough treatment seems to indicate the following result:

(i) Given $m \geq 0$ and $n \geq m$ we have $H_m(GL_n(k)) = H_m(GL_{n+1}(k))$ for almost all finite fields k .

(ii) For $m \geq 0$ the map $H_m(GL_{m-1}(k)) \rightarrow H_m(GL_m(k))$ is surjective for almost all finite fields k .

In fact Quillen proved – using other methods – that $H_m(GL_n(k)) \rightarrow H_m(GL_{n+1}(k))$ is an isomorphism for all fields k with more than 2 elements and $n > m$ [6]. Similar results with weaker bounds are due to Maazen and van der Kallen [4].

Acknowledgement

The above results were obtained in the Seminar Müller-Stach 2004 at the University of Mainz. I am indebted to Philippe Elbaz-Vincent, Stefan Müller-Stach, Oliver Petras, Wilberd van der Kallen and Volkmar Welker for their comments.

Especially I have to thank Volkmar Welker for his hospitality in Marburg.

References

- [1] Björner, Anders; Wachs, Michelle *On lexicographically shellable posets.* Trans. Amer. Math. Soc. **277** (1983), no. 1, 323–341.
- [2] Brown, Kenneth S. *Cohomology of groups.* Corrected reprint of the 1982 original. Graduate Texts in Mathematics, 87. Springer-Verlag, New York, 1994
- [3] Farmer, Frank D. *Cellular homology for posets.* Math. Japon. **23** (1978/79), no. 6, 607–613.
- [4] van der Kallen, Wilberd *Homology stability for linear groups.* Invent. Math. **60** (1980), no. 3, 269–295.
- [5] Nakaoka, Minoru *Decomposition theorem for homology groups of symmetric groups.* Ann. of Math. (2) 71 1960 16–42.
- [6] Quillen, Daniel; unpublished.
- [7] Stanley, Richard P. *Combinatorics and commutative algebra.* Second edition. Progress in Mathematics, 41. Birkhäuser Boston, Inc., Boston, MA, 1996.
- [8] Suslin, A. A. *Homology of GL_n , characteristic classes and Milnor K -theory.* Algebraic K -theory, number theory, geometry and analysis (Bielefeld, 1982), 357–375, Lecture Notes in Math., 1046, Springer, Berlin, 1984.
- [9] Suslin, A. A. *K_3 of a field, and the Bloch group.* Proc. Steklov Inst. Math. 1991, no. 4, 217–239.

This article may be accessed via WWW at <http://www.rmi.acnet.ge/hha/>
or by anonymous ftp at
[ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2005/n1a5/v7n1a5.\(dvi,ps,pdf\)](ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2005/n1a5/v7n1a5.(dvi,ps,pdf))

Moritz C. Kerz mokerz@students.uni-mainz.de
Institut für Mathematik (Fachbereich 8)
Johannes Gutenberg-Universität Mainz
Staudingerweg 9
55099 Mainz
Germany