

## 12. Two types of complete discrete valuation fields

*Masato Kurihara*

In this section we discuss results of a paper [Ku1] which is an attempt to understand the structure of the Milnor  $K$ -groups of complete discrete valuation fields of mixed characteristics in the case of an arbitrary residue field.

### 12.0. Definitions

Let  $K$  be a complete discrete valuation field of mixed characteristics  $(0, p)$  with the ring of integers  $\mathcal{O}_K$ . We consider the  $p$ -adic completion  $\widehat{\Omega}_{\mathcal{O}_K}^1$  of  $\Omega_{\mathcal{O}_K/\mathbb{Z}}^1$  as in section 9.

Note that

- (a) If  $K$  is a finite extension of  $\mathbb{Q}_p$ , then

$$\widehat{\Omega}_{\mathcal{O}_K}^1 = (\mathcal{O}_K/\mathcal{D}_{K/\mathbb{Q}_p})d\pi$$

where  $\mathcal{D}_{K/\mathbb{Q}_p}$  is the different of  $K/\mathbb{Q}_p$ , and  $\pi$  is a prime element of  $K$ .

- (b) If  $K = k\{\{t_1\}\} \cdots \{\{t_{n-1}\}\}$  with  $|k : \mathbb{Q}_p| < \infty$  (for the definition see subsection 1.1), then

$$\widehat{\Omega}_{\mathcal{O}_K}^1 = (\mathcal{O}_k/\mathcal{D}_{k/\mathbb{Q}_p})d\pi \oplus \mathcal{O}_K dt_1 \oplus \cdots \oplus \mathcal{O}_K dt_{n-1}$$

where  $\pi$  is a prime element of  $\mathcal{O}_k$ .

But in general, the structure of  $\widehat{\Omega}_{\mathcal{O}_K}^1$  is a little more complicated. Let  $F$  be the residue field of  $K$ , and consider a natural map

$$\varphi: \widehat{\Omega}_{\mathcal{O}_K}^1 \longrightarrow \Omega_F^1.$$

**Definition.** Let  $\text{Tors } \widehat{\Omega}_{\mathcal{O}_K}^1$  be the torsion part of  $\widehat{\Omega}_{\mathcal{O}_K}^1$ . If  $\varphi(\text{Tors } \widehat{\Omega}_{\mathcal{O}_K}^1) = 0$ ,  $K$  is said to be of *type I*, and said to be of *type II* otherwise.

So if  $K$  is a field in (a) or (b) as above,  $K$  is of type I.

Let  $\pi$  be a prime element and  $\{t_i\}$  be a lifting of a  $p$ -base of  $F$ . Then, there is a relation

$$ad\pi + \sum b_i dt_i = 0$$

with  $a, b_i \in \mathcal{O}_K$ . The field  $K$  is of type I if and only if  $v_K(a) < \min_i v_K(b_i)$ , where  $v_K$  is the normalized discrete valuation of  $K$ .

**Examples.**

- (1) If  $v_K(p)$  is prime to  $p$ , or if  $F$  is perfect, then  $K$  is of type I.
- (2) The field  $K = \mathbb{Q}_p\{\{t\}\}(\pi)$  with  $\pi^p = pt$  is of type II. In this case we have

$$\widehat{\Omega}_{\mathcal{O}_K}^1 \simeq \mathcal{O}_K/p \oplus \mathcal{O}_K.$$

The torsion part is generated by  $dt - \pi^{p-1}d\pi$  (we have  $pdt - p\pi^{p-1}d\pi = 0$ ), so  $\varphi(dt - \pi^{p-1}d\pi) = dt \neq 0$ .

## 12.1. The Milnor $K$ -groups

Let  $\pi$  be a prime element, and put  $e = v_K(p)$ . Section 4 contains the definition of the homomorphism

$$\rho_m: \Omega_F^{q-1} \oplus \Omega_F^{q-2} \longrightarrow \text{gr}_m K_q(K).$$

**Theorem.** Put  $\ell = \text{length}_{\mathcal{O}_K}(\text{Tors } \widehat{\Omega}_{\mathcal{O}_K}^1)$ .

- (a) If  $K$  is of type I, then for  $m \geq \ell + 1 + 2e/(p-1)$

$$\rho_m|_{\Omega_F^{q-1}}: \Omega_F^{q-1} \longrightarrow \text{gr}_m K_q(K)$$

is surjective.

- (b) If  $K$  is of type II, then for  $m \geq \ell + 2e/(p-1)$  and for  $q \geq 2$

$$\rho_m|_{\Omega_F^{q-2}}: \Omega_F^{q-2} \longrightarrow \text{gr}_m K_q(K)$$

is surjective.

For the proof we used the exponential homomorphism for the Milnor  $K$ -groups defined in section 9.

**Corollary.** Define the subgroup  $U_i K_q(K)$  of  $K_q(K)$  as in section 4, and define the subgroup  $V_i K_q(K)$  as generated by  $\{1 + \mathcal{M}_K^i, \mathcal{O}_K^*, \dots, \mathcal{O}_K^*\}$  where  $\mathcal{M}_K$  is the maximal ideal of  $\mathcal{O}_K$ .

- (a) If  $K$  is of type I, then for sufficiently large  $m$  we have  $U_m K_q(K) = V_m K_q(K)$ .
- (b) If  $K$  is of type II, then for sufficiently large  $m$ , we have  $V_m K_q(K) = U_{m+1} K_q(K)$ .  
Especially,  $\text{gr}_m K_q(K) = 0$  for sufficiently large  $m$  prime to  $p$ .

**Example.** Let  $K = \mathbb{Q}_p\{\{t\}\}(\pi)$  where  $\pi^p = pt$  as in Example (2) of subsection 12.0, and assume  $p > 2$ . Then, we can determine the structures of  $\text{gr}_m K_q(K)$  as follows ([Ku2]).

For  $m \leq p + 1$ ,  $\text{gr}_m K_q(K)$  is determined by Bloch and Kato ([BK]). We have an isomorphism  $\text{gr}_0 K_2(K) = K_2(K)/U_1 K_2(K) \simeq K_2(F) \oplus F^*$ , and  $\text{gr}_p K_q(K)$  is a certain quotient of  $\Omega_F^1/dF \oplus F$  (cf. [BK]). The homomorphism  $\rho_m$  induces an isomorphism from

$$\left\{ \begin{array}{ll} \Omega_F^1 & \text{if } 1 \leq m \leq p - 1 \text{ or } m = p + 1 \\ 0 & \text{if } i \geq p + 2 \text{ and } i \text{ is prime to } p \\ F/F^p & \text{if } m = 2p \\ & (x \mapsto \{1 + p\pi^p x, \pi\} \text{ induces this isomorphism}) \\ F^{p^{n-2}} & \text{if } m = np \text{ with } n \geq 3 \\ & (x \mapsto \{1 + p^n x, \pi\} \text{ induces this isomorphism}) \end{array} \right.$$

onto  $\text{gr}_m K_2(K)$ .

## 12.2. Cyclic extensions

For cyclic extensions of  $K$ , by the argument using higher local class field theory and the theorem of 12.1 we have (cf. [Ku1])

**Theorem.** *Let  $\ell$  be as in the theorem of 12.1.*

- (a) *If  $K$  is of type I and  $i \geq 1 + \ell + 2e/(p - 1)$ , then  $K$  does not have ferociously ramified cyclic extensions of degree  $p^i$ . Here, we call an extension  $L/K$  ferociously ramified if  $|L : K| = |k_L : k_K|_{\text{ins}}$  where  $k_L$  (resp.  $k_K$ ) is the residue field of  $L$  (resp.  $K$ ).*
- (b) *If  $K$  is of type II and  $i \geq \ell + 2e/(p - 1)$ , then  $K$  does not have totally ramified cyclic extensions of degree  $p^i$ .*

The bounds in the theorem are not so sharp. By some consideration, we can make them more precise. For example, using this method we can give a new proof of the following result of Miki.

**Theorem (Miki, [M]).** *If  $e < p - 1$  and  $L/K$  is a cyclic extension, the extension of the residue fields is separable.*

For  $K = \mathbb{Q}_p\{\{t\}\}(\sqrt[p]{pt})$  with  $p > 2$ , we can show that it has no cyclic extensions of degree  $p^3$ .

Miki also showed that for any  $K$ , there is a constant  $c$  depending only on  $K$  such that  $K$  has no ferociously ramified cyclic extensions of degree  $p^i$  with  $i > c$ .

For totally ramified extensions, we guess the following. Let  $F^{p^\infty}$  be the maximal perfect subfield of  $F$ , namely  $F^{p^\infty} = \bigcap F^{p^n}$ . We regard the ring of Witt vectors  $W(F^{p^\infty})$  as a subring of  $\mathcal{O}_K$ , and write  $k_0$  for the quotient field of  $W(F^{p^\infty})$ , and write  $k$  for the algebraic closure of  $k_0$  in  $K$ . Then,  $k$  is a finite extension of  $k_0$ , and is a complete discrete valuation field of mixed characteristics  $(0, p)$  with residue field  $F^{p^\infty}$ .

**Conjecture.** *Suppose that  $e(K|k) > 1$ , i.e. a prime element of  $\mathcal{O}_k$  is not a prime element of  $\mathcal{O}_K$ . Then there is a constant  $c$  depending only on  $K$  such that  $K$  has no totally ramified cyclic extension of degree  $p^i$  with  $i > c$ .*

### References

- [BK] S. Bloch and K. Kato,  $p$ -adic etale cohomology, Publ. Math. IHES, 63(1986), 107–152.
- [Ku1] M. Kurihara, On two types of complete discrete valuation fields, Comp. Math., 63(1987), 237–257.
- [Ku2] M. Kurihara, On the structure of the Milnor  $K$ -group of a certain complete discrete valuation fields, preprint.
- [M] H. Miki, On  $\mathbb{Z}_p$ -extensions of complete  $p$ -adic power series fields and function fields, J. Fac. Sci. Univ. Tokyo, Sect. 1A Math., 21(1974), 377–393.

*Department of Mathematics Tokyo Metropolitan University  
Minami-Osawa 1-1, Hachioji, Tokyo 192-03, Japan  
E-mail: m-kuri@comp.metro-u.ac.jp*