

THIN SEVERI–BRAUER VARIETIES

MAX-ALBERT KNUS AND JEAN-PIERRE TIGNOL

À Jacques Tits, pour son 80^{ème} anniversaire

ABSTRACT. Severi–Brauer varieties are twisted forms of projective spaces (in the sense of Galois cohomology) and are associated in a functorial way to central simple algebras. Similarly quadrics are related to algebras with involution. Since thin projective spaces are finite sets, thin Severi–Brauer varieties are finite sets endowed with a Galois action; they are associated to étale algebras. Similarly, thin quadrics are étale algebras with involution. We discuss embeddings of thin Severi–Brauer varieties and thin quadrics in Severi–Brauer varieties and quadrics as geometric analogues of embeddings of étale algebras into central simple algebras (with or without involution), and consider the geometric counterpart of the Clifford algebra construction.

1. INTRODUCTION

Paraphrasing Tits [18], “thin” geometric objects are characterised by the fact that their automorphism groups are the Weyl groups of the groups of automorphisms of the corresponding classical geometric objects.¹ Thus a thin $(n - 1)$ -dimensional projective space is a finite set of n elements and a $2(n - 1)$ -dimensional thin quadric is a double covering of a set of n elements.

Our starting point is to consider twisted forms (in the sense of Galois cohomology) of projective spaces and quadrics. Fixing a base field, a twist is given by a continuous action of the absolute Galois group of the field. Twisted forms of projective spaces over a field F occur as Severi–Brauer varieties associated to central simple algebras over F , and twisted quadrics correspond to central simple algebras with quadratic pair. Similarly, twisted forms of finite sets are associated to algebraic objects, namely étale algebras, by a correspondence known as Grothendieck’s version of Galois theory. Double coverings correspond to étale algebras with involution. Moreover the Grassmannian of maximal linear subspaces of a split quadric of dimension $2(n - 1)$ corresponds as a geometric object to the Clifford algebra of a central algebra with quadratic pair as algebraic object. We call this Grassmannian the Clifford set of the quadric.

The first aim of the paper is to study equivariant embeddings of finite twisted objects into corresponding classical objects. For example a general embedding of a finite set of n elements in a projective space of dimension $n - 1$ corresponds to the embedding of an étale algebra of rank n in a central simple algebra of degree n (or to the embedding of a torus of rank $n - 1$ into a twisted form of SL_n , see [13, S2]). The second aim is to discuss in detail the geometric side of the Clifford

The second author is supported in part by the F.R.S.–FNRS (Belgium).

¹In [16] Tits says that such objects are defined over the “field of characteristic 1”.

construction, and to relate it to Clifford algebras for étale algebras with involution and for central simple algebras with quadratic pairs.

In Section 2 we consider Severi–Brauer varieties, in Section 3 quadrics, and in Section 4 the various types of Clifford constructions, where we introduce a special type of Severi–Brauer variety to describe the geometry of a Clifford algebra. Finally we sketch in Section 5 a discussion of two exceptional phenomena. First we describe consequences of the fact that the Dynkin diagrams A_3 and D_3 are equal. This is related on the geometric side with the Klein quadric and on the algebraic side with the resolution of the equation of degree 4. Then we look at triality, that is, outer automorphisms of order 3 of D_4 .

One origin of this paper is the observation of Tits [16, p. 287] that the two phenomena described above also occur over the “field of characteristic 1”. Also, his short geometric proof of the fact that two quaternion algebras whose tensor product is not division share a common quadratic subfield [19] is a beautiful illustration of the power of geometric insights into algebraic questions. Another source of inspiration was [14], where Saltman discusses the embedding of étale algebras with involution into central simple algebras with involution. Pairs consisting of a maximal étale subalgebra of a central simple algebra (and corresponding pairs of algebras with involutions) are also considered in [1] and [13]. Triality in relation with étale algebras is extensively discussed in [9]. We plan to come back to other aspects, in particular to unitary and symplectic analogues, in [10].

2. SEVERI–BRAUER PROJECTIVE SPACES

Γ -projective spaces and central simple algebras. Throughout most of this work, F is an arbitrary field. We denote by F_s a separable closure of F and by Γ the absolute Galois group $\text{Gal}(F_s/F)$, which is a profinite group.

A Γ -projective space P of dimension $n - 1$ over a field F is a projective space $\mathbb{P}^{n-1}(F_s)$ of dimension $n - 1$ endowed with the discrete topology and a continuous action of Γ by collineations, denoted by $x \mapsto \gamma x$, $x \in P$, $\gamma \in \Gamma$. For every integer $k \in \{1, \dots, n\}$, the Grassmannian $Gr_{k-1}(P)$ of $(k - 1)$ -dimensional linear varieties in P carries an action of Γ . We view the set:

$$\overline{P} = \{x \in P \mid \gamma x = x \text{ for all } \gamma \in \Gamma\} \subset P$$

as a scheme over F . The embedding $\overline{P} \subset P$ induces an isomorphism

$$\beta: \overline{P} \times_{\text{Spec } F} \text{Spec } F_s \xrightarrow{\sim} P$$

of Γ -spaces. From now on we shall not distinguish between P and the pair (\overline{P}, β) . We have a similar construction for Grassmannians.

Continuous actions of Γ are related with Galois cohomology, as we now illustrate in the case of Γ -projective spaces. There is a standard action of Γ on $\mathbb{P}^{n-1}(F_s)$ induced by the action of Γ on F_s . We denote it by $x \mapsto \tilde{\gamma}x$. This standard action induces in turn a Γ -action on the group $\text{PGL}_n(F_s)$ of automorphisms of the projective space $\mathbb{P}^{n-1}(F_s)$, given by $\varphi \mapsto \tilde{\gamma}(\varphi) = \tilde{\gamma}\varphi\tilde{\gamma}^{-1}$, $\varphi \in \text{PGL}_n(F_s)$.

A 1-cocycle of Γ with values in PGL_n is a continuous map $\Gamma \rightarrow \text{PGL}_n(F_s)$, $\gamma \mapsto a_\gamma$, such that

$$a_{\gamma\delta} = a_\gamma \tilde{\gamma}(a_\delta) \quad \text{for all } \gamma, \delta \in \Gamma.$$

Two cocycles a_γ, a'_γ are *cohomologous or equivalent* if there exists $a \in \mathrm{PGL}_n(F_s)$ such that $a'_\gamma = aa_\gamma\bar{\gamma}(a)^{-1}$. We denote by $H^1(\Gamma, \mathrm{PGL}_n)$ the set of equivalence classes of 1-cocycles.

Let $P = (\bar{P}, \beta)$ be any Γ -projective space. The map $a_\gamma = \beta\gamma\beta^{-1}\bar{\gamma}^{-1}$, $\gamma \in \Gamma$, is an element of $\mathrm{PGL}_n(F_s)$ and defines a 1-cocycle. Moreover one shows that isomorphic Γ -projective spaces lead to cohomologous cocycles. Let $\mathrm{Proj}_\Gamma^{n-1}$ be the groupoid² of Γ -projective spaces and let $\mathrm{Iso}(\mathrm{Proj}_\Gamma^{n-1})$ be its set of isomorphism classes.

Proposition 2.1. *The map $(P, \beta) \mapsto a_\gamma$ induces a bijection*

$$\mathrm{Iso}(\mathrm{Proj}_\Gamma^{n-1}) \xrightarrow{\sim} H^1(\Gamma, \mathrm{PGL}_n).$$

Proof. We describe a map in the opposite direction and refer to [15, p. 161] for details. Given a 1-cocycle $(a_\gamma)_{\gamma \in \Gamma}$, we define a Γ -action on $P = \mathbb{P}^{n-1}(F_s)$ as

$$x \mapsto^\gamma x = a_\gamma \bar{\gamma} x, \quad x \in P. \quad \square$$

The Γ -projective spaces we consider arise from the following construction. Recall that an F -algebra A is *central simple of degree n* if there exists an isomorphism $\alpha: A \otimes_F F_s \xrightarrow{\sim} \mathrm{End}_{F_s}(V)$ for a vector space V of dimension n over F_s . Since the group of F_s -automorphisms of $\mathrm{End}_{F_s}(V)$ is isomorphic to $\mathrm{PGL}_n(F_s)$, a 1-cocycle defining a Γ -projective space of dimension $n - 1$ is also a 1-cocycle defining a central simple algebra of degree n . Thus denoting by CSA_F^n the groupoid of central simple algebras of degree n , the sets $\mathrm{Iso}(\mathrm{Proj}_\Gamma^{n-1})$ and $\mathrm{Iso}(\mathrm{CSA}_F^n)$ are in bijection, since they are in bijection with $H^1(\Gamma, \mathrm{PGL}_n)$.

There is a canonical way to associate a Γ -projective space to a central simple algebra. Let $\alpha: A \otimes_F F_s \xrightarrow{\sim} \mathrm{End}_{F_s}(V)$ as above. Since the n -dimensional right ideals of $\mathrm{End}_{F_s}(V)$ are of the form $\mathrm{Hom}_{F_s}(V, U)$ for $U \subset V$ a 1-dimensional subspace, any isomorphism $A_s \simeq \mathrm{End}_{F_s}(V)$ defines a one-to-one correspondence between n -dimensional right ideals of A_s and 1-dimensional subspaces of V . Therefore, we may canonically associate to A a projective space $\mathcal{SB}(A)$ over F_s , whose $(k - 1)$ -dimensional linear varieties are the nk -dimensional right ideals of A_s . The action of the group Γ of F_s over F induces a continuous action on A_s , hence also on the right ideals of A_s , and endows $\mathcal{SB}(A)$ with a Γ -projective space structure. We call $\mathcal{SB}(A)$ (or its associated scheme $\underline{\mathcal{SB}}(A)$ over F) the *Severi–Brauer variety of A* . Its dimension is $n - 1$, where $n = \deg A$.

Proposition 2.2. *The rule $A \mapsto \mathcal{SB}(A)$ induces an anti-equivalence*

$$\mathrm{CSA}_F^n \equiv \mathrm{Proj}_\Gamma^{n-1}.$$

Proof. Applying [11, Theorem 1, p. 93] it suffices to show that the functor \mathcal{SB} is fully faithful and essentially surjective. We refer to [6] for details. \square

Γ -sets and étale algebras. Finite sets with a continuous left action of Γ (for the discrete topology) are called (finite) Γ -sets. They form a category Set_Γ whose morphisms are the Γ -equivariant maps. Finite direct products and direct sums (= disjoint unions) are defined in this category. We denote by $|X|$ the cardinality of any finite set X .

²A groupoid is a category in which all morphisms are isomorphisms.

Following Tits [18] we define a *thin* Γ -projective space of dimension $n - 1$ as a Γ -set X of n elements. A *thin* $(k - 1)$ -dimensional linear subvariety is a Γ -subset of X with k elements. For every integer k , $1 \leq k \leq n$, the Γ -set

$$\begin{aligned} Gr_{k-1}(X) &= \{(k-1)\text{-dimensional linear varieties contained in } X\} \\ &= \{k\text{-element subsets of } X\} \end{aligned}$$

is a *thin* Γ -Grassmannian. As for Γ -projective spaces over F , we shall not distinguish between a finite Γ -set X and its scheme \overline{X} over F of fixed points of the Γ -action. Like Γ -projective spaces over F , thin Γ -projective spaces are related with algebraic structures. We recall that a finite-dimensional commutative F -algebra L is called *étale* (over F) if $L \otimes_F F_s$ is isomorphic to the F_s -algebra $F_s^n = F_s \times \cdots \times F_s$ (n factors) for some $n \geq 1$. Étale F -algebras are the direct products of finite separable field extensions of F . We refer to [7, §18.A] for various equivalent characterizations of étale F -algebras. These algebras (with F -algebra homomorphisms) form a category $\acute{E}t_F$ in which finite direct products and finite direct sums (= tensor products) are defined.

For any étale F -algebra L of dimension n , the set of F -algebra homomorphisms

$$\mathbf{X}(L) = \text{Hom}_{F\text{-alg}}(L, F_s)$$

is a Γ -set of n elements since Γ acts on F_s . Conversely, if X is a Γ -set of n elements, the F -algebra $\mathbf{M}(X)$ of Γ -equivariant maps $X \rightarrow F_s$ is an étale F -algebra of dimension n ,

$$\mathbf{M}(X) = \{f: X \rightarrow F_s \mid \gamma(f(x)) = f(\gamma x) \text{ for } \gamma \in \Gamma, x \in X\}.$$

As first observed by Grothendieck, there are canonical isomorphisms

$$\mathbf{M}(\mathbf{X}(L)) \cong L, \quad \mathbf{X}(\mathbf{M}(X)) \cong X,$$

so that the functors \mathbf{M} and \mathbf{X} define an anti-equivalence of categories³

$$(2.3) \quad \text{Set}_\Gamma \cong \acute{E}t_F$$

(see [2, Proposition (4.3), p. 25] or [7, (18.4)]), which we refer to as the *Grothendieck correspondence*. Under this anti-equivalence, the cardinality of Γ -sets corresponds to the dimension of étale F -algebras. The disjoint union \sqcup in Set_Γ corresponds to the product \times in $\acute{E}t_F$ and the product \times in Set_Γ to the tensor product \otimes in $\acute{E}t_F$. For any integer $n \geq 1$, we let $\acute{E}t_F^n$ denote the groupoid whose objects are n -dimensional étale F -algebras and whose morphisms are F -algebra isomorphisms, and Set_Γ^n the groupoid of Γ -sets with n elements. The anti-equivalence (2.3) restricts to an anti-equivalence $\text{Set}_\Gamma^n \cong \acute{E}t_F^n$. Isomorphism classes are in bijection with $H^1(\Gamma, \mathfrak{S}_n)$, where Γ acts trivially on \mathfrak{S}_n (see [7, 29.9]).

The split étale algebra F^n corresponds to the Γ -set \mathbf{n} of n elements with trivial Γ -action. Étale algebras of dimension 2 are also called *quadratic* étale algebras.

Embeddings of Γ -sets in Severi–Brauer varieties. For X a Γ -set of n elements and a Γ -projective space P of dimension $n - 1$, we define a *general embedding*

$$\varepsilon: X \hookrightarrow P$$

³We let morphisms of Γ -sets act on the right of the arguments (with the exponential notation) and use the usual function notation for morphisms in the anti-equivalent category of étale algebras.

to be a Γ -equivariant injective map whose image consists of n points in general position, i.e., not contained in a proper subspace⁴. The map ε then induces a map

$$\text{Gr}_{k-1}\varepsilon: \text{Gr}_{k-1}(X) \hookrightarrow \text{Gr}_{k-1}(P) \quad \text{for } 1 \leq k \leq n,$$

which carries each k -element subset of X to the linear variety spanned by its image. In the rest of this subsection, we relate F -algebra embeddings of n -dimensional étale F -algebras in a central simple F -algebra A with general embeddings of Γ -sets of n elements in $\mathbf{SB}(A)$. We first discuss the case where the étale F -algebra is split.

Proposition 2.4. *Let L be a split étale F -algebra of dimension n , and let A be a central simple F -algebra of degree n . For every F -algebra embedding $\varepsilon: L \hookrightarrow A$, there is an F -algebra isomorphism $\varphi: A \rightarrow \text{End}_F L$ such that $\varphi \circ \varepsilon: L \hookrightarrow \text{End}_F L$ is the regular representation, which maps each $x \in L$ to multiplication by x .*

Proof. Let $(e_i)_{1 \leq i \leq n}$ be the collection of primitive idempotents of L . We have

$$A = \varepsilon(e_1)A \oplus \cdots \oplus \varepsilon(e_n)A.$$

Since the dimension of every right ideal of A is a multiple of $n \text{ Ind } A$, we must have

$$\dim \varepsilon(e_i)A = n \text{ for } 1 \leq i \leq n, \quad \text{and} \quad \text{Ind } A = 1.$$

Identifying $A = \text{End}_F V$ for some F -vector space V of dimension n , we may view $\varepsilon(e_1), \dots, \varepsilon(e_n)$ as projections on 1-dimensional subspaces V_1, \dots, V_n such that $V = V_1 \oplus \cdots \oplus V_n$. Any F -vector space isomorphism $V \rightarrow L$ that maps V_i to $e_i L$ for all $1 \leq i \leq n$ determines an F -algebra isomorphism $\varphi: A \rightarrow \text{End}_F L$ as required. \square

Now, let L be an arbitrary n -dimensional étale F -algebra. Consider an F -algebra embedding into a central simple F -algebra of degree n :

$$\varepsilon: L \hookrightarrow A.$$

For every $\xi \in \mathbf{X}(L) := \text{Hom}_{F\text{-alg}}(L, F_s)$, set

$$(2.5) \quad \varepsilon_*(\xi) = \{x \in A_s \mid (1 \otimes \xi(\ell)) \cdot x = (\varepsilon(\ell) \otimes 1) \cdot x \text{ for all } \ell \in L\}.$$

Proposition 2.6. *The right ideal $\varepsilon_*(\xi)$ is n -dimensional, and ε_* is a general embedding*

$$\varepsilon_*: \mathbf{X}(L) \hookrightarrow \mathbf{SB}(A).$$

Proof. Let $L_s = L \otimes_F F_s$, a split étale F_s -algebra, and $\varepsilon_s = \varepsilon \otimes \text{Id}_{F_s}: L_s \hookrightarrow A_s$. By Proposition 2.4, we may identify $A_s = \text{End}_{F_s} L_s$ in such a way that $\varepsilon_s: L_s \hookrightarrow \text{End}_{F_s} L_s$ is the regular representation. Let $(e_i)_{1 \leq i \leq n}$ be the collection of primitive idempotents of L_s . Using the canonical correspondence $\text{Hom}_{F\text{-alg}}(L, F_s) = \text{Hom}_{F_s\text{-alg}}(L_s, F_s)$, we may set

$$\mathbf{X}(L) = \{\xi_1, \dots, \xi_n\}$$

where $\xi_i: L_s \rightarrow F_s$ maps e_i to 1 and e_j to 0 for $j \neq i$. For $f \in \text{End}_{F_s} L_s$, the condition

$$(1 \otimes \xi_i(\ell)) \cdot f = (\varepsilon(\ell) \otimes 1) \circ f \quad \text{for all } \ell \in L$$

is equivalent to

$$(1 \otimes \xi_i(\ell)) \cdot f = \varepsilon_s(\ell) \circ f \quad \text{for all } \ell \in L_s.$$

⁴Such embeddings are called *frames* in [18].

For $\ell = e_1, \dots, e_n$, this condition can be rewritten as

$$e_j \cdot f(x) = 0 \quad \text{and} \quad e_i \cdot f(x) = f(x) \quad \text{for all } x \in L_s.$$

Therefore,

$$\varepsilon_*(\xi_i) = \text{Hom}_{F_s}(L_s, e_i L_s),$$

which shows that $\dim \varepsilon_*(\xi_i) = n$. Moreover,

$$\varepsilon_*(\xi_1) + \dots + \varepsilon_*(\xi_n) = \text{Hom}_{F_s}(L_s, e_1 L_s + \dots + e_n L_s) = \text{End}_{F_s} L_s.$$

Therefore $\varepsilon_*(\xi_1), \dots, \varepsilon_*(\xi_n)$ are in general position in the projective space $\mathbf{SB}(A)$. To complete the proof, we show ε_* is Γ -equivariant. For $\gamma \in \Gamma$, we have

$$\gamma(\varepsilon_*(\xi)) = \{x \in A_s \mid (1 \otimes \xi(\ell)) \cdot \gamma^{-1}(x) = (\varepsilon(\ell) \otimes 1) \cdot \gamma^{-1}(x) \text{ for all } \ell \in L\}.$$

By applying γ to each side, we see that the condition

$$(1 \otimes \xi(\ell)) \cdot \gamma^{-1}(x) = (\varepsilon(\ell) \otimes 1) \cdot \gamma^{-1}(x)$$

is equivalent to

$$(1 \otimes \gamma \xi(\ell)) \cdot x = (\varepsilon(\ell) \otimes 1) \cdot x.$$

Thus, $\gamma(\varepsilon_*(\xi)) = \varepsilon_*(\gamma \xi)$. \square

Conversely, every general embedding $X \hookrightarrow \mathbf{SB}(A)$ of a Γ -set X of n elements arises from an F -algebra embedding, as we now show.

Given n -dimensional right ideals $I_1, \dots, I_n \subset A_s$ such that $\{I_1, \dots, I_n\}$ is stable under Γ and $I_1 + \dots + I_n = A_s$, let

$$L_s = \{x \in A_s \mid xI_i \subset I_i \text{ for } 1 \leq i \leq n\}.$$

The set L_s is an F_s -subalgebra of A_s stable under the action of Γ . Let $L = L_s^\Gamma$ be the fixed F -subalgebra.

Proposition 2.7. *The F -algebra L is étale of dimension n and $\{I_1, \dots, I_n\}$ is the image of the general embedding $\varepsilon_*: \mathbf{X}(L) \hookrightarrow \mathbf{SB}(A)$ induced by the inclusion $\varepsilon: L \hookrightarrow A$.*

Proof. We may identify $A_s = M_n(F_s)$ in such a way that for $1 \leq i \leq n$ the ideal I_i consists of matrices whose nonzero entries are on the i -th row. For $x = (x_{ij})_{1 \leq i, j \leq n}$ we have $xI_j \subset I_j$ if and only if $x_{ij} = 0$ for $i \neq j$. Therefore, L_s is the algebra of diagonal matrices. Since $L_s = L \otimes_F F_s$, it follows that L is an étale F -algebra of dimension n . Under the canonical identification $\mathbf{X}(L) = \text{Hom}_{F_s\text{-alg}}(L_s, F_s)$ we have $\mathbf{X}(L) = \{\xi_1, \dots, \xi_n\}$ where $\xi_i: L_s \rightarrow F_s$ yields the i -th diagonal entry. Matrix computation shows that $\varepsilon_*(\xi_i) = I_i$. \square

Remark 2.8. The definition of the map ε_* is inspired by the correspondence between maximal subfields of a division algebra and closed points on the corresponding Severi–Brauer variety set up by Merkurjev in [12].

3. QUADRICS AND ALGEBRAS WITH INVOLUTION

Γ -quadratics and quadratic pairs. As in the preceding section, let F be an arbitrary field and Γ the Galois group of some separable closure F_s of F . We call Γ -quadratic in a $(2n - 1)$ -dimensional Γ -projective space P over F any quadric in P defined by a hyperbolic quadratic form and stable under the action of Γ . The Γ -quadratics we consider in the sequel arise from quadratic pairs on central simple algebras, which we now recall from [7, §5].

Let B be an arbitrary F -algebra. An *involution*⁵ on B is an F -linear anti-automorphism of B of order 2. The sets of symmetric, skew-symmetric, symmetrized and alternating elements are the following subspaces of B :

$$\begin{aligned}\mathrm{Sym}(B, \sigma) &= \{x \in B \mid \sigma(x) = x\} \\ \mathrm{Skew}(B, \sigma) &= \{x \in B \mid \sigma(x) = -x\} \\ \mathrm{Symd}(B, \sigma) &= \{x + \sigma(x) \mid x \in B\} \\ \mathrm{Alt}(B, \sigma) &= \{x - \sigma(x) \mid x \in B\}.\end{aligned}$$

Now, let A be a central simple F -algebra of degree $2n$. A *quadratic pair* on A is a pair (σ, f) where σ is an involution on A and $f: \mathrm{Sym}(A, \sigma) \rightarrow F$ is a linear map, subject to the following conditions:

- (1) $\dim_F \mathrm{Sym}(A, \sigma) = n(2n + 1)$ and $\mathrm{Trd}_A(\mathrm{Skew}(A, \sigma)) = \{0\}$;
- (2) $f(x + \sigma(x)) = \mathrm{Trd}_A(x)$ for all $x \in A$.

If $A = \mathrm{End}_F V$, any quadratic pair (σ, f) on A is attached to a nonsingular quadratic form q on V . The involution σ is the involution adjoint to the quadratic form q and we refer to [7, §5B] for the definition of f .

For a central simple algebra A over a field of characteristic not 2 a quadratic pair (σ, f) is determined by the involution σ , which has to be of orthogonal type (see [7, §5B]).

Any quadratic pair (σ, f) on A extends to a quadratic pair (σ_s, f_s) on $A_s = A \otimes_F F_s$. Since A_s is split, we may identify $A_s = \mathrm{End}_{F_s}(V)$ for some $2n$ -dimensional F_s -vector space V . Then (σ_s, f_s) is adjoint to some hyperbolic quadratic form q on V , which is uniquely determined up to a scalar factor, see [7, (5.11)]. Recall from [7, (6.5)] that a right ideal $I \subset A_s$ is called *isotropic for (σ_s, f_s)* when

$$\sigma_s(I) \cdot I = \{0\} \quad \text{and} \quad f_s(I \cap \mathrm{Sym} \sigma_s) = \{0\}.$$

Isotropic ideals have the form $\mathrm{Hom}_{F_s}(V, W)$ where $W \subset V$ is a totally isotropic subspace of V for the form q , by [7, (6.6)]. Therefore, we may consider the quadric $\mathbf{Q}(\sigma, f)$ in $\mathbf{SB}(A)$ whose points are the $2n$ -dimensional right ideals in A_s that are isotropic for (σ_s, f_s) . The quadric is stable under the action of the Galois group Γ of F_s/F and is given by the equation $q = 0$ in $\mathbb{P}(V)$ under the identification $\mathbf{SB}(A) = \mathbb{P}(V)$.

Let QCSA_F^{2n} be the groupoid of central simple algebras over F of degree $2n$ with quadratic pairs and let $\mathrm{Quad}_\Gamma^{2(n-1)}$ be the groupoid of Γ -quadrics of dimension $2(n-1)$ embedded in a Γ -projective space of dimension $2n-1$. As in Proposition 2.2, we have:

Proposition 3.1. *The rule $(A, \sigma, f) \mapsto \mathbf{Q}(\sigma, f)$ induces an anti-equivalence of categories $\mathrm{QCSA}_F^{2n} \cong \mathrm{Quad}_\Gamma^{2(n-1)}$.*

The automorphism group of a quadric of dimension $2(n-1)$ over F_s or of a central simple algebra of degree $2n$ with a quadratic pair over F_s is the projective orthogonal group $\mathrm{PGO}_{2n}(F_s)$ thus Galois cohomology leads to bijections of the sets $\mathrm{Iso}(\mathrm{QCSA}_F^{2n})$ and $\mathrm{Iso}(\mathrm{Quad}_\Gamma^{2(n-1)})$ with $H^1(\Gamma, \mathrm{PGO}_{2n})$.

⁵We only consider F -linear involutions, which are sometimes called *of the first kind*, when semi-linear involutions are allowed.

Thin Γ -quadrics and quadratic étale extensions. A morphism of Γ -sets $Y_0 \xleftarrow{\pi} Y$ is called a Γ -covering if the number of elements in each fiber $y_0^{\pi^{-1}} \subset Y$ does not depend on $y_0 \in Y_0$. This number is called the *degree* of the covering. Of particular importance in the sequel are coverings of degree 2, which are also called *double coverings*. Each such covering $Y_0 \xleftarrow{\pi} Y$ defines a canonical automorphism $Y \xleftarrow{\sigma} Y$ of order 2, which interchanges the elements in each fiber of π . Clearly, this automorphism has no fixed points. We call *involution* of a Γ -set with an even number of elements any automorphism of order 2 without fixed points. If Y is a Γ -set and $Y \xleftarrow{\sigma} Y$ is an involution, the set of orbits

$$Y/\sigma = \{\{y, y^\sigma\} \mid y \in Y\}$$

is a Γ -set and the canonical map $(Y/\sigma) \leftarrow Y$ is a double covering. Thus, the notions of double covering and involution of Γ -sets are equivalent. Paraphrasing Tits [18, 7.3] we call a double Γ -covering $Y_0 \xleftarrow{\pi} Y$ with $|Y_0| = n$, resp. a Γ -set with involution (Y, σ) with $|Y| = 2n$, a *thin $2(n-1)$ -dimensional Γ -quadric*. Putting $Y = \{a_1, b_1, a_2, b_2, \dots, a_n, b_n\}$ where $a_i^\pi = b_i^\pi$ for all i , we view Y_0 as the set of pairs $Y_0 = \{\{a_1, b_1\}, \{a_2, b_2\}, \dots, \{a_n, b_n\}\}$. An *isotropic linear subvariety of dimension $k-1$* is a k -element subset of Y not containing any pair $\{a_i, b_i\}$. Thus, every isotropic linear subvariety is contained in a maximal isotropic linear subvariety, which has dimension $n-1$. The maximal isotropic linear subvarieties may be viewed as the sections of the map π ; they form a Γ -set $C(Y/Y_0)$ (or $C(Y, \sigma)$) of 2^n elements, called the *Clifford Γ -set* of Y/Y_0 , which is studied in more detail in §4.

Étale extensions are defined so as to correspond to Γ -coverings under the anti-equivalence (2.3):

Proposition 3.2. *For a homomorphism $L_0 \xrightarrow{\varepsilon} L$ of étale F -algebras, the following conditions are equivalent:*

- (a) ε endows L with a structure of free L_0 -module of rank d ;
- (b) the morphism of Γ -sets $\mathbf{X}(L_0) \xleftarrow{\mathbf{X}(\varepsilon)} \mathbf{X}(L)$ is a covering of degree d .

Proof. Let e_1, \dots, e_r be the primitive idempotents of L_0 , so that $L_0 \cong M_1 \times \dots \times M_r$ with $M_i = e_i L_0$ a finite separable field extension of F . Then $\varepsilon(e_1), \dots, \varepsilon(e_r)$ are orthogonal idempotents of L and, letting $L_i = \varepsilon(e_i)L$ we have $L \cong L_1 \times \dots \times L_r$ where each L_i is an M_i -vector space. Condition (a) holds if and only if

$$\dim_{M_1} L_1 = \dots = \dim_{M_r} L_r = d.$$

To prove (a) and (b) are equivalent, we may extend scalars to F_s and therefore assume $F = F_s = M_i$ for all i . Then $(e_i)_{i=1}^r$ is a base of L_0 , and we may identify $\mathbf{X}(L_0)$ with the dual base $(e_i^*)_{i=1}^r$. The number of elements of $\mathbf{X}(L)$ in the fiber above e_i^* is the dimension $\dim_{M_i} L_i$, so (a) and (b) are equivalent. \square

When the equivalent conditions of Proposition 3.2 hold, the map ε is injective; identifying L_0 to a subalgebra of L , we call L an *étale extension* of L_0 of degree d . Étale extensions of degree 2 are called *quadratic étale extensions*. They can also be defined by means of automorphisms, as we now show.

Let $\sigma: L \rightarrow L$ be an automorphism of order 2 of an étale F -algebra L , and let $L_0 = \text{Sym}(L, \sigma) \subset L$ denote the F -subalgebra of fixed elements, which is necessarily étale.

Proposition 3.3. *The following conditions are equivalent:*

- (a) the inclusion $L_0 \hookrightarrow L$ is a quadratic étale extension of F -algebras;
- (b) the automorphism $\mathbf{X}(\sigma)$ is an involution on $\mathbf{X}(L)$.

Proof. By the anti-equivalence (2.3) we have

$$\mathbf{X}(L_0) = \mathbf{X}(L)/\mathbf{X}(\sigma).$$

Therefore, (a) is equivalent by Proposition 3.2 to the condition that the canonical map $\mathbf{X}(L)/\mathbf{X}(\sigma) \leftarrow \mathbf{X}(L)$ is a Γ -covering of degree 2, hence also to (b). \square

When the equivalent conditions of Proposition 3.3 hold, the automorphism σ is called an *involution* of the étale F -algebra L , whose dimension is then necessarily even. Thus, the notions of quadratic étale extension and involution of étale algebras are equivalent.

For $n \geq 1$ we let QCov_Γ^n denote the groupoid whose objects are thin Γ -quadratics of dimension $2(n-1)$, i.e., double coverings $Y_0 \xleftarrow{\pi} Y$ of Γ -sets with $|Y_0| = n$, and whose morphisms are isomorphisms of Γ -coverings. Let $\mathrm{QÉtex}_F^n$ denote the groupoid of quadratic étale extensions L/L_0 of F -algebras with $\dim_F L_0 = n$. From (2.3) we obtain an anti-equivalence of groupoids

$$(3.4) \quad \mathrm{QÉtex}_F^n \cong \mathrm{QCov}_\Gamma^n.$$

Isomorphism classes of these groupoids are in bijection with $H^1(\Gamma, \mathfrak{S}_2^n \rtimes \mathfrak{S}_n)$ (see [8]). We denote by $\mathbf{Y}(L/L_0)$ or $\mathbf{Y}(L, \sigma)$ the double covering (i.e., the thin $2(n-1)$ -dimensional quadric) attached to the quadratic extension L/L_0 :

$$\mathbf{Y}(L/L_0) = \mathbf{X}(L_0) \leftarrow \mathbf{X}(L), \quad \mathbf{Y}(L, \sigma) = \mathbf{X}(L)/\mathbf{X}(\sigma) \leftarrow \mathbf{X}(L).$$

The following observation on symmetric and alternating elements in an étale algebra with involution will be used in the next subsection:

Proposition 3.5. *For any étale F -algebra of even dimension with involution (L, σ) , we have*

$$\mathrm{Sym}(L, \sigma) = \mathrm{Symd}(L, \sigma) \quad \text{and} \quad \mathrm{Skew}(L, \sigma) = \mathrm{Alt}(L, \sigma).$$

Proof. The proposition is clear if $\mathrm{char} F \neq 2$, for then

$$x = \frac{x}{2} + \sigma_L\left(\frac{x}{2}\right) \quad \text{and} \quad y = \frac{y}{2} - \sigma_L\left(\frac{y}{2}\right) \quad \text{for } x \in \mathrm{Sym}(L, \sigma) \text{ and } y \in \mathrm{Skew}(L, \sigma).$$

For the rest of the proof, we may thus assume $\mathrm{char} F = 2$. The linear map $L \rightarrow \mathrm{Symd}(L, \sigma)$ that carries $\ell \in L$ to $\ell + \sigma(\ell)$ fits into the exact sequence

$$0 \rightarrow \mathrm{Sym}(L, \sigma) \rightarrow L \rightarrow \mathrm{Symd}(L, \sigma) \rightarrow 0.$$

Since $\dim_F \mathrm{Sym}(L, \sigma) = \frac{1}{2} \dim_F L$, it follows that $\dim \mathrm{Symd}(L, \sigma) = \dim \mathrm{Sym}(L, \sigma)$. It follows that the obvious inclusion $\mathrm{Symd}(L, \sigma) \subseteq \mathrm{Sym}(L, \sigma)$ is an equality, which proves the first equation. The second equation is the same as the first since $\mathrm{char} F = 2$. \square

Embeddings of quadratics and of algebras with involution. Let A be a central simple F -algebra of even degree $2n$, and let (σ, f) be a quadratic pair on A . If (L, σ_L) is an étale F -algebra of dimension $2n$ with involution, we define an *embedding of algebras with involution*

$$\varepsilon: (L, \sigma_L) \hookrightarrow (A, \sigma, f)$$

to be an F -algebra embedding $\varepsilon: L \hookrightarrow A$ satisfying the following conditions:

- (a) $\varepsilon \circ \sigma_L = \sigma \circ \varepsilon$, and

(b) $f(x) = \text{Trd}_A(\varepsilon(\ell)x)$ for all $x \in \text{Sym}(\sigma)$ and all $\ell \in L$ such that $\ell + \sigma_L(\ell) = 1$.

Note that Proposition 3.5 shows that there exists $\ell \in L$ such that $\ell + \sigma_L(\ell) = 1$, since $1 \in \text{Sym}(\sigma_L)$. If $\ell, \ell' \in L$ satisfy

$$\ell + \sigma_L(\ell) = \ell' + \sigma_L(\ell') = 1,$$

then $\ell - \ell' \in \text{Skew}(\sigma_L)$, and Proposition 3.5 yields $\ell - \ell' \in \text{Alt}(\sigma_L)$. If (a) holds, then $\varepsilon(\ell) - \varepsilon(\ell') \in \text{Alt}(\sigma)$, hence, by [7, (2.3)],

$$\text{Trd}_A(x(\varepsilon(\ell) - \varepsilon(\ell'))) = 0 \quad \text{for all } x \in \text{Sym}(\sigma).$$

Therefore, when (a) holds, condition (b) is equivalent to

(b') there exists $\ell \in L$ such that $\ell + \sigma_L(\ell) = 1$ and $f(x) = \text{Trd}_A(\varepsilon(\ell)x)$ for all $x \in \text{Sym}(\sigma)$.

It can be further weakened to

(b'') there exists $\ell \in L$ such that $f(x) = \text{Trd}_A(\varepsilon(\ell)x)$ for all $x \in \text{Sym}(\sigma_L)$,

because this condition on ℓ implies $\varepsilon(\ell) + \sigma\varepsilon(\ell) = 1$, see [7, (5.7)]. Note that condition (b') (hence also (b'')) holds with $\ell = \frac{1}{2}$ if $\text{char } F \neq 2$.

Example 3.6. Let (L, σ_L) be an arbitrary étale F -algebra with involution of the first kind. Let $L_0 = \text{Sym}(\sigma_L) \subset L$ be the fixed subalgebra under σ_L , let $T_0: L_0 \rightarrow F$ be the trace, and let $t: L \rightarrow F$ be the quadratic form given by $t(x) = T_0(\sigma(x)x)$ for $x \in L$. Let (σ_t, f_t) denote the adjoint quadratic pair on $\text{End}_F L$. The regular representation is an embedding

$$(L, \sigma_L) \hookrightarrow (\text{End}_F L, \sigma_t, f_t).$$

Indeed, condition (a) holds because the polar form of t is

$$b_t(x, y) = T_{L_0/F}(\sigma_L(x)y + \sigma_L(y)x) \quad \text{for } x, y \in L,$$

and we have

$$b_t(xy, z) = b_t(y, \sigma_L(x)z) \quad \text{for } x, y, z \in L.$$

Condition (b) also is easily checked, see [7, (5.12)].

Proposition 3.7. *Use the same notation as in Example 3.6. Suppose L is split and let (A, σ, f) be a central simple F -algebra with quadratic pair. For any embedding*

$$\varepsilon: (L, \sigma_L) \hookrightarrow (A, \sigma, f)$$

there is an isomorphism $\varphi: (A, \sigma, f) \rightarrow (\text{End}_F L, \sigma_t, f_t)$ such that $\varphi \circ \varepsilon: (L, \sigma_L) \hookrightarrow (\text{End}_F L, \sigma_t, f_t)$ is the regular representation.

Proof. Let $e_1, \dots, e_n, e'_1, \dots, e'_n$ be the collection of primitive idempotents of L , labelled in such a way that $e'_i = \sigma_L(e_i)$ for $i \in \mathbf{n}$. As observed in the proof of Proposition 2.4, we may identify $A = \text{End}_F V$ and we have a decomposition into 1-dimensional subspaces

$$V = V_1 \oplus \dots \oplus V_n \oplus V'_1 \oplus \dots \oplus V'_n$$

such that $\varepsilon(e_i)$ (resp. $\varepsilon(e'_i)$) is the projection $V \rightarrow V_i$ (resp. $V \rightarrow V'_i$). Let $q: V \rightarrow F$ be a quadratic form such that (σ, f) is adjoint to q , and let b be the polar form of q . Since σ is adjoint to b and $\sigma\varepsilon(e_i) = \varepsilon\sigma_L(e_i) = \varepsilon(e'_i)$, we have for all $i, j \in \mathbf{n}$

$$b(V_i, V_j) = b(\varepsilon(e_i)(V_i), V_j) = b(V_i, \varepsilon(e'_i)(V_j)) = \{0\}.$$

Similarly, $b(V'_i, V'_j) = \{0\}$ for all $i, j \in \mathbf{n}$, and $b(V_i, V'_j) = \{0\}$ if $i \neq j$. Since b is nonsingular, it follows that $b(V_i, V'_i) = F$ for all $i \in \mathbf{n}$, hence we may find $v_i \in V_i$, $v'_i \in V'_i$ such that

$$b(v_i, v'_i) = 1.$$

Let $S \in \text{End}_F V$ be the map such that $S(v_i) = v'_i$ and $S(v'_i) = v_i$ for all $i \in \mathbf{n}$. It is readily verified that for $i \in \mathbf{n}$

$$S \circ \varepsilon(e'_i): x \mapsto v_i b(v_i, x) \quad \text{and} \quad S \circ \varepsilon(e_i): x \mapsto v'_i b(v'_i, x) \quad \text{for all } x \in V.$$

Therefore, by [7, (5.11)],

$$q(v_i) = f(S \circ \varepsilon(e'_i)) \quad \text{and} \quad q(v'_i) = f(S \circ \varepsilon(e_i)).$$

If $\ell \in L$ is such that $f(x) = \text{Tr}(\varepsilon(\ell) \circ x)$ for all $x \in \text{Sym}(\sigma)$, then

$$f(S \circ \varepsilon(e'_i)) = \text{Tr}(\varepsilon(\ell) \circ S \circ \varepsilon(e'_i)) = \text{Tr}(S \circ \varepsilon(e'_i \ell)).$$

None of the base vectors v_1, \dots, v'_n is mapped to a nonzero multiple of itself by $S \circ \varepsilon(e'_i \ell)$, hence $\text{Tr}(S \circ \varepsilon(e'_i \ell)) = 0$ and therefore $q(v_i) = 0$. Likewise, $q(v'_i) = 0$ for $i \in \mathbf{n}$. Therefore, mapping $v_i \mapsto e_i$ and $v'_i \mapsto e'_i$ for $i \in \mathbf{n}$ defines an isometry $(V, q) \rightarrow (L, t)$, and the induced isomorphism of algebras with quadratic pairs $\varphi: (\text{End}_F V, \sigma, f) \rightarrow (\text{End}_F L, \sigma_t, f_t)$ is such that $\varphi \circ \varepsilon$ is the regular representation. \square

We now turn to the geometric counterpart of the notion of embeddings of algebras with involution. Let $Q \subset P$ be a Γ -quadric of dimension $2(n-1)$ in a Γ -projective space of dimension $2n-1$ over a field F . We let $C(Q)$ denote the Γ -set of $(n-1)$ -dimensional linear varieties that lie in Q , and call this Γ -set the *Clifford Γ -set* of Q . If (Y, σ) is a thin Γ -quadric of dimension $2(n-1)$, we define an embedding of quadrics

$$\varepsilon: (Y, \sigma) \hookrightarrow Q$$

as a general embedding $\varepsilon: Y \hookrightarrow P$ that carries isotropic linear subvarieties of (Y, σ) to isotropic linear subvarieties of P , i.e., to linear subvarieties of Q . In particular, the image of Y lies on Q . Since every isotropic linear subvariety is contained in a maximal isotropic linear subvariety, it suffices to require that the induced map $Gr_{n-1}\varepsilon: Gr_{n-1}Y \hookrightarrow Gr_{n-1}(P)$ carry the set of maximal isotropic linear subvarieties $C(Y, \sigma)$ to $C(Q)$. We let $C(\varepsilon)$ denote the Γ -equivariant map induced by $Gr_{n-1}\varepsilon$,

$$C(\varepsilon): C(Y, \sigma) \hookrightarrow C(Q).$$

Let (L, σ_L) be an arbitrary étale F -algebra of dimension $2n$ with involution of the first kind, and let

$$\varepsilon: (L, \sigma_L) \hookrightarrow (A, \sigma, f)$$

be an embedding in a central simple F -algebra of degree $2n$ with quadratic pair. Since $\varepsilon: L \hookrightarrow A$ is an F -algebra embedding, we may consider the general embedding of Proposition 2.6:

$$\varepsilon_*: \mathbf{X}(L) \hookrightarrow \mathbf{SB}(A).$$

Let $\mathbf{Y}(L, \sigma_L)$ be the thin Γ -quadric associated with (L, σ_L) and $\mathbf{Q}(A, \sigma, f)$ the Γ -quadric associated with (A, σ, f) .

Proposition 3.8. *The map ε_* defines an embedding of quadrics*

$$\varepsilon_*: \mathbf{Y}(L, \sigma_L) \hookrightarrow \mathbf{Q}(A, \sigma, f).$$

Proof. As usual, let $A_s = A \otimes_F F_s$, and let (σ_s, f_s) be the quadratic pair on A_s extending (σ, f) . We have to show that for every $\omega \in C(\mathbf{Y}(L, \sigma_L))$ we have $Gr_{n-1}\varepsilon_*(\omega) \in C(Q(\sigma, f))$. Let $\xi, \eta \in \mathbf{X}(L)$. For $x \in \varepsilon_*(\xi)$, $y \in \varepsilon_*(\eta)$ and $\ell \in L$ we have

$$(1 \otimes \xi(\ell)) \cdot x = (\varepsilon(\ell) \otimes 1) \cdot x \quad \text{and} \quad (1 \otimes \eta(\ell)) \cdot y = (\varepsilon(\ell) \otimes 1) \cdot y,$$

hence also

$$\sigma_s(x) \cdot (\varepsilon(\ell) \otimes 1) = \sigma_s(x) \cdot (1 \otimes \xi\sigma_L(\ell)),$$

and

$$\sigma_s(x) \cdot y \cdot (1 \otimes \eta(\ell)) = \sigma_s(x) \cdot (\varepsilon(\ell) \otimes 1) \cdot y = \sigma_s(x) \cdot y \cdot (1 \otimes \xi\sigma_L(\ell)).$$

If $\eta \neq \xi\sigma_L$, we can find $\ell \in L$ such that $\eta(\ell) \neq \xi\sigma_L(\ell)$, hence the last equation shows that $\sigma_s(x) \cdot y = 0$. Therefore,

$$\sigma_s(\varepsilon_*(\xi)) \cdot \varepsilon_*(\eta) = \{0\}, \quad \text{unless } \eta = \xi\sigma_L.$$

In particular, for every $\omega \in C(\mathbf{Y}(L, \sigma_L))$,

$$(3.9) \quad \sigma_s\left(\bigoplus_{\xi \in \omega} \varepsilon_*(\xi)\right) \cdot \bigoplus_{\xi \in \omega} \varepsilon_*(\xi) = \{0\}, \quad \text{i.e.,} \quad \sigma_s(Gr_{n-1}\varepsilon_*(\omega)) \cdot Gr_{n-1}\varepsilon_*(\omega) = \{0\}.$$

Now, let $\ell \in L$ be such that $f(x) = \text{Tr}(\varepsilon(\ell)x)$ for all $x \in \text{Sym}(\sigma)$. For $x \in Gr_{n-1}\varepsilon_*(\omega) \cap \text{Sym}(\sigma_s)$ we have

$$(3.10) \quad f_s(x) = \text{Tr}((\varepsilon(\ell) \otimes 1) \cdot x) \quad \text{since } x \in \text{Sym}(\sigma_s),$$

and

$$(\varepsilon(\ell) \otimes 1) \cdot x \in Gr_{n-1}\varepsilon_*(\omega) \quad \text{since } x \in Gr_{n-1}\varepsilon_*(\omega).$$

On the other hand, we have $x = \sigma_s(x) \in \sigma_s(Gr_{n-1}\varepsilon_*(\omega))$, hence by (3.9),

$$x \cdot (\varepsilon(\ell) \otimes 1) \cdot x = 0.$$

It follows that $((\varepsilon(\ell) \otimes 1) \cdot x)^2 = 0$, hence $\text{Tr}((\varepsilon(\ell) \otimes 1) \cdot x) = 0$ and (3.10) shows that $f_s(x) = 0$. Thus, $Gr_{n-1}\varepsilon_*(\omega)$ is isotropic for all $\omega \in C(\mathbf{Y}(L, \sigma_L))$, and the proposition is proved. \square

Conversely, every embedding

$$\varepsilon: (Y, \varphi) \hookrightarrow Q(\sigma, f)$$

of a Γ -set Y of $2n$ elements with an involution φ of the first kind in the quadric associated with a quadratic pair on a central simple algebra A of degree $2n$ arises from the embedding of the étale F -algebra with involution $\mathbf{M}(Y, \varphi)$ into (A, σ, f) , as we now show.

Let $\varepsilon(Y) = \{I_1, \dots, I_n, I'_1, \dots, I'_n\}$, where the isotropic $2n$ -dimensional right ideals I_1, \dots, I'_n of A_s are indexed in such a way that $I'_i = \varepsilon(\xi^\varphi)$ if $\varepsilon(\xi) = I_i$. As in §2, let

$$L_s = \{x \in A_s \mid x \cdot I_i \subset I_i \text{ and } x \cdot I'_i \subset I'_i \text{ for all } i \in \mathbf{n}\}$$

and $L = L_s^\Gamma$. Proposition 2.7 shows that we may identify X with $\mathbf{X}(L)$ in such a way that ε is the general embedding $\mathbf{X}(L) \hookrightarrow \mathbf{SB}(A)$ induced by the inclusion $L \hookrightarrow A$. Let σ_L be the involution on L such that $\mathbf{X}(\sigma_L) = \varphi$.

Proposition 3.11. *The inclusion $L \hookrightarrow A$ is an embedding $(L, \sigma_L) \hookrightarrow (A, \sigma, f)$.*

Proof. The primitive idempotents $e_1, \dots, e_n, e'_1, \dots, e'_n$ of L_s are in one-to-one correspondence with $I_1, \dots, I_n, I'_1, \dots, I'_n$ in such a way that $I_i = e_i \cdot A_s$ and $I'_i = e'_i \cdot A_s$ for $i \in \mathbf{n}$. The indexing of I_1, \dots, I'_n is such that σ_{L_s} interchanges e_i and e'_i for $i \in \mathbf{n}$. To check that σ restricts to σ_L , observe that I'_i is the only element among $\{I_1, \dots, I'_n\}$ that does not belong to any $\omega^{C(\varepsilon)}$ for any $\omega \in C(X, *)$ containing i . Therefore,

$$\sigma_s(I_i) \cdot I_j = \sigma_s(I_i) \cdot I'_k = \{0\} \quad \text{for all } j, k \in \mathbf{n} \text{ with } k \neq i,$$

hence for all $i \in \mathbf{n}$

$$\sigma_s(e_i) \cdot e_j = \sigma_s(e_i) \cdot e'_k = 0 \quad \text{for all } j, k \in \mathbf{n} \text{ with } k \neq i.$$

Similarly, for all $i \in \mathbf{n}$,

$$\sigma_s(e'_i) \cdot e_j = \sigma_s(e'_i) \cdot e'_k = 0 \quad \text{for all } j, k \in \mathbf{n} \text{ with } j \neq i.$$

Since $e_1 + \dots + e_n + e'_1 + \dots + e'_n = 1$, it follows that for all $i \in \mathbf{n}$ we have

$$e'_i = \left(\sum_{j \in \mathbf{n}} \sigma_s(e_j) + \sigma_s(e'_j) \right) \cdot e'_i = \sigma_s(e_i) \cdot e'_i$$

and, on the other hand,

$$\sigma_s(e_i) \cdot (1 - e'_i) = \sigma_s(e_i) \cdot \left(\sum_{j \in \mathbf{n}} e_j + \sum_{k \in \mathbf{n} \setminus \{i\}} e'_k \right) = 0,$$

hence $\sigma_s(e_i) = \sigma_s(e_i) \cdot e'_i$. Therefore, $\sigma_s(e_i) = e'_i$ for $i \in \mathbf{n}$.

To complete the proof that $(L, \sigma_L) \hookrightarrow (A, \sigma, f)$, it remains to show that for all $x \in \text{Sym}(\sigma)$ and all $\ell \in L$ such that $\ell + \sigma_L(\ell) = 1$ we have $f(x) = \text{Trd}_A(\ell x)$. To show that this equation holds for all $x \in \text{Sym}(\sigma_s)$ and all $\ell \in L_s$ such that $\ell + \sigma_s(\ell) = 1$, it actually suffices to prove it for a single ℓ , as was observed in the equivalence of conditions (b) and (b') defining an embedding. Let $\ell = \sum_{i \in \mathbf{n}} e_i$. Since $\ell + \sigma_s(\ell) = 1$ we have for all $y \in A_s$

$$\text{Trd}_{A_s}(\ell \cdot (y + \sigma_s(y))) = \text{Trd}_{A_s}((\ell + \sigma_s(\ell)) \cdot y) = \text{Trd}_{A_s}(y).$$

Since $f_s(y + \sigma_s(y)) = \text{Trd}_{A_s}(y)$, we have

$$(3.12) \quad f_s(x) = \text{Trd}_{A_s}(\ell x) \quad \text{for all } x \in \text{Symd}(\sigma_s).$$

Let $I = \ell \cdot A_s = I_1 + \dots + I_n$ and $I' = (1 - \ell) \cdot A_s = I'_1 + \dots + I'_n$. These right ideals are isotropic since $\{I_1, \dots, I_n\}, \{I'_1, \dots, I'_n\} \in C(\varepsilon(X, *))$. For $x \in I \cap \text{Sym}(\sigma_s)$ we have $f_s(x) = 0$ and, on the other hand, $\ell x \in I$ and $x \ell x \in \sigma_s(I) \cdot I = \{0\}$, hence $(\ell x)^2 = 0$. Therefore,

$$(3.13) \quad f_s(x) = \text{Trd}_{A_s}(\ell x) \quad \text{for all } x \in I \cap \text{Sym}(\sigma_s).$$

Similarly, for $x \in I' \cap \text{Sym}(\sigma_s)$ we have $\ell x = 0$, hence

$$(3.14) \quad f_s(x) = \text{Trd}_{A_s}(\ell x) \quad \text{for all } x \in I' \cap \text{Sym}(\sigma_s).$$

Now, for $x \in \text{Sym}(\sigma_s)$ we have

$$x = \ell x(1 - \ell) + (1 - \ell)x\ell + \ell x \ell + (1 - \ell)x(1 - \ell).$$

Since $\ell \in I$, $1 - \ell \in I'$, and $\sigma_s(\ell) = 1 - \ell$, we have

$$\ell x(1 - \ell) \in I \cap \text{Sym}(\sigma_s), \quad (1 - \ell)x\ell \in I' \cap \text{Sym}(\sigma_s),$$

and

$$\ell x \ell + (1 - \ell)x(1 - \ell) = \ell x \ell + \sigma_s(\ell x \ell) \in \text{Symd}(\sigma_s).$$

Therefore,

$$f_s(x) = f_s(\ell x(1 - \ell)) + f_s((1 - \ell)x\ell) + f_s(\ell x\ell + (1 - \ell)x(1 - \ell)),$$

and it follows from (3.12), (3.13), and (3.14) that

$$f_s(x) = \text{Trd}_{A_s}(\ell x).$$

□

4. CLIFFORD ALGEBRAS AND Γ -SETS

As in the previous sections, we let F denote an arbitrary field and Γ the Galois group of a separable closure F_s of F .

Clifford constructions. Recall from §3 that for any $2(n - 1)$ -dimensional thin Γ -quadric Y/Y_0 we let $C(Y/Y_0)$ denote the Γ -set of maximal linear isotropic subvarieties, which can be identified with the 2^n -element set of sections of the covering $Y_0 \leftarrow Y$. We call $C(Y/Y_0)$ the *Clifford Γ -set* of Y/Y_0 . The canonical involution σ of Y/Y_0 induces an involution $\underline{\sigma}$ on $C(Y/Y_0)$, which maps every section to its complement in Y . If $n = 1$, then $C(Y/Y_0) = Y$ and $\underline{\sigma} = \sigma$. The following proposition is clear:

Proposition 4.1. *If Y/Y_0 and Z/Z_0 are thin Γ -quadrics, then $(Y \sqcup Z)/(Y_0 \sqcup Z_0)$ is a thin Γ -quadric and there is a canonical isomorphism*

$$C(Y/Y_0) \times C(Z/Z_0) \xrightarrow{\sim} C(Y \sqcup Z/Y_0 \sqcup Z_0)$$

which maps any pair of isotropic linear varieties (ω, ω') to $\omega \sqcup \omega'$.

We now turn to the analogue of this construction for quadratic étale extensions. Let $\varepsilon: L_0 \hookrightarrow L$ be a quadratic étale extension of étale F -algebras, with $\dim_F L_0 = n$. In view of the Grothendieck correspondence there exists an étale algebra $C(L/L_0)$ of dimension 2^n , with involution $\underline{\sigma}$, such that $C \mathbf{Y}(L/L_0) = \mathbf{Y}(C(L/L_0), \underline{\sigma})$. This étale algebra is constructed explicitly in [8], where it is designated by $\Omega(L/L_0)$. We briefly recall the construction: the algebra $L_0^{\otimes n}$ contains an idempotent $e_n^{L_0}$ that corresponds under the composition of canonical isomorphisms

$$L_0^{\otimes n} \xrightarrow{\sim} \mathbf{M}(\mathbf{X}(L_0^{\otimes n})) \xrightarrow{\sim} \mathbf{M}(\mathbf{X}(L_0)^n)$$

to the characteristic function χ of the complement of the “fat diagonal” of $\mathbf{X}(L_0)^n$:

$$\chi(\xi_1, \dots, \xi_n) = \begin{cases} 1 & \text{if } \xi_1, \dots, \xi_n \text{ are all different;} \\ 0 & \text{if } \xi_i = \xi_j \text{ for some } i, j \text{ with } i \neq j. \end{cases}$$

Then, letting $(L^{\otimes n})^{\mathfrak{S}_n}$ denote the subalgebra of $L^{\otimes n}$ fixed under the action of \mathfrak{S}_n by permutation of the factors, we set

$$C(L/L_0) = \varepsilon^{\otimes n}(e_n^{L_0}) \cdot (L^{\otimes n})^{\mathfrak{S}_n}.$$

We call the algebra $C(L/L_0)$ the *Clifford algebra of L/L_0* . If $n = 1$, then $C(L/L_0) = L$, and the canonical involution $\underline{\sigma}$ is the canonical involution of L/L_0 .

Proposition 4.2. *If L/L_0 and M/M_0 are quadratic étale extensions, there is a canonical isomorphism*

$$P: C(L \times M/L_0 \times M_0) \xrightarrow{\sim} C(L/L_0) \otimes C(M/M_0),$$

which corresponds under the Grothendieck correspondence to the canonical isomorphism of Proposition 4.1.

By contrast to Γ -sets, we have an F -linear map

$$c_L: L \rightarrow C(L/L_0)$$

defined by

$$(4.3) \quad c_L(x) = \varepsilon^{\otimes n}(e_n^{L_0}) \cdot ((x \otimes 1 \otimes \cdots \otimes 1) + (1 \otimes x \otimes \cdots \otimes 1) + (1 \otimes \cdots \otimes 1 \otimes x))$$

for $x \in L$.

Proposition 4.4. *Let L/L_0 and M/M_0 be quadratic étale extensions and let $\rho: L \times M \rightarrow L \otimes M$ be the linear map given by $\rho(x, y) = x \otimes 1 + 1 \otimes y$. The diagram*

$$\begin{array}{ccc} C(L \times M/L_0 \times M_0) & \xrightarrow{P} & C(L/L_0) \otimes C(M/M_0) \\ \uparrow c_{L \times M} & & \uparrow c_L \otimes c_M \\ L \times M & \xrightarrow[\rho]{} & L \otimes M \end{array}$$

is commutative.

Proof. The claim follows easily from the definition of the canonical map P . \square

Corollary 4.5. *The space $c_L(L)$ generates $C(L/L_0)$ as an algebra.*

Proof. By extending F we may assume that L/L_0 has a nontrivial decomposition $L/L_0 = M/M_0 \times N/N_0$. The claim then follows from (4.4) by induction on the dimension. \square

Lemma 4.6. *Let $L_0 = \text{Sym}(\sigma_L)$ and $T_0: L_0 \rightarrow F$ be the trace map. The kernel of $c_L: L \rightarrow C(L, \sigma_L)$ is*

$$\ker c_L = \{x \in L_0 \mid T_0(x) = 0\}.$$

Proof. Extend scalars to F_s and use induction on $\dim L$. \square

Discriminants. Let Q be a $2(n-1)$ -dimensional Γ -quadric. The Clifford Γ -set $C(Q)$, consisting of maximal linear subvarieties of Q , decomposes into two classes $C_1(Q), C_2(Q)$: two linear varieties v_1, v_2 lie in the same class if and only if $\dim(v_1 \cap v_2) \equiv n-1 \pmod{2}$, see for example [3, §§II.6]. The equivalence relation is clearly preserved under the action of Γ , hence Γ acts on the set

$$\Delta(Q) := \{C^1(Q), C^2(Q)\},$$

which we call the *discriminant* of Q . We let $\delta: C(Q) \rightarrow \Delta(Q)$ denote the Γ -equivariant map that carries each linear variety in $C(Q)$ to its class.

Likewise, if (Y, σ) is a thin Γ -quadric of dimension $2(n-1)$, the Clifford Γ -set $C(Y, \sigma)$ of $(n-1)$ -dimensional isotropic linear subvarieties decomposes into two classes as for “classical” quadrics, hence we also have a discriminant $\Delta(Y, \sigma)$ and a canonical map $\delta: C(Y, \sigma) \rightarrow \Delta(Y, \sigma)$. An alternative description of this discriminant is given in [9, Proposition 2.5]: it is canonically isomorphic to the set of orbits under the alternating group \mathfrak{A}_{2n} of the set

$$\Sigma_{2n}(Y) = \{(y_1, \dots, y_{2n}) \mid Y = \{y_1, \dots, y_{2n}\}\}.$$

Therefore, $\Delta(Y, \sigma)$ depends only on Y , and not on the involution σ ; we may thus use the notation $\Delta(Y)$ for $\Delta(Y, \sigma)$.

Proposition 4.7. *Let $\varepsilon: (Y, \sigma) \hookrightarrow Q$ be an embedding of quadrics. There is an induced isomorphism of Γ -sets $\Delta(\varepsilon): \Delta(Y) \rightarrow \Delta(Q)$ that fits into the following commutative diagram:*

$$\begin{array}{ccc} C(Y, \sigma) & \xrightarrow{C(\varepsilon)} & C(Q) \\ \delta \downarrow & & \downarrow \delta \\ \Delta(Y) & \xrightarrow{\Delta(\varepsilon)} & \Delta(Q). \end{array}$$

Proof. For $\omega_1, \omega_2 \in C(Y, \sigma)$, we have

$$|\omega_1 \cap \omega_2| = \dim(C(\varepsilon)(\omega_1) \cap C(\varepsilon)(\omega_2)),$$

hence the map $C(\varepsilon)$ is compatible with the division of $C(Y, \sigma)$ and $C(Q)$ into two classes. Therefore, we may define $\Delta(\varepsilon)$ by mapping the equivalence class of $\omega \in C(Y, \sigma)$ to the equivalence class of $C(\varepsilon)(\omega) \in C(Q)$. \square

Under the Grothendieck correspondence, the discriminant of Γ -sets yields a discriminant of étale algebras, see [8, §18]. If (L, σ) is an étale algebra with involution, we have a canonical map

$$\delta: \Delta(L) \rightarrow C(L, \sigma),$$

so we may consider the Clifford algebra $C(L, \sigma)$ as an étale extension of $\Delta(L)$.

Clifford algebra embeddings. Let (A, σ, f) be a central simple algebra of even degree $2n$ with quadratic pair over an arbitrary field F . Recall from [7, §8] that the Clifford algebra $C(A, \sigma, f)$ is defined in such a way that if $A = \text{End}_F V$ and (σ, f) is induced by some quadratic form $q: V \rightarrow F$, then $C(A, \sigma, f)$ is the even Clifford algebra

$$C(A, \sigma, f) = C_0(V, q).$$

The center of $C(A, \sigma, f)$ is a quadratic étale F -algebra $Z(A, \sigma, f)$, which is related to the discriminant of (σ, f) , see [7, (8.10)], and $C(A, \sigma, f)$ is an Azumaya algebra of constant rank 2^n over $Z(A, \sigma, f)$. The Clifford algebra $C(A, \sigma, f)$ is defined as a quotient of the tensor algebra $T(A)$ of the underlying vector space of A , hence there is a canonical F -linear map

$$c_A: A \rightarrow C(A, \sigma, f).$$

In the split case $A = \text{End}_F V$, the bilinear polar form of q yields an F -linear isomorphism $V \rightarrow V^*$ which allows us to identify

$$V \otimes_F V = V \otimes_F V^* = \text{End}_F V;$$

under this identification, the map c_A carries $v \otimes w \in V \otimes_F V$ to $v \cdot w \in C_0(V, q)$. Now, let $\deg A = 2n$, and let

$$\varepsilon: (L, \sigma_L) \hookrightarrow (A, \sigma, f)$$

be an embedding of an étale F -algebra with involution of the first kind of dimension $2n$ in (A, σ, f) .

Theorem 4.8. *There is an F -algebra embedding*

$$(4.9) \quad C(\varepsilon): C(L, \sigma_L) \hookrightarrow C(A, \sigma, f)$$

such that the following diagram commutes:

$$(4.10) \quad \begin{array}{ccc} L & \xrightarrow{\varepsilon} & A \\ c_L \downarrow & & \downarrow c_A \\ C(L, \sigma_L) & \xrightarrow{C(\varepsilon)} & C(A, \sigma, f). \end{array}$$

The F -algebra homomorphism $C(\varepsilon)$ is uniquely determined by this property. Moreover, the following diagram commutes:

$$(4.11) \quad \begin{array}{ccc} C(L, \sigma_L) & \xrightarrow{C(\varepsilon)} & C(A, \sigma, f) \\ \underline{\sigma}_L \downarrow & & \downarrow \underline{\sigma} \\ C(L, \sigma_L) & \xrightarrow{C(\varepsilon)} & C(A, \sigma, f). \end{array}$$

Furthermore, there is an F -algebra isomorphism $\overline{C(\varepsilon)}$ which makes the following diagram commute:

$$\begin{array}{ccc} C(L, \sigma_L) & \xrightarrow{C(\varepsilon)} & C(A, \sigma, f) \\ \delta \uparrow & & \uparrow \\ \Delta(L) & \xrightarrow{\overline{C(\varepsilon)}} & Z(A, \sigma, f). \end{array}$$

The proof uses the following general principle:

Principle 4.12. *Let C and C' be F -algebras, and let $U \subset C$ be a subspace which generates C as an F -algebra. Let also $f: U \rightarrow C'$ be a linear map. If there exists a field extension K/F and a K -algebra homomorphism $\hat{f}_K: C \otimes_F K \rightarrow C' \otimes_F K$ such that $\hat{f}_K|_{U \otimes K} = f \otimes \text{Id}_K$, then f extends to an F -algebra homomorphism $\hat{f}: C \rightarrow C'$.*

Proof. The restriction of \hat{f}_K to C is an F -algebra homomorphism $C \rightarrow C' \otimes_F K$. Since the restriction of \hat{f}_K to U is f , and U generates C , the image of this homomorphism is in fact in C' . \square

Proof of Theorem 4.8. Since $c(L)$ generates the algebra $C(L, \sigma_L)$, it is clear that, if it exists, $C(\varepsilon)$ is uniquely determined by the commutativity of diagram (4.10). By [7, (8.14)], we have

$$\ker(c_A \circ \varepsilon) = \{x \in L \mid \varepsilon(x) \in \text{Sym}(\sigma) \text{ and } f(\varepsilon(x)) = 0\}.$$

Since ε embeds (L, σ_L) in (A, σ, f) , the condition $\varepsilon(x) \in \text{Sym}(\sigma_L)$ holds if and only if $x \in L_0$, and then

$$f(\varepsilon(x)) = \text{Trd}_A(\varepsilon(\ell x)) = T(\ell x) \quad \text{for any } \ell \in L \text{ such that } \ell + \sigma_L(\ell) = 1.$$

Since for $x \in L_0$ we have

$$T(\ell x) = T_0(\ell x + \sigma_L(\ell x)) = T_0(x),$$

it follows from Lemma 4.6 that $\ker c_L = \ker(c_A \circ \varepsilon)$. Therefore, ε induces an injective F -linear map $\varepsilon': c_L(L) \hookrightarrow C(A, \sigma, f)$. Since $c_L(L)$ generates $C(L, \sigma_L)$, we may apply the general principle above with $U = c_L(L)$, $C = C(L, \sigma_L)$, and $C' = C(A, \sigma, f)$, taking for K a separable closure F_s of F . To prove the existence of $C(\varepsilon)$, it suffices to show that $\varepsilon'_s := \varepsilon \otimes \text{Id}_{F_s}$ extends to an F_s -algebra embedding

$$C(\varepsilon_s): C(L_s, \sigma_{L_s}) \rightarrow C(A_s, \sigma_s, f_s).$$

By Proposition 3.7, we may assume $A_s = \text{End}_{F_s}(L_s)$, (σ_s, f_s) is the quadratic pair adjoint to the quadratic form $t: L_s \rightarrow F_s$ of Example 3.6, and ε_s is the regular representation, hence $C(A_s, \sigma_s, f_s) = C_0(L_s, t)$.

We argue by induction on n . If $n = 1$ (i.e. $L_s = F_s \times F_s$) the map c_L is the identity

$$c_L: L_s \xrightarrow{\sim} C(L_s, \sigma_{L_s}) = L_s,$$

and $c_A \circ \varepsilon_s: L_s \rightarrow C_0(L_s, t)$ is an F_s -algebra isomorphism, so the result is clear.

If $n > 1$ we may find a nontrivial decomposition $(L_s, \sigma_s) = (L_1, \sigma_1) \times (L_2, \sigma_2)$. We then have, by Proposition 4.2,

$$C(L_s, \sigma_s) = C(L_1, \sigma_1) \otimes_F C(L_2, \sigma_2),$$

and, letting $c_i: L \rightarrow C(L_i, \sigma_i)$ for $i = 1, 2$ denote the canonical F_s -linear map,

$$c_L(\ell_1, \ell_2) = c_1(\ell_1) \otimes c_2(\ell_2) \quad \text{for } \ell_1 \in L_1 \text{ and } \ell_2 \in L_2.$$

On the other hand, defining quadratic forms t_1 and t_2 on L_1 and L_2 respectively in the same way as t on L_s , we have an orthogonal sum decomposition

$$(L_s, t) = (L_1, t_1) \bigoplus^\perp (L_2, t_2),$$

hence a canonical F -algebra embedding

$$C_0(L_1, t_1) \otimes_F C_0(L_2, t_2) \hookrightarrow C_0(L_s, t).$$

By induction, we may assume $(c \circ \varepsilon)_1: L_1 \rightarrow C_0(L_1, t_1)$ and $(c \circ \varepsilon)_2: L_2 \rightarrow C_0(L_2, t_2)$ extend to F_s -algebra embeddings

$$C(\varepsilon_1): C(L_1, \sigma_1) \hookrightarrow C_0(L_1, t_1) \quad \text{and} \quad C(\varepsilon_2): C(L_2, \sigma_2) \hookrightarrow C_0(L_2, t_2).$$

Then the composition

$$C(L_s, \sigma_{L_s}) = C(L_1, \sigma_1) \otimes_F C(L_2, \sigma_2) \xrightarrow{C(\varepsilon_1) \otimes C(\varepsilon_2)} C_0(L_1, t_1) \otimes_F C_0(L_2, t_2) \hookrightarrow C_0(L_s, t)$$

is an F_s -algebra embedding that extends $c_A \circ \varepsilon_s$.

To prove that diagram (4.11) commutes, it suffices to prove the commutativity of

$$\begin{array}{ccc} c_L(L) & \xrightarrow{C(\varepsilon)} & c_A(A) \\ \sigma_L \downarrow & & \downarrow \sigma \\ c_L(L) & \xrightarrow{C(\varepsilon)} & c_A(A) \end{array}$$

since $c_L(L)$ generates $C(L, \sigma_L)$. Commutativity of the latter diagram follows from commutativity of the diagrams

$$\begin{array}{ccccc} L & \xrightarrow{\varepsilon} & A & L & \xrightarrow{c_L} & C(L, \sigma_L) & A & \xrightarrow{c_A} & C(A, \sigma, f) \\ \sigma_L \downarrow & & \downarrow \sigma & \sigma_L \downarrow & & \downarrow \sigma_L & \text{and } \sigma \downarrow & & \downarrow \sigma \\ L & \xrightarrow{\varepsilon} & A, & L & \xrightarrow{c_L} & C(L, \sigma_L), & A & \xrightarrow{c_A} & C(A, \sigma, f). \end{array}$$

To complete the proof, it remains to show that $C(\varepsilon)$ maps $\Delta(L)$ to $Z(A, \sigma, f)$. It suffices to see that the image of $\Delta(L)$ centralizes $c_A(A)$ since $c_A(A)$ generates $C(A, \sigma, f)$. Again, we may extend scalars to reduce to the case where L is split, and use induction on $\dim L$. Details are left to the reader. \square

Remark 4.13. The situation described in Theorem 4.8 is related to results about Cartan subspaces given in [17]. Assume that F has characteristic different from 2. A *Cartan subspace* of a central simple algebra A of even degree $2n$ with an orthogonal involution σ is by definition a subspace of $\text{Skew}(A, \sigma_A)$ such that there exists an element $a \in H$ which is invertible in A , is separable (i.e., generates an étale F -subalgebra of A) and such that $a^{-1}H$ is an étale commutative subalgebra of A of dimension $2n$. Equivalently, if $a \in \text{Skew}(A, \sigma_A)$ is invertible and separable in A , then $H = \{x \in \text{Skew}(A, \sigma_A) \mid xa = ax\}$ is a Cartan subspace for a . It can be shown that the algebra $L_0 = a^{-1}H$ is independent of the choice of the element a . Moreover the subalgebra $L = L_0 \oplus H$ of A is maximal commutative étale in A and has an involution σ_L which is the restriction of σ_A to L . Thus the set of skew-symmetric elements in L is a Cartan subspace of A . It follows from [17, Théorème 3] that $c_A(L)$ generates a maximal commutative étale subalgebra of $C(A, \sigma_A)$. This algebra is isomorphic to $C(L, \sigma_L)$.

The Severi–Brauer variety of a Clifford algebra. We discuss in this subsection the geometric analogue of Theorem 4.8, showing that an embedding of a thin quadric into a quadric induces an embedding of the corresponding Clifford Γ -sets. Recall that the Severi–Brauer variety of a central simple algebra is a Γ -projective space. To take in account the fact that $C(A, \sigma, f)$ is an Azumaya algebra of constant rank over the étale F -algebra $Z(A, \sigma, f)$ we introduce a variant of the Severi–Brauer variety for Azumaya algebras over quadratic étale algebras.⁶ We consider disjoint unions $P_1 \sqcup P_2$ of two projective F -spaces of the same dimension, endowed with a continuous action of a profinite group Γ by collineations that may interchange the two components. We call these unions *double Γ -projective spaces*. They are equipped with a canonical structure map to a Γ -set Z of two elements

$$Z := \{\{P_1\}, \{P_2\}\} \leftarrow P_1 \sqcup P_2.$$

If $\dim P_1 = \dim P_2 = n - 1$, we may define for every integer $k \in \mathbf{n}$ two different sets:

$$\begin{aligned} gr_{k-1}(P_1 \sqcup P_2) &= Gr_{k-1}(P_1) \cup Gr_{k-1}(P_2) \quad \text{and} \\ Gr_{k-1}(P_1 \sqcup P_2) &= \{v_1 \sqcup v_2 \mid v_1 \in Gr_{k-1}(P_1), v_2 \in Gr_{k-1}(P_2)\}. \end{aligned}$$

Note that, even though $Gr_{k-1}(P_1)$ and $Gr_{k-1}(P_2)$ are not Γ -sets since they may be interchanged under the Γ -action, Γ naturally acts on $gr_{k-1}(P_1 \sqcup P_2)$ and $Gr_{k-1}(P_1 \sqcup P_2)$, and $gr_{k-1}(P_1 \sqcup P_2)$ carries a canonical Γ -equivariant map

$$Z \leftarrow gr_{k-1}(P_1 \sqcup P_2).$$

If $\Gamma_0 \subset \Gamma$ is the subgroup fixing Z , we clearly have an isomorphism of Γ_0 -sets

$$Gr_{k-1}(P_1 \sqcup P_2) = Gr_{k-1}(P_1) \times Gr_{k-1}(P_2).$$

If $\Gamma_0 \neq \Gamma$, then $Gr_{k-1}(P_1 \sqcup P_2)$ may be identified with the norm of the Γ_0 -set $Gr_{k-1}(P_1)$ (or $Gr_{k-1}(P_2)$) following Ferrand’s general definition in [4].

For $(Y \xleftarrow{\pi} X)$ a Γ -covering with $|Y| = 2$ and $|X| = 2n$, a *general embedding*

$$\varepsilon: (Y \xleftarrow{\pi} X) \hookrightarrow (Z \leftarrow P_1 \sqcup P_2)$$

is a pair of Γ -equivariant maps $\varepsilon: X \rightarrow gr_0(P_1 \sqcup P_2)$, $\bar{\varepsilon}: Y \rightarrow Z$ such that

⁶For the sake of simplicity we give an ad-hoc construction.

(a) the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{\varepsilon} & \text{gr}_0(P_1 \sqcup P_2) \\ \pi \downarrow & & \downarrow \\ Y & \xrightarrow{\bar{\varepsilon}} & Z, \end{array}$$

(b) $\bar{\varepsilon}$ is bijective, and

(c) the image of each fiber of Y is a set of n points in general position in P_1 and P_2 respectively.

The embedding ε thus induces Γ -equivariant maps

$$\text{Gr}_{k-1}\varepsilon: \text{Gr}_{k-1}(X/Y) \hookrightarrow \text{gr}_{k-1}(P_1 \sqcup P_2) \quad \text{for all } k \in \mathbf{n}.$$

These maps fit into commutative diagrams

$$\begin{array}{ccc} \text{Gr}_{k-1}(X/Y) & \xrightarrow{\lambda^k(\varepsilon)} & \text{gr}_{k-1}(P_1 \sqcup P_2) \\ \downarrow & & \downarrow \\ Y & \xrightarrow{\bar{\varepsilon}} & Z. \end{array}$$

Double Γ -projective space arise from the following construction. Let Z be a quadratic étale algebra over an arbitrary field F , and let B be an Azumaya Z -algebra of constant rank. As usual, let F_s denote a separable closure of F , let $Z_s = Z \otimes_F F_s$ and $B_s = B \otimes_F F_s$ the F_s -algebras obtained by scalar extension, and let Γ be the Galois group of F_s/F . If $\dim_F B = 2n^2$, then $B_s \simeq M_n(F_s) \times M_n(F_s)$; its minimal right ideals have dimension n . We let $\mathbb{S}\mathbb{B}(B)$ be the double Γ -projective space whose points are the n -dimensional right ideals of B_s , with the natural Γ -action. These ideals are divided into two classes, which correspond to the two primitive idempotents of Z_s : to each n -dimensional right ideal I , we associate the primitive idempotent $e_I \in Z_s$ such that $e_I \cdot I = I$. Alternatively, we associate to I the element $I^\delta \in \mathbf{X}(Z)$ such that $I^\delta(e_I) = 1$. Thus, the canonical structure map of the double Γ -projective space $\mathbb{S}\mathbb{B}(B)$ can be identified with

$$\mathbf{X}(Z) \xleftarrow{\delta} \mathbb{S}\mathbb{B}(B).$$

Now, let $K \xrightarrow{i} L$ be an étale extension of degree n of a quadratic étale F -algebra. (Thus, $\dim_F L = 2n$.) By definition, an embedding

$$\varepsilon: (L/K) \hookrightarrow (B/Z)$$

is an F -algebra embedding $L \hookrightarrow B$ that restricts to an isomorphism $\bar{\varepsilon}: K \xrightarrow{\sim} Z$.

Proposition 4.14. *If, in the situation above, L is a split étale F -algebra, then for every embedding $\varepsilon: (L/K) \hookrightarrow (B/Z)$ there is an F -algebra isomorphism $\varphi: B \xrightarrow{\sim} \text{End}_K L$ whose restriction to Z is $\bar{\varepsilon}^{-1}$ such that $\varphi \circ \varepsilon: L \hookrightarrow \text{End}_K L$ is the regular representation.*

Proof. By hypothesis K is split, hence Z also. Using $\bar{\varepsilon}$ as an identification, let

$$K = Z = F \times F,$$

and consider the corresponding decompositions $L = L_1 \times L_2$, $B = B_1 \times B_2$, and $\varepsilon = (\varepsilon_1, \varepsilon_2)$. By Proposition 2.4, we may find F -algebra isomorphisms $\varphi_i: B_i \xrightarrow{\sim} \text{End}_F L_i$, for $i = 1, 2$, such that $\varphi_i \circ \varepsilon_i: L_i \hookrightarrow \text{End}_F L_i$ is the regular representation. The proposition holds with $\varphi = (\varphi_1, \varphi_2): B \rightarrow \text{End}_F L_1 \times \text{End}_F L_2 = \text{End}_K L$. \square

Now, consider an arbitrary embedding $\varepsilon: (L/K) \hookrightarrow (B/Z)$ of an étale extension of degree n . For every $\xi \in \mathbf{X}(L)$ we let as in(2.5)

$$\varepsilon_*(\xi) = \{x \in B_s \mid (1 \otimes \xi(\ell)) \cdot x = (\varepsilon(\ell) \otimes 1) \cdot x \text{ for all } \ell \in L\}.$$

Proposition 4.15. *The right ideal $\varepsilon_*(\xi)$ is n -dimensional, and ε_* defines a general embedding*

$$\varepsilon_*: (\mathbf{X}(K) \xleftarrow{\mathbf{X}(i)} \mathbf{X}(L)) \hookrightarrow (\mathbf{X}(Z) \leftarrow \mathbb{S}\mathbb{B}(B)).$$

Conversely, every general embedding of a Γ -covering of degree n with 2-element base in $\mathbb{S}\mathbb{B}(B)$ arises from an embedding of an étale extension in B .

Proof. The proof of the first (resp. second) part is similar to that of Proposition 2.6 (resp. 2.7). \square

Proposition 4.15 applies in particular in the situation of Theorem 4.8: for the rest of this subsection, suppose A is a central simple F -algebra of degree $2n$ with a quadratic pair (σ, f) . If (L, σ_L) is an étale F -algebra with involution of dimension $2n$ and

$$\varepsilon: (L, \sigma_L) \hookrightarrow (A, \sigma, f)$$

is an embedding of algebras with involution, then Theorem 4.8 yields an F -algebra embedding of the étale extension $C(L, \sigma_L)/\Delta(L)$ into $C(A, \sigma, f)/Z(A, \sigma, f)$. By Proposition 4.15 there is a general embedding

$$(4.16) \quad C(\varepsilon)_*: \mathbf{X}C(L, \sigma_L) \hookrightarrow \mathbb{S}\mathbb{B}(C(A, \sigma, f)).$$

On the other hand, the embedding ε also yields by Proposition 3.8 an embedding of quadrics

$$(4.17) \quad \varepsilon_*: \mathbf{Y}(L, \sigma_L) \hookrightarrow Q(\sigma, f).$$

To complete the picture, we now relate the Clifford Γ -set $C(Q(\sigma, f))$ to the Severi-Brauer double projective space $\mathbb{S}\mathbb{B}(C(A, \sigma, f))$.

Recall from [7, §8] that there is a Clifford bimodule $B(A, \sigma, f)$ over $C(A, \sigma, f)$, which is equipped with a canonical F -linear map $b: A \rightarrow B(A, \sigma, f)$, see [7, §9]. In the split case $A = \text{End}_F V$, we have $B(A, \sigma, f) = V \otimes_F C_1(V, q)$. The bilinear polar form of q yields an F -linear isomorphism $V \rightarrow V^*$ which allows us to identify

$$V \otimes_F V = V \otimes_F V^* = \text{End}_F V;$$

under this identification, the map b carries $v \otimes w \in V \otimes_F V$ to $v \otimes w \in V \otimes_F C_1(V, q)$. Since $\deg A = 2n$, we have $\dim_F C(A, \sigma, f) = 2^{2n-1}$. For every $2n$ -dimensional isotropic right ideal $I \subset A_s$, let $\psi(I) \subset C(A_s, \sigma_s, f_s)$ denote the right annihilator of the F -vector space $(\sigma_s(I))^b \subset B(A_s, \sigma_s, f_s)$, i.e.,

$$\psi(I) = \{x \in C(A_s, \sigma_s, f_s) \mid (\sigma_s(I))^b \cdot x = 0 \text{ for all } y \in I\}.$$

Proposition 4.18. *The map ψ defines an embedding*

$$\psi: (\Delta(Q(\sigma, f)) \leftarrow C(Q(\sigma, f))) \hookrightarrow (\mathbf{X}(Z(A, \sigma, f)) \leftarrow \mathbb{S}\mathbb{B}(C(A, \sigma, f))).$$

Proof. We first show $\dim_F \psi(I) = 2^{n-1}$. Let $A_s = \text{End}_{F_s} V$ for some $2n$ -dimensional F_s -vector space V . The quadratic pair (σ_s, f_s) is adjoint to some hyperbolic quadratic form q on V , and $I = \text{Hom}_{F_s}(V, U)$ for some totally isotropic n -dimensional subspace $U \subset V$. We may therefore identify V with $U \oplus U^*$, and q with the quadratic form $q(u, \varphi) = \varphi(u)$ for $u \in U$, $\varphi \in U^*$. Then the full Clifford algebra of q

can be identified with the algebra $\text{End}_{F_s}(\bigwedge U)$ of F_s -linear endomorphisms of the exterior algebra of U , in such a way that the canonical map

$$V = U \oplus U^* \rightarrow C(V, q) = \text{End}_{F_s}(\bigwedge U)$$

maps $u \in U$ to the left exterior multiplication by u and $\varphi \in U^*$ to its associated derivation d_φ , see [7, (8.3)]. Let

$$\bigwedge_0 U = \bigoplus_{i \text{ even}} \bigwedge^i U \subset \bigwedge U \quad \text{and} \quad \bigwedge_1 U = \bigoplus_{i \text{ odd}} \bigwedge^i U \subset U.$$

Then

$$C(A_s, \sigma_s, f_s) = C_0(V, q) = \text{End}_{F_s}(\bigwedge_0 U) \times \text{End}_{F_s}(\bigwedge_1 U)$$

and

$$B(A_s, \sigma_s, f_s) = V \otimes_{F_s} C_1(V, q) = V \otimes_{F_s} (\text{Hom}_{F_s}(\bigwedge_0 U, \bigwedge_1 U) \times \text{Hom}_{F_s}(\bigwedge_1 U, \bigwedge_0 U)).$$

Under the identification $A_s = \text{End}_{F_s} V = V \otimes_{F_s} V$ we have $I = U \otimes_{F_s} V$, hence

$$(\sigma_s(I))^b = V \otimes_{F_s} U \subset V \otimes_{F_s} C_1(V, q).$$

Therefore, $\psi(I)$ consists of the linear maps $f \in \text{End}_{F_s}(\bigwedge U)$ such that $f(\bigwedge_0 U) \subset \bigwedge_0 U$, $f(\bigwedge_1 U) \subset \bigwedge_1 U$, and

$$u \wedge f(x) = 0 \quad \text{for all } u \in U \text{ and all } x \in \bigwedge U.$$

The last condition implies $f(x) \in \bigwedge^n U$, hence

$$\psi(I) = \begin{cases} \text{Hom}_{F_s}(\bigwedge_0 U, \bigwedge^n U) & \text{if } n \text{ is even,} \\ \text{Hom}_{F_s}(\bigwedge_1 U, \bigwedge^n U) & \text{if } n \text{ is odd.} \end{cases}$$

It follows that $\dim_{F_s} \psi(I) = \dim_{F_s} \bigwedge_0 U = \dim_{F_s} \bigwedge_1 U = 2^{n-1}$, hence $\psi(I) \in \mathbb{S}\mathbb{B}(C(A, \sigma, f))$. Moreover, for every nonzero $\varphi \in U^*$ we have $d_\varphi(\bigwedge^n U) \neq \{0\}$, hence

$$\{v \in V \mid v \cdot f = 0 \text{ in } C(V, q) \text{ for all } f \in \psi(I)\} = U.$$

It follows that

$$I = \{y \in A \mid (\sigma(y))^b \cdot x = 0 \text{ for all } x \in \psi(I)\},$$

hence the map ψ is injective.

It is clear from the definition that the map ψ is Γ -equivariant. To complete the proof, it remains to show that ψ induces an isomorphism

$$\bar{\psi}: \Delta(Q(\sigma, f)) \xrightarrow{\sim} X(Z(A, \sigma, f)).$$

As above, we identify $A_s = \text{End}_{F_s} V$ for some $2n$ -dimensional F_s -vector space V . Let $I, I' \in C(Q(\sigma, f))$, so

$$I = \text{Hom}_{F_s}(V, U), \quad I' = \text{Hom}_{F_s}(V, U')$$

for some totally isotropic n -dimensional subspaces $U, U' \subset V$. By a theorem of Witt, there is an orthogonal transformation $g \in \text{O}(V, q)$ such that $g(U) = U'$. This transformation induces an automorphism \hat{g} of $C_0(V, q)$ such that $\hat{g}(\psi(I)) = \psi(I')$. If e (resp. e') is the primitive central idempotent of $C_0(V, q)$ such that $e \cdot \psi(I) = \psi(I)$ (resp. $e' \cdot \psi(I') = \psi(I')$), we therefore have $\hat{g}(e) = e'$. Now, we have $\dim(U \cap U') \equiv n \pmod{2}$ if and only if $g \in \text{O}^+(V, q)$ by [3, §§ II.6, II.10], and this condition holds if and only if the restriction of \hat{g} to the center of $C_0(V, q)$ is the identity, by [7, (13.2)]. Therefore, $e = e'$ if and only if the ideals I and I' belong to the same class of $C(Q(\sigma, f))$. This completes the proof. \square

Corollary 4.19. *If (σ, f) is a hyperbolic quadratic pair on A , then the Clifford algebra $C(A, \sigma, f)$ decomposes into a direct product of simple F -algebras $C(A, \sigma, f) \simeq C_1 \times C_2$, and one of the factors C_1, C_2 is split.*

Proof. If (σ, f) is hyperbolic, then its discriminant is trivial, hence $Z(A, \sigma, f) \simeq F \times F$ and $C(A, \sigma, f)$ decomposes into a direct product of simple F -algebras as above. Moreover, there is a maximal isotropic linear variety in $Q(\sigma, f)$ that is defined over F , hence fixed under the action of Γ . Its image under ψ is a point on $\mathbb{S}\mathbb{B}(C(A, \sigma, f)) = \mathbf{S}\mathbb{B}(C_1) \sqcup \mathbf{S}\mathbb{B}(C_2)$ fixed under the action of Γ . Its existence shows that C_1 or C_2 is split. \square

Remark 1. This corollary was first proved by D.C. Van Drooge and H.P. Allen, see also Tits [17, Proposition 8].

We may now relate the maps (4.16) and (4.17) associated to an embedding ε of algebras with involution:

Proposition 4.20. *The following diagram is commutative:*

$$(4.21) \quad \begin{array}{ccc} C\mathbf{Y}(L, \sigma_L) & \xrightarrow{C(\varepsilon_*)} & C(Q(\sigma, f)) \\ \parallel & & \downarrow \psi \\ \mathbf{X}C(L, \sigma_L) & \xrightarrow{C(\varepsilon)_*} & \mathbb{S}\mathbb{B}(C(A, \sigma, f)). \end{array}$$

Proof. Extending scalars, we reduce to the case where L is split, $A = \text{End}_F L$, and ε is the regular representation as in Example 3.6 (see Proposition 3.7). Let $\omega \in C\mathbf{Y}(L, \sigma_L)$, and let $e_1, \dots, e_n, e'_1, \dots, e'_n$ be the primitive idempotents of L , labelled in such a way that $\sigma(e_i) = e'_i$ for $i \in \mathbf{n}$ and $\xi(e_i) = 1, \xi(e'_i) = 0$ for $\xi \in \omega$. Under the identification $C\mathbf{Y}(L, \sigma_L) = \mathbf{X}C(L, \sigma_L)$, the corresponding F -algebra homomorphism $\xi_\omega: C(L, \sigma_L) \rightarrow F$ satisfies

$$\xi_\omega(c_L(e_i)) = 1 \quad \text{and} \quad \xi_\omega(c_L(e'_i)) = 0 \quad \text{for all } i \in \mathbf{n}.$$

In $C(A, \sigma, f) = C_0(L, t)$ we have $\varepsilon(e_i) = e_i e'_i$ and $\varepsilon(e'_i) = e'_i e_i$ for $i \in \mathbf{n}$, hence

$$C(\varepsilon)_*(\xi_\omega) = \{x \in C_0(L, t) \mid e_i e'_i x = x \text{ and } e'_i e_i x = 0 \text{ for all } i \in \mathbf{n}\}.$$

On the other hand, $C(\varepsilon_*)(\omega) = \text{Hom}_F(L, \bigoplus_{i \in \mathbf{n}} e_i L)$, hence

$$\psi(C(\varepsilon_*)(\omega)) = \{x \in C_0(L, t) \mid e_i \cdot x = 0 \text{ for } i \in \mathbf{n}\}.$$

In $C(L, t)$, we have $e_i^2 = 0$ and $e_i e'_i + e'_i e_i = 1$, hence $e_i e'_i e_i = e_i$ for $i \in \mathbf{n}$. Therefore, if $e'_i e_i x = 0$, then $e_i x = 0$. Conversely, if $e_i x = 0$, then $e'_i e_i x = 0$ and

$$x = (e_i e'_i + e'_i e_i)x = e_i e'_i x.$$

Therefore, $\psi(C(\varepsilon_*)(\omega)) = C(\varepsilon)_*(\omega)$. \square

5. DYNKIN DIAGRAMS AND SEVERI–BRAUER VARIETIES

The projective linear group PGL_n has Dynkin diagram of type A_{n-1} and the group \mathfrak{S}_n is its Weyl group. The special orthogonal group PGO_{2n}^+ , with Weyl group $\mathfrak{S}_2^{n-1} \times \mathfrak{S}_n$, has Dynkin diagram of type D_n . Accordingly we call groupoids occurring in Proposition 2.2 and (2.3) *groupoids of type A_{n-1}* , and groupoids occurring in (3.4) and Proposition 3.1 *groupoids of type D_n* . Many properties of these groupoids

are related to properties of the corresponding Dynkin diagrams. After introducing oriented quadrics and étale algebras, we briefly consider two examples: the equivalence $A_3 = D_3$ and triality.

Special quadrics. A *special or oriented* Γ -quadric is a pair (Q, ∂) , where Q is Γ -quadric (or a thin Γ -quadric) and ∂ is a fixed isomorphism of Γ -sets $\mathbf{2} \xleftarrow{\sim} \Delta(Q)$. In particular the Γ -action on $\Delta(Q)$ is trivial. There are two possible choices for ∂ . A choice is an *orientation of* Q . If Q is oriented we may use the given isomorphism $\partial: \mathbf{2} \xleftarrow{\sim} \Delta(Q)$ to decompose the Clifford Γ -set:

$$C(Z/Z_0) = C_1(Z/Z_0, \partial) \sqcup C_2(Z/Z_0, \partial)$$

where

$$C_i(Z/Z_0, \partial) = \{\omega \in C(Z/Z_0) \mid \omega^{\delta\partial} = i\} \quad \text{for } i = 1, 2.$$

Similarly, we define *oriented extensions of étale F -algebras* as pairs $(L/L_0, \partial)$ where L/L_0 is an extension of étale F -algebras and ∂ is an orientation of L , i.e., a fixed isomorphism $\partial: \Delta(L) \xrightarrow{\sim} F \times F$. We also have *oriented central simple algebras with quadratic pairs*, defined as 4-tuples (A, σ, f, ∂) , where ∂ is a fixed isomorphism $\partial: Z(A, \sigma, f) \xrightarrow{\sim} F \times F$. In the corresponding decomposition

$$C(A, \sigma, f, \partial) = C_1(A, \sigma, f, \partial) \times C_2(A, \sigma, f, \partial),$$

$C_1(A, \sigma, f, \partial)$ and $C_2(A, \sigma, f, \partial)$ are central simple algebras over F . The classes of morphisms considered in the previous sections can be restricted to preserve the orientations. For example an embedding of oriented quadrics

$$\varepsilon: (Y, \sigma, \partial_Y) \hookrightarrow (Q, \partial_Q)$$

is an embedding $\varepsilon: Y \hookrightarrow Q$ such that $\Delta(\varepsilon) \circ \partial_Q = \partial_Y$. The diagram (4.21) restricts to a commutative diagram

$$\begin{array}{ccc} C_1 \mathbf{Y}(L/L_0, \partial_L) & \xrightarrow{C_1(\varepsilon_*)} & C_1(Q(\sigma, f), \partial_Q) \\ \parallel & & \downarrow \psi_1 \\ \mathbf{X} C_1(L/L_0, \partial_L) & \xrightarrow{C_1(\varepsilon)_*} & \mathbf{SB}(C_1(A, \sigma, f, \partial)). \end{array}$$

As in 2.2, 2.3, 3.4 and 3.1, we have anti-equivalences of groupoids

$$\text{OQCSA}_F^{2n} \equiv \text{OQuad}_\Gamma^{2(n-1)} \quad \text{and} \quad \text{OQÉtex}_F^n \equiv \text{OQCov}_\Gamma^n,$$

using the following notations:

- (1) OQCSA_F^{2n} for the groupoid of oriented central simple algebras over F of degree $2n$ with quadratic pairs,
- (2) $\text{OQuad}_\Gamma^{2(n-1)}$ for the groupoid of oriented Γ -quadrics of dimension $2(n-1)$ over F ,
- (3) OQÉtex_F^n for the groupoid of oriented quadratic étale extensions L/L_0 of F -algebras with $\dim_F L_0 = n$.
- (4) OQCov_Γ^n for the groupoid of oriented thin Γ -quadrics of dimension $2(n-1)$ over F , i.e., double Γ -coverings Y/Y_0 with $|Y_0| = n$.

Isomorphism classes are in bijection with $H^1(\Gamma, \text{PGO}_{2n}^+)$, resp. $H^1(\Gamma, \mathfrak{S}_2^{n-1} \times \mathfrak{S}_n)$.

The exceptional isomorphism $A_3 = D_3$. The fact that the Dynkin diagrams A_3 and D_3 are identical can be interpreted at the level of the corresponding groupoids. For any 4-element Γ -set X , the Γ -set $Gr_1(X)$ of 2-element subsets of X is equipped with a canonical involution σ , which maps each 2-element subset of X to its complement. Thus, it defines a functor $\text{Set}_\Gamma^4 \rightarrow \text{QCov}_\Gamma^3$, which we also designate by Gr_1 . The sections $\{S_1, S_2, S_3\} \in C(Gr_1(X), \sigma)$ (where S_1, S_2, S_3 are 2-element subsets of X such that $\{\sigma(S_1), \sigma(S_2), \sigma(S_3)\} \cap \{S_1, S_2, S_3\} = \emptyset$) naturally fall into two classes, depending on whether $S_1 \cap S_2 \cap S_3$ is empty or is a 1-element subset of X . Therefore, there is a natural orientation on $Gr_1(X)$, and we may view Gr_1 as a functor $\text{Set}_\Gamma^4 \rightarrow \text{OQCov}_\Gamma^3$.

The analogue of Gr_1 for étale algebras under the Grothendieck correspondence is an “exterior algebra” functor $\lambda^2: \acute{\text{E}}t_F^4 \rightarrow \text{OQ}\acute{\text{E}}\text{tex}_F^3$ studied in [8]. There is also an “exterior algebra” functor $\lambda^2: \text{CSA}_F^4 \rightarrow \text{OQCSA}_F^6$ discussed in [7, §15D].

Proposition 5.1. *The functors Gr_1 (resp. λ^2) and C_1 induce equivalences:*


$$\begin{array}{ccc} \text{Set}_\Gamma^4 & \xrightleftharpoons[C_1]{Gr_1} & \text{OQCov}_\Gamma^3, & \acute{\text{E}}t_F^4 & \xrightleftharpoons[C_1]{\lambda^2} & \text{OQ}\acute{\text{E}}\text{tex}_F^3, \\ \text{Proj}_\Gamma^3 & \xrightleftharpoons[C_1]{Gr_1} & \text{OQuad}_\Gamma^4, & \text{CSA}_F^4 & \xrightleftharpoons[C_1]{\lambda^2} & \text{OQCSA}_F^6. \end{array}$$

Moreover these equivalences are compatible with general embeddings.

Proof. The (equivalent) first two cases are discussed in [8], the third is equivalent to the last and the last follows from [7, (15.32)]. \square

Remark 5.2. The equivalence between groupoids of thin objects of type A_3 and of type D_3 is induced at the cohomological level by the well-known isomorphism $\mathfrak{S}_4 \xrightarrow{\sim} \mathfrak{S}_2^2 \rtimes \mathfrak{S}_3$. Classically, for any separable field extension E of degree 4, the subalgebra of $\lambda^2 E$ fixed under the involution is the *cubic resolvent* (see [8]). On the geometric side, the quadric $Gr_1(P)$ can be viewed as a 4-dimensional quadric parametrised by a 3-dimensional projective space. This quadric is known as *Klein’s quadric*.

Triality. All Dynkin diagrams but one admit at most automorphisms of order two, which are related to duality in algebra and geometry. The Dynkin diagram of D_4 :



is special, in the sense that it admits automorphisms of order 3. Algebraic and geometric objects related to D_4 are of particular interest as they also usually admit exceptional automorphisms of order 3, which are called trialitarian. This is the case for the four groupoids OQCov_Γ^4 , $\text{OQ}\acute{\text{E}}\text{tex}_F^4$, OQuad_Γ^6 and OQCSA_F^8 of type D_4 . For (E, ∂_E) in $\text{OQ}\acute{\text{E}}\text{tex}_F^4$ or in OQCSA_F^8 , the objects $C_1(E, \partial_E)$ and $C_2(E, \partial_E)$ belong to $\text{Q}\acute{\text{E}}\text{tex}_F^4$, resp. to QCSA_F^8 (see [9], resp. [7, §35]). Corresponding results hold for objects in OQCov_Γ^4 or OQuad_Γ^6 since these groupoids of quadrics are anti-equivalent to the corresponding groupoids of algebras. Isomorphism classes of these groupoids are in bijection with $H^1(\Gamma, \text{PGO}_8^+)$ or $H^1(\Gamma, \mathfrak{S}_2^3 \rtimes \mathfrak{S}_4)$. The groups PGO_8^+ and $\mathfrak{S}_2^3 \rtimes \mathfrak{S}_4$ admit up to inner automorphisms one outer action of the cyclic group of

order 3 (this is well known for PGO_3^+) and is in [5] for $\mathfrak{S}_2^3 \rtimes \mathfrak{S}_4$). Thus the cyclic group of order 3 acts on the corresponding sets of isomorphism classes. In fact these automorphisms of order 3 are induced by automorphisms that can already be defined at the level of the groupoids. Let D_4 be any of the four groupoids above, let \tilde{D}_4 be the corresponding groupoid of unoriented objects, and let $\mathcal{F}: (E, \partial_E) \mapsto E$ be the forgetful functor $D_4 \rightarrow \tilde{D}_4$.

Theorem 5.3. *The functors C_1, C_2 factor through the forgetful functor, i.e., there are functors*

$$C_1^+, C_2^+: D_4 \rightarrow D_4$$

such that $\mathcal{F} \circ C_i^+ = C^i$ for $i = 1, 2$. These functors satisfy natural equivalences:

$$(C_1^+)^3 = \text{Id}, \quad (C_1^+)^2 = C_2^+$$

and induce the trialitarian action on the sets of isomorphism classes.

Proof. The claim for thin objects is in [9] (under the hypothesis that $\text{char}(F) \neq 2$) and the classical case (for algebras) is in [7, §35]. \square

Observe that triality is compatible with general embeddings. Triality for thin objects is extensively discussed in [9]. A very nice presentation of classical (geometric and algebraic) triality is in [20].

REFERENCES

- [1] R. Brusamarello, P. Chuard-Koulmann, and J. Morales. Orthogonal groups containing a given maximal torus. *J. Algebra*, 266:87–101, 2003.
- [2] P. Deligne. *Séminaire de géométrie algébrique du Bois-Marie, SGA 4 1/2, Cohomologie étale, avec la collaboration de J. F. Boutot, A. Grothendieck, L. Illusie et J. L. Verdier*, volume 569 of *Lecture Notes in Mathematics*. Springer-Verlag, 1977.
- [3] J. Dieudonné. *La géométrie des groupes classiques*. Seconde édition, revue et corrigée. Springer-Verlag, Berlin, 1963.
- [4] D. Ferrand. Un foncteur norme. *Bull. Soc. math. France*, 126:1–49, 1998.
- [5] W. N. Franzsen and R. B. Howlett. Automorphisms of nearly finite Coxeter groups. *Adv. Geom.*, 3(3):301–338, 2003.
- [6] J. Jahnel. The Brauer-Severi variety associated with a central simple algebra: A survey. Preprint 52, Linear Algebraic Groups and Related Structures Preprint Server, <http://www.mathematik.uni-bielefeld.de/LAG/man/052.html>, 2000.
- [7] M-A. Knus, A. A. Merkurjev, M. Rost, and J-P. Tignol. *The Book of Involutions*. Number 44 in American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, R.I., 1998. With a preface in French by J. Tits.
- [8] M-A. Knus and J-P. Tignol. Quartic exercises. *Inter. J. Math. Math. Sci.*, 2003:4263–4323, 2003.
- [9] M-A. Knus and J-P. Tignol. Triality and étale algebras. In: J.-L. Colliot-Thélène, S. Garibaldi, R. Sujatha, and V. Suresh, editors. *Quadratic forms, linear algebraic groups, and cohomology*, Developments in Mathematics. Springer, 2010.
- [10] M-A. Knus and J-P. Tignol. Severi-Brauer varieties over the field of characteristic one. In preparation, 2010.
- [11] S. Mac Lane. *Categories for the Working Mathematician*, volume 5 of Graduate Texts in Mathematics. Springer-Verlag, 2nd ed., 1998.
- [12] A. S. Merkurjev. Closed points of Severi-Brauer varieties. *Vestnik S.-Peterburg. Univ. Mat. Mekh. Astronom.*, (vyp. 2):51–53, 124, 1993.
- [13] G. Prasad and A. S. Rapinchuk. Local-global principles for embedding of fields with involution into simple algebras with involution. *Commentarii mathematici Helvetici* 85, 2010 (to appear).
- [14] D. J Saltman. Triality, cocycles, crossed products, involutions, Clifford algebras and invariants, 88 p. Austin.

- [15] J-P. Serre. *Corps locaux*. Number VIII in Publications de l'Institut de Mathématique de l'Université de Nancago. Hermann, Paris, 1968.
- [16] J. Tits. Sur les analogues algébriques des groupes semi-simples complexes. In *Colloque d'algèbre supérieure, tenu à Bruxelles du 19 au 22 décembre 1956*, pages 261–289. Centre Belge de Recherches Mathématiques, Établissements Ceuterick, Louvain; Librairie Gauthier-Villars, Paris, 1957.
- [17] J. Tits. Formes quadratiques, groupes orthogonaux et algèbres de Clifford. *Invent. Math.*, 5:19–41, 1968.
- [18] J. Tits. *Buildings of Spherical Type and Finite BN-Pairs*, volume 386 of *Lecture Notes in Mathematics*. Springer-Verlag, 1974.
- [19] J. Tits. Sur les produits tensoriels de deux algèbres de quaternions. *Bull. Soc. Math. Belg. Sér. B* **45** (1993), no. 3, 329–331.
- [20] F. van der Blij and T.A. Springer. Octaves and triality. *Nieuw Arch. Wiskd., III. Ser.*, 8:158–169, 1960.

DEPARTMENT MATHEMATIK, ETH ZENTRUM, CH-8092 ZÜRICH, SWITZERLAND
E-mail address: knus@math.ethz.ch

INSTITUT DE MATHÉMATIQUE PURE ET APPLIQUÉE, UNIVERSITÉ CATHOLIQUE DE LOUVAIN, B-1348
LOUVAIN-LA-NEUVE, BELGIUM
E-mail address: jean-pierre.tignol@uclouvain.be