

ON A CONJECTURE OF ERDŐS AND SIMONOVITS:
EVEN CYCLESPETER KEEVASH*, BENNY SUDAKOV†,
JACQUES VERSTRAËTE‡

Received July 24, 2011

Let \mathcal{F} be a family of graphs. A graph is \mathcal{F} -free if it contains no copy of a graph in \mathcal{F} as a subgraph. A cornerstone of extremal graph theory is the study of the *Turán number* $\text{ex}(n, \mathcal{F})$, the maximum number of edges in an \mathcal{F} -free graph on n vertices. Define the *Zarankiewicz number* $\text{z}(n, \mathcal{F})$ to be the maximum number of edges in an \mathcal{F} -free bipartite graph on n vertices. Let C_k denote a cycle of length k , and let \mathcal{C}_k denote the set of cycles C_ℓ , where $3 \leq \ell \leq k$ and ℓ and k have the same parity. Erdős and Simonovits conjectured that for any family \mathcal{F} consisting of bipartite graphs there exists an odd integer k such that $\text{ex}(n, \mathcal{F} \cup \mathcal{C}_k) \sim \text{z}(n, \mathcal{F})$ – here we write $f(n) \sim g(n)$ for functions $f, g: \mathbb{N} \rightarrow \mathbb{R}$ if $\lim_{n \rightarrow \infty} f(n)/g(n) = 1$. They proved this when $\mathcal{F} = \{C_4\}$ by showing that $\text{ex}(n, \{C_4, C_5\}) \sim \text{z}(n, C_4)$. In this paper, we extend this result by showing that if $\ell \in \{2, 3, 5\}$ and $k > 2\ell$ is odd, then $\text{ex}(n, \mathcal{C}_{2\ell} \cup \{C_k\}) \sim \text{z}(n, \mathcal{C}_{2\ell})$. Furthermore, if $k > 2\ell + 2$ is odd, then for infinitely many n we show that the extremal $\mathcal{C}_{2\ell} \cup \{C_k\}$ -free graphs are bipartite incidence graphs of generalized polygons. We observe that this exact result does not hold for any odd $k < 2\ell$, and furthermore the asymptotic result does not hold when (ℓ, k) is $(3, 3)$, $(5, 3)$ or $(5, 5)$. Our proofs make use of pseudorandomness properties of nearly extremal graphs that are of independent interest.

1. Introduction

Given a family \mathcal{F} of graphs, a graph is \mathcal{F} -free if it contains no copy of a graph in \mathcal{F} as a subgraph. The *Turán number* $\text{ex}(n, \mathcal{F})$ is the maximum number of

Mathematics Subject Classification (2000): 05C35, 05C38, 05C50, 05D40

* Research supported in part by ERC grant 239696 and EPSRC grant EP/G056730/1.

† Research supported in part by NSF grant DMS-1101185 and by USA-Israeli BSF grant.

‡ Research supported in part by an Alfred P. Sloan Research Fellowship and NSF Grant DMS-0800704.

edges in an \mathcal{F} -free graph on n vertices. The *Zarankiewicz number* $z(n, \mathcal{F})$ is the maximum number of edges in an \mathcal{F} -free *bipartite* graph on n vertices.

When $\mathcal{F} = \{F\}$ consists of a single forbidden graph we denote the Turán number by $\text{ex}(n, F)$. A classical theorem of Turán [39] gives an exact result for $\text{ex}(n, K_t)$, where K_t is the complete graph on t vertices: the unique largest K_t -free graph on n vertices is the complete $(t-1)$ -partite graph with part sizes as equal as possible. In general, Erdős, Stone and Simonovits [14,15] showed that $\text{ex}(n, \mathcal{F}) \sim (1 - 1/r) \binom{n}{2}$, where $r = \min\{\chi(F) - 1 : F \in \mathcal{F}\}$. Here we write $f(n) \sim g(n)$ for functions $f, g: \mathbb{N} \rightarrow \mathbb{R}$ if $\lim_{n \rightarrow \infty} f(n)/g(n) = 1$. This determines the Turán number asymptotically when \mathcal{F} consists of non-bipartite graphs. However, much less is known concerning the Turán numbers of bipartite graphs. There is no bipartite graph F containing a cycle such that $\text{ex}(n, F)$ is known exactly for all n . In fact, even the order of magnitude of $\text{ex}(n, F)$ is not known for quite simple bipartite graphs, such as the complete bipartite graph with four vertices in each part, the cycle of length eight, and the three-dimensional cube graph.

In this paper, we study the effect of forbidding short odd cycles on bipartite extremal problems, in particular, the extremal problem for even cycles. Let C_k denote a cycle of length k and let \mathcal{C}_k denote the set of cycles C_ℓ where $3 \leq \ell \leq k$ and ℓ and k have the same parity. Bondy and Simonovits [9] showed that $\text{ex}(n, \mathcal{C}_{2\ell}) = O(n^{1+1/\ell})$. A better bound when all even cycles of length at most 2ℓ are forbidden was obtained by Lam and Verstraëte [26], who showed that

$$\text{ex}(n, \mathcal{C}_{2\ell}) \leq \frac{1}{2} n^{1+1/\ell} + O(n). \quad (1)$$

It is a notoriously difficult problem to determine even the order of magnitude of $\text{ex}(n, \mathcal{C}_{2\ell})$ or $z(n, \mathcal{C}_{2\ell})$ for $\ell \notin \{2, 3, 5\}$. For $\ell \in \{2, 3, 5\}$, the existence of structures from projective geometry called *generalized polygons* show $\text{ex}(n, \mathcal{C}_{2\ell}) = \Theta(n^{1+1/\ell})$. Erdős and Simonovits [13] conjectured more generally that $\text{ex}(n, \mathcal{C}_{2\ell}) = \Theta(n^{1+1/\ell})$ for all $\ell \geq 2$, and this conjecture remains open. For large ℓ , the densest known $\mathcal{C}_{2\ell}$ -free graphs on n vertices are the Ramanujan graphs of Lubotzky, Phillips, and Sarnak [29], and the slightly denser graphs of Lazebnik, Ustimenko and Woldar [27]. The constructions have $n^{1+\theta}$ edges where $\theta \sim \frac{2}{3\ell}$ as $\ell \rightarrow \infty$. For $\ell = 4$, the current best bounds are $c_1 n^{6/5} \leq \text{ex}(n, \mathcal{C}_{2\ell}) \leq c_2 n^{5/4}$ for some constants $c_1, c_2 > 0$ – the lower bound is from the constructions of Benson [7] and Singleton [36], whereas the upper bound follows from (1) with $\ell = 4$. In this paper, we study the relationship between extremal $\mathcal{C}_{2\ell} \cup \{C_k\}$ -free graphs with k odd, and extremal $\mathcal{C}_{2\ell}$ -free bipartite graphs.

1.1. Main Results

Define the *Zarankiewicz number* $z(n, \mathcal{F})$ to be the maximum number of edges in an \mathcal{F} -free bipartite graph on n vertices. Erdős and Simonovits [13, Conjecture 3] conjectured that for any family \mathcal{F} consisting of bipartite graphs there exists an odd integer k such that $\text{ex}(n, \mathcal{F} \cup \mathcal{C}_k) \sim z(n, \mathcal{F})$. They proved this when $\mathcal{F} = \{C_4\}$ by showing that $\text{ex}(n, \{C_4, C_5\}) \sim z(n, \{C_4\})$. In this paper, we extend this by proving the following theorem which verifies the Erdős-Simonovits conjecture when $\mathcal{F} = \mathcal{C}_{2\ell}$ for $\ell \in \{2, 3, 5\}$.

Theorem 1.1. *Suppose $\ell \in \{2, 3, 5\}$ and $k > 2\ell$ is odd. Let G be a $\mathcal{C}_{2\ell} \cup \{C_k\}$ -free graph on n vertices with average degree $d = \Theta(n^{1/\ell})$. Then G has a bipartite subgraph H with at least $d^{\ell+1} - o(n^{1+1/\ell})$ edges. If also $d \geq (1 + o(1))(n/2)^{1/\ell}$ then $d \sim (n/2)^{1/\ell}$ and $e(H) \sim e(G) \sim (n/2)^{1+1/\ell}$. In particular*

$$\text{ex}(n, \mathcal{C}_{2\ell} \cup \{C_k\}) \sim z(n, \mathcal{C}_{2\ell}).$$

The last statement in the theorem was proved in the case $k=5$ and $\ell=2$ by Erdős and Simonovits [13]. The theorem for the first time determines the asymptotic behavior of $\text{ex}(n, \mathcal{C}_{2\ell} \cup \{C_k\})$ for all other odd values of $k > 2\ell$ and $\ell \in \{2, 3, 5\}$. Note that the statement in Theorem 1.1 is stronger than that of the Erdős-Simonovits conjecture in two ways. Firstly, we replace \mathcal{C}_k by C_k , i.e. we forbid a single odd cycle rather than all short odd cycles. Secondly, we obtain a *stability theorem*: any graph which is close to extremal for $\mathcal{C}_{2\ell} \cup \{C_k\}$ must be close to bipartite. This is the first ingredient in applying the *stability method*, in which one first obtains approximate structure, and then eliminates any imperfections to obtain exact structure. Using this theorem we can prove the following exact result. The reader unfamiliar with generalized $(\ell + 1)$ -gons can find a brief description in Section 3.6: these can be viewed for infinitely many prime powers q as $(q + 1)$ -regular $\mathcal{C}_{2\ell}$ -free extremal bipartite graphs with $q^\ell + q^{\ell-1} + \dots + 1$ vertices in each part.

Theorem 1.2. *Let $\ell \in \{2, 3, 5\}$, $k > 2\ell + 2$ be odd, $n > n_k$ be sufficiently large, and define $q \in \mathbb{R}^+$ by $n = 2(q^\ell + q^{\ell-1} + \dots + 1)$. Then any $\mathcal{C}_{2\ell}$ -free graph G on n vertices with at least $\frac{1}{2}(q + 1)n$ edges contains a cycle of length k , unless q is an integer and G is the bipartite incidence graph of a generalized $(\ell + 1)$ -gon of order q . Furthermore, if q is an integer, $n > n_k$ and there is a generalized $(\ell + 1)$ -gon of order q , then*

$$\text{ex}(n, \mathcal{C}_{2\ell} \cup \{C_k\}) = z(n, \mathcal{C}_{2\ell}).$$

The same statement holds when $\ell=2$ and $k=5$.

1.2. Chromatic number

A classical result of Andrásfai, Erdős and Sós [6] states that a triangle-free n -vertex graph with minimum degree more than $2n/5$ is 2-colorable. Generalizations of this theorem have been studied extensively, for example see [1,2,5,21,30] and their references. Here we address a similar question for a $\mathcal{C}_{2\ell} \cup \{C_k\}$ -free graph when $k \geq 4\ell + 1$ is odd. We use $\chi(G)$ to denote the chromatic number of G .

Theorem 1.3. *Let $\ell \geq 2$ be an integer, and let $k \geq 4\ell + 1$ be an odd integer, let c be a positive real number, and let G be a $\mathcal{C}_{2\ell} \cup \{C_k\}$ -free graph on n vertices with minimum degree at least $cn^{1/\ell}$. Then $\chi(G) < (4k)^{\ell+1}/c^\ell$.*

This theorem is effective when $k = o(n^{1/(\ell+1)})$. Forbidding short odd cycles in a graph generally has little effect on the chromatic number if the graph is too sparse. A well-known construction of Erdős in random graphs shows that there are graphs of arbitrarily large girth and chromatic number. Theorem 1.3 in contrast shows that the chromatic number becomes bounded for very dense $\mathcal{C}_{2\ell}$ -free graphs with a forbidden long odd cycle.

1.3. Short odd cycles

We start by observing that the second statement of Theorem 1.2 does not hold whenever $5 \leq k < 2\ell$ and k is odd, as in this case we have

$$\text{ex}(n, \mathcal{C}_{2\ell} \cup \{C_k\}) \geq z(n, \mathcal{C}_{2\ell}) + 1.$$

To see this, consider an extremal $\mathcal{C}_{2\ell}$ -free bipartite graph H on n vertices. Let G be obtained from H by adding an edge joining a pair of vertices $\{x, y\}$ at distance two in one part of H . We claim that G has no C_k for odd k with $5 \leq k < 2\ell$. For such a cycle would have to contain the edge xy , so we would have a path P of length $k - 1$ in H from x to y . Adding the edges xz and yz , where z is a common neighbor of x and y , we obtain a closed walk of length $k + 1$. Furthermore, this walk is not acyclic, so it must contain an even cycle of length at most 2ℓ in H , contradicting the fact that H is $\mathcal{C}_{2\ell}$ -free. It would be interesting to see if Theorem 1.2 can be extended to the last remaining case, namely $k = 2\ell + 1$.

The following proposition determines an upper bound for $z(n, \mathcal{C}_{2\ell})$. Its proof will follow easily from the counting arguments in Section 3.6.

Proposition 1.4. *Suppose $n \in \mathbb{N}$, and let $q \in \mathbb{R}^+$ be defined by $n = 2(q^\ell + q^{\ell-1} + \dots + 1)$. Then $z(n, \mathcal{C}_{2\ell}) \leq \frac{1}{2}(q + 1)n$.*

Now we can describe a much stronger discrepancy between Turán and Zarankiewicz numbers when one forbids a short odd cycle. Consider the *polarity graphs* constructed by Lazebnik, Ustimenko and Woldar [28]. These are $\mathcal{C}_{2\ell}$ -free graphs on n vertices with $(1/2+o(1))n^{1+1/\ell}$ edges, such that for $\ell=3$ they have no triangles and no even cycles of length at most six, and for $\ell=5$ they have no triangles and no cycles of length five and no even cycles of length at most ten. Thus when (ℓ, k) is $(3, 3)$, $(5, 3)$ or $(5, 5)$ we have

$$\liminf_{n \rightarrow \infty} \frac{\text{ex}(n, \mathcal{C}_{2\ell} \cup \{C_k\})}{z(n, \mathcal{C}_{2\ell})} \geq \liminf_{n \rightarrow \infty} \frac{(n^{1+1/\ell})/2}{(n/2)^{1+1/\ell}} = 2^{1/\ell},$$

so in these cases one does not even have an asymptotic result similar to Theorem 1.1. For the case $\ell=2$ and $k=3$ we do not know of such a strong discrepancy. Parsons [33] constructed $\{C_3, C_4\}$ -free graphs showing that

$$\text{ex}(n, \{C_3, C_4\}) > z(n, C_4) + \frac{7}{32}n - O(\sqrt{n})$$

when $n = \binom{q}{2}$ and $q = 1 \pmod{4}$ is prime. On the other hand, Erdős [10,11] suggested that in the case of $\{C_3, C_4\}$ -free graphs there should not be a stronger discrepancy, and conjectured that $\text{ex}(n, \{C_3, C_4\}) \sim z(n, C_4)$ – this conjecture remains open. One may also ask in general whether or not $\text{ex}(n, \mathcal{C}_{2\ell} \cup \{C_{2\ell-1}\}) \sim z(n, \mathcal{C}_{2\ell})$ for $\ell \geq 2$.

1.4. Organization

This paper is organized as follows. In the next section we illustrate our ideas by sketching the proofs of Theorems 1.1 and 1.2 in the case of quadrilateral-free graphs. Section 3 contains the essential results on counting walks and paths in graphs. These are used throughout the paper, in particular to deduce some bounds on Turán numbers and Zarankiewicz numbers for cycles in the same section. In Section 4 we show that nearly extremal graphs contain large subgraphs which are almost regular. We prove a pseudorandomness property for nearly extremal graphs in Section 5. Section 6 contains the proof of Theorem 1.1 and Section 7 the proof of Theorem 1.2. Section 8 contains the short proof of Theorem 1.3. We make some concluding remarks in the final section.

1.5. Notation

We write $e(G)$ for the number of edges in a graph G . If $S \subseteq V(G)$, then $e(S)$ denotes the number of edges of G joining vertices of S . In a graph

G , let $N_r(v)$ denote the set of vertices at distance exactly r from v , and let $d_r(v) = |N_r(v)|$. For $r = 1$, we omit the subscript r , so that $d(v)$ is the degree of v and $N(v)$ is the neighborhood of v . We write $d_B(v)$ for the number of neighbors of a vertex v in a set B . We write $G[S]$ for the subgraph of G induced by a set $S \subseteq V(G)$ and $e(S)$ for the number of edges in $G[S]$. Given two sets $S, T \subseteq V(G)$, not necessarily disjoint, we write $e(S, T)$ for the number of ordered pairs (s, t) with $s \in S, t \in T$ and $st \in E(G)$. For example $e(S, S) = 2e(S)$. In addition to the *Turán number* $\text{ex}(n, \mathcal{F})$ and *Zarankiewicz number* $z(n, \mathcal{F})$ defined above, we use $z(a, b, \mathcal{F})$ for the maximum number of edges in an \mathcal{F} -free bipartite graph that has a vertices in one part and b vertices in the other part. We let \mathbb{R}^+ denote the positive reals and \mathbb{N} the positive integers. Our asymptotic notation assumes that $n \rightarrow \infty$; we write $f(n) \sim g(n)$ if $\lim_{n \rightarrow \infty} f(n)/g(n) = 1$, and $f(n) = o(g(n))$ if $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$.

2. Sketch proof for 4-cycles

In this section we outline the proofs of Theorems 1.1 and 1.2 in the case $\ell = 2$, i.e. 4-cycles. This introduces the main ideas of our approach, without some technicalities that arise in the other cases.

2.1. Stability

We start with Theorem 1.1 for $\ell = 2$. The idea is that we can take H to be the bipartite subgraph of G , containing all the edges between $N_2(v)$ and $N_3(v)$ for a suitable vertex v . Suppose that G is a graph on n vertices with average degree d and G does not contain C_4 or C_k for some odd $k \geq 5$. The proof proceeds by the following steps.

1. Controlling the maximum degree. Lemma 4.1(i) will show that we can delete at most $2n$ edges from G to obtain a subgraph G' with maximum degree $\Delta \leq 2\sqrt{n}$. In particular, if d' is the average degree of G' then $d' \geq d - 4$.

2. Enumeration of walks and paths. The Blakley-Roy inequality (Proposition 3.1) gives a lower bound on the number of walks. It implies that we can find a vertex v that is the start of at least d'^3 walks of length 3, where $d' \geq d - 4$ is the average degree of G' . Also, the bound on the maximum degree implies that all but $O(n)$ of these walks are paths, i.e. we have $d^3 - O(n)$ paths of length 3 starting at v .

3. Finding odd cycles. We shall show in Section 3.5 that if $G[N_2(v)]$ has average degree at least $\max(6, 2k - 8)$, then G contains C_k , so we conclude

that $G[N_2(v)]$ has average degree less than $\max(6, 2k-8)$. This implies that the number of paths of length 3 which start at v and end in $N_2(v)$ is $O(n)$. Since G is C_4 -free, we also have no path of length 3 starting at v and ending in $N_1(v)$. Therefore, all but $O(n)$ of the paths found in step 2 go from v to $N_3(v)$.

4. Conclusion. Since G is C_4 -free, each edge between $N_2(v)$ and $N_3(v)$ is contained in at most one path of length 3 from v to $N_3(v)$. Thus we obtain $d^3 - O(n)$ edges between $N_2(v)$ and $N_3(v)$, and these constitute the bipartite subgraph H needed for the first statement in Theorem 1.1. For the second statement, note that Proposition 1.4 implies $z(n, C_4) \leq (1 + o(1))(n/2)^{3/2}$. Since H is a bipartite C_4 -free graph we must have $d \leq (1 + o(1))(n/2)^{1/2}$. If also $d \geq (1 + o(1))(n/2)^{1/2}$ then we have $d \sim (n/2)^{1/2}$ and $e(H) \sim e(G) \sim (n/2)^{3/2}$, which proves Theorem 1.1 for $\ell=2$.

2.2. Exact result

Now we sketch the proof of Theorem 1.2 for $\ell=2$. Suppose that n is large, G is a graph on n vertices with $e(G) \geq (q+1)n/2$, where $q \in \mathbb{R}^+$ is defined by $n = 2(q^2 + q + 1)$, and G does not contain C_4 or C_k for some odd $k \geq 7$. The proof proceeds by the following steps.

1. Pseudorandomness. Not much is known about the structure of nearly extremal graphs for bipartite Turán problems. Here we obtain a result in this direction. It says that the number of edges between any two large sets in nearly extremal graph is close to what one would expect in a random graph with the same edge density: more precisely we shall show that if G is a C_4 -free bipartite graph on n vertices with parts X and Y and average degree $d \sim (n/2)^{1/2}$, then for any $S \subseteq X$ and $T \subseteq Y$ we have $e(S, T) = \frac{2d}{n}|S||T| + o(n^{3/2})$. As pointed out by a referee, it is possible to prove this result by direct counting. However, we shall obtain this result as a special case of Theorem 5.1.

2. Controlling the minimum degree. We reduce the proof to the case when the minimum degree satisfies $\delta(G) > q/4$. This uses a vertex deletion argument that is quite standard in extremal graph theory. We consider a sequence of graphs $G = G_n, G_{n-1}, \dots, G_t$ for some $0 \leq t \leq n$, where G_{i-1} is obtained from G_i by deleting a vertex of degree at most $q/4$, while possible. If this process continues at least $n/2$ times, then we are left with a graph G_t with at most $n/2$ vertices and $e(G_t) \geq e(G) - \frac{qn}{8} \geq \frac{3}{8}qn$ edges. However, $\text{ex}(n/2, C_4) \sim \frac{1}{2}(n/2)^{3/2} < \frac{3}{8}qn$, so this is a contradiction. Now suppose that we know Theorem 1.2 holds under the additional assumption $\delta(G) > q/4$.




Figure 1. Constructing a cycle of any specified odd length $k \geq 7$

Applying this to G_t gives $e(G_t) \leq (r + 1)t/2$, where r is defined by $t = 2(r^2 + r + 1)$. Furthermore, we have $e(G) \leq e(G_t) + (n - t)q/4$, and calculations show that this is less than $(q + 1)n/2$, unless $t = n$. Since $e(G) \geq (q + 1)n/2$, we must have $t = n$, so $\delta(G) > q/4$, and we are justified in assuming this when proving Theorem 1.2.

3. Refining the approximate structure. Now we have $e(G) \geq (q + 1)n/2$ and can assume that $\delta(G) > q/4$. Let H be a bipartite subgraph of G with maximum size. Theorem 1.1 implies that $e(H) \sim e(G) \sim (n/2)^{3/2}$. Furthermore, maximality of H implies that $\delta(H) > q/8$. Actually, we only need the fact that $\delta(H) > cn^{1/2}$, where $c > 0$ is independent of n . Now we will show that $G = H$. Label the parts of H as A and B . Suppose for a contradiction that $G[A]$ contains an edge ac . Let b be a neighbor of c in B . Now we ‘explore’ the graph until we reach two sets of linear size where we can apply pseudorandomness (see Figure 1). By the minimum degree assumption, we can take $|A_b|, |B_a| \sim cn^{1/2}$ such that all vertices in A_b are neighbors of b and all vertices in B_a are neighbors of a . Next, taking all the neighbors of vertices in A_b we get a set B_b with $|B_b| = \Theta(n)$ since G is C_4 -free. By taking neighbors of B_a we similarly get a set A_a with $|A_a| = \Theta(n)$. Then Step 1 (Pseudorandomness) gives

$$e(A_a, B_b) = \frac{2d}{n}|A_a||B_b| + o(n^{3/2}) = \Theta(n^{3/2}).$$

In particular, we can find a path of length $k - 6$ using only edges between A_a and B_b . By construction this can be completed to a cycle of length k in G , which is a contradiction. We deduce that A is an independent set in G . Similarly B is independent, so $G = H$ is bipartite. The characterization of

equality now follows from a result on Zarankiewicz numbers for even cycles – see Proposition 3.8 – so this proves Theorem 1.2 for quadrilaterals.

3. Counting walks

The basis of estimates for both Turán numbers and Zarankiewicz numbers for even cycles is counting various types of walks in graphs. These counts are also used in Step 2 of the stability result (Enumeration of walks and paths) and Step 1 of the exact result (Pseudorandomness). A *walk of length t* in a graph G is a sequence $v_0e_0v_1e_1\dots v_{t-1}e_{t-1}v_t$ such that $v_i \in V(G)$, $e_i \in E(G)$ and $e_i = v_iv_{i+1}$ for $0 \leq i < t$. Note that edges may be repeated in a walk, even on consecutive steps.

3.1. The Blakley-Roy Inequality

Let $w_k(G)$ denote the number of walks of length k in a graph G divided by the number of vertices in G – this is the average number of walks of length k starting at a vertex. If G is an d -regular graph on n vertices, then clearly $w_k(G) = d^k$. Blakley and Roy [8] proved a matrix version of Hölder's Inequality, which shows (as a special case) that any graph of average degree d has at least as many walks of a given length as an d -regular graph on the same number of vertices:

Proposition 3.1. *Suppose G is a graph of average degree d . Then $w_k(G) \geq d^k$.*

There are many proofs of this inequality in the literature; the original proof of Blakley and Roy uses eigenvalues. We now briefly discuss the tight connection between walks and eigenvalues.

3.2. Walks and eigenvalues

Let G be a graph on n vertices with adjacency matrix A . Then we may write

$$w_k(G) = e^\top A^k e \quad \text{where} \quad e = n^{-1/2}(1, 1, \dots, 1)^\top.$$

In this notation, the Blakley-Roy inequality (Proposition 3.1) mentioned earlier can be stated as $w_k \geq w_1^k$.

We can also use spectral theory to count closed walks. Let w_k° be the number of closed walks of length k in a graph G divided by the number

of vertices in G – this is the average number of closed walks of length k starting at a vertex. Suppose A has eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. We need the following well-known formula.

Proposition 3.2. $w_k^\circ := \frac{1}{n} \operatorname{Tr}(A^k) = \frac{1}{n} \sum_{i=1}^n \lambda_i^k$.

This gives rise to a standard method for establishing a spectral gap in a d -regular graph: if w_k° is close to $\lambda_1^k = d^k$, then all $|\lambda_i|$ for $i > 1$ must be small. This idea will be extended to proving bounds on eigenvalues of the adjacency matrix of regular and almost regular bipartite graphs. We have the following bound on $w_{2\ell+2}^\circ$ in dense $\mathcal{C}_{2\ell}$ -free bipartite graphs with roughly equal part sizes.

Lemma 3.3. *Suppose G is a bipartite graph on n vertices with part sizes $n/2 + o(n)$, maximum degree Δ and girth at least $2\ell + 2$. Then*

$$w_{2\ell+2}^\circ(G) < (1/2 + o(1))n\Delta^2 + (4\Delta)^{\ell+1}.$$

Proof. Consider any vertex v and a closed walk W of length $2\ell+2$ from v . If W is not a cycle then the girth assumption implies that its underlying graph is acyclic. The number of such walks W is therefore at most the number of closed walks of length $2\ell+2$ from the root in the complete Δ -ary tree. A crude upper bound on this number is $\binom{2\ell+2}{\ell+1} \Delta^{\ell+1} < (4\Delta)^{\ell+1}$, as may be seen by choosing the $\ell+1$ times when the walk moves towards the root and multiplying Δ choices for the $\ell+1$ times when the walk moves away from the root. (The exact formula is $\frac{1}{\ell+2} \binom{2\ell+2}{\ell+1} \Delta^{\ell+1}$ but we do not need this.) In the case when W is a cycle, we estimate the possibilities by considering the neighbors a and b of v on W , and the opposite vertex u at distance $\ell+1$ from v on W . Note that W is uniquely determined by a , b and u , as the girth assumption implies that there is at most one path of length ℓ between any specified pair of vertices. We can choose each of a and b in at most Δ ways, and u in at most $n/2 + o(n)$ ways (using the assumption on the part sizes). Thus the number of possibilities is at most $(1/2 + o(1))n\Delta^2$, which also takes into account the orientation of the cycle. Combining the two cases, the number of closed walks of length $2\ell+2$ from v is at most $(1/2 + o(1))n\Delta^2 + (4\Delta)^{\ell+1}$. Since v was arbitrary, this bound also holds for the average $w_{2\ell+2}^\circ(G)$. ■

3.3. Non-returning walks

A *non-returning walk* of length k in a graph G is a walk $v_0 e_0 v_1 e_1 \dots v_{k-1} e_{k-1} v_k$ of length k such that $e_i \neq e_{i+1}$ for $0 \leq i < k$. Let $\nu_k(G)$ denote the number of non-returning walks of length k in a graph G divided by

the number of vertices in G . If G is an r -regular graph on n vertices then clearly $\nu_k(G) = r(r-1)^{k-1}$. Alon, Hoory and Linial [3] gave an analogue of Proposition 3.1 for non-returning walks.

Proposition 3.4. *Suppose G is a graph of average degree $r \geq 2$. Then $\nu_k(G) \geq r(r-1)^{k-1}$. If $r \in \mathbb{N}$ then equality holds if and only if G is r -regular.*

In fact, they proved a slightly stronger form that the result given, which is sensitive to variations in the degrees of the vertices in the graph. Sidorenko [35] gave a bipartite analogue of Proposition 3.1. Here we need a bipartite analogue of Proposition 3.4, which was proved by Hoory [23]. We will only need the result for walks of odd length.

Proposition 3.5. *Suppose G is a bipartite graph with parts A and B and average degree d . Let α be the average degree of vertices in A and β the average degree of vertices in B . Then for any $t \in \mathbb{N}$ we have*

$$\nu_{2t+1} \geq d \prod_{v \in A \cup B} (d(v) - 1)^{td(v)/e(G)} \geq d(\alpha - 1)^t(\beta - 1)^t.$$

If $\alpha, \beta \in \mathbb{N}$ then equality holds if and only if every vertex of A has degree α and every vertex of B has degree β .

3.4. Paths

A *path* is a walk which has no repeated vertices or edges. Let $p_\ell(G)$ denote the number of paths of length ℓ in a graph G divided by the number of vertices in G .

Lemma 3.6. *Suppose G is a graph with maximum degree Δ and average degree d , and $\ell \in \mathbb{N}$. Then*

$$p_\ell(G) \geq d^\ell - \ell^2 \Delta^{\ell-1}.$$

Proof. By the Blakley-Roy inequality there are at least nd^ℓ walks of length ℓ in G if G has n vertices. Let $(v_1, v_2, \dots, v_\ell)$ be a walk of length ℓ which is not a path. Then $v_i = v_j$ for some distinct $i, j \in \{1, 2, \dots, \ell\}$. Fixing $i < j$, there are at most $n\Delta^{j-1}$ choices for the part of the walk up to v_{j-1} , then the next step is determined since $v_j = v_i$, and then there are at most $\Delta^{\ell-j}$ choices for the remaining steps. There are at most ℓ^2 choices for i and j , and therefore the number of walks of length ℓ which are not paths is at most $n\ell^2 \Delta^{\ell-1}$. Dividing by n , we obtain the lemma. ▀

3.5. Finding odd cycles

We use the following lemma which is implicit in [40].

Lemma 3.7. *Suppose G is a graph, v is a vertex of G and $G[N_r(v)]$ has average degree at least $2s-4$, for some $r \geq 1$ and odd $s \geq 5$. Then G contains cycles $C_{2m+1}, C_{2m+3}, \dots, C_{2m+s}$ for some $1 \leq m \leq r$. In particular, if G is a graph not containing a cycle C_k , for some fixed odd $k > 2r$, then $G[N_r(v)]$ has average degree less than $\max(6, 2k-8)$.*

Proof. To give a self-contained proof of this lemma would require the duplication of large parts of [40], so instead we just sketch the argument, and refer the reader to [40] for the omitted details. We use the fact that any graph with average degree at least $2d$ has a subgraph of minimum degree at least $d+1$. Following the proof of [40, Lemma 3], since $G[N_r(v)]$ has average degree at least $2d=2(s-2)$, there is a subgraph of $G[N_r(v)]$ with minimum degree at least $d=s-1$. It is well-known (and easy to see by considering a longest path) that a graph of minimum degree at least $d \geq 3$ has a cycle of length at least $d+1$ with at least one chord. Therefore $G[N_r(v)]$ contains a subgraph H which is a cycle of length at least s with at least one chord. Next, as in the proof of [40, Theorem 1], we consider a minimal subtree T of the breadth first search tree rooted at v that contains $V(H)$. By minimality T branches at its root (which may not be v). We let A be the set of vertices of H belonging to one branch of T and let $B=V(H) \setminus A$. Next we apply [40, Lemma 2], which tells us that H contains paths from A to B of every length $\ell \leq |V(H)|$, unless H is bipartite with bipartition (A, B) . Even if H is bipartite with bipartition (A, B) we still have paths from A to B of every odd length $\ell \leq |V(H)|$. These may be completed to cycles via paths in T , which all have the same length $2m$, where m is the distance from the root of T to $N_r(v)$. The second statement of the lemma is proved by taking $s = \max(5, k-2)$. ■

3.6. Zarankiewicz numbers of even cycles

In this section, we describe extremal bipartite $\mathcal{C}_{2\ell}$ -free graphs and generalized polygons, and give a proof of extremality by using the walk counting arguments covered in the previous sections. In the cases $\ell \in \{2, 3, 5\}$, the bound (1) for $\text{ex}(n, \mathcal{C}_{2\ell})$ is asymptotically tight, i.e. $\text{ex}(n, \mathcal{C}_{2\ell}) \sim \frac{1}{2}n^{1+1/\ell}$. This is shown by constructions from projective geometry, due to Erdős and Rényi [12] for $\ell=2$, and to Benson [7] and Singleton [36] for $\ell=3$ and $\ell=5$, based on generalized polygons. We briefly describe these here, referring the

reader to [28] for more details. Suppose P is a set of points, L a set of lines, and I an incidence relation between P and L . The *bipartite incidence graph* is the bipartite graph with parts P and L such that $p \in P$ is adjacent to $\ell \in L$ if and only if $(p, \ell) \in I$. Let $q \geq 2$ and $\ell \geq 2$ be integers. A *generalized $(\ell + 1)$ -gon of order q* consists of a set of $q^\ell + q^{\ell-1} + \dots + 1$ points P and a set of $q^\ell + q^{\ell-1} + \dots + 1$ lines L with an incidence relation I such that the bipartite incidence graph of (P, L, I) is $(q + 1)$ -regular and has girth $2\ell + 2$ and diameter $\ell + 1$. Feit and Higman [16] showed that generalized $(\ell + 1)$ -gons of order q only exist for $\ell = 2, 3, 5$. These are known as generalized triangles, generalized quadrangles and generalized hexagons respectively. Generalized triangles are precisely projective planes, whereas the generalized quadrangles and generalized hexagons were first constructed by Tits in his seminal paper [39].

We start with the proof of an upper bound for $z(n, C_{2\ell})$, and the characterization of equality by generalized polygons described above. For $\ell = 2$ this result is well-known (see [4], pp. 273–274 for a nice exposition). Here it is convenient to just give the proof for odd $\ell > 2$, as we do not need the other cases.

Proposition 3.8. *Let $\ell = 2$ or $\ell > 2$ be odd. Suppose $n \in \mathbb{N}$, and let $q \in \mathbb{R}^+$ be defined by $n = 2(q^\ell + q^{\ell-1} + \dots + 1)$. Then $z(n, C_{2\ell}) \leq \frac{1}{2}(q + 1)n$, with equality if and only if q is a positive integer and there exists a generalized $(\ell + 1)$ -gon on n points.*

Proof. Let $\ell > 2$ be odd and G be a bipartite graph on n vertices with $e(G) \geq (q + 1)n/2$ containing no even cycle of length at most 2ℓ . Then G has average degree at least $q + 1$. Also, G has diameter at least $\ell + 1$ since G has no cycles of length less than $2\ell + 2$. Let A and B be the parts of G . Then $|A| + |B| = n$. By the girth assumption, for any ordered pair of vertices x, y there is at most one non-returning walk of length at most ℓ from x to y . Since ℓ is odd, we need only consider pairs with x and y in different parts, of which there are $2|A||B| \leq n^2/2$. By Proposition 3.4 (using that $(\alpha - 1)(\beta - 1) \leq (\frac{\alpha + \beta}{2} - 1)^2$ together with $\frac{\alpha + \beta}{2} = q + 1$), the number of non-returning walks of odd length at most ℓ is at least

$$n(q + 1) \sum_{i=0}^{(\ell-1)/2} q^{2i} = n \sum_{j=0}^{\ell} q^j = n^2/2.$$

We conclude that the number of non-returning walks of length at most ℓ is exactly $2|A||B| = n^2/2$, so $|A| = |B| = n/2$. Since equality only holds in Proposition 3.4 for regular graphs, every vertex of G has degree $q + 1$. It

also follows that every pair of vertices from different parts is connected by a path of length at most ℓ . Thus every pair of vertices is connected by a path of length at most $\ell + 1$. Since G has diameter at least $\ell + 1$, we conclude G has diameter exactly $\ell + 1$. Therefore G is the bipartite incidence graph of a generalized $(\ell + 1)$ -gon. ■

We also require the following bound on unbalanced Zarankiewicz numbers $z(a, b, \mathcal{C}_{2\ell})$.

Proposition 3.9. *Suppose $\ell = 2$ or $\ell \geq 3$ is odd, and $a, b \geq 1$. Then*

$$z(a, b, \mathcal{C}_{2\ell}) \leq (ab)^{\frac{1}{2} + \frac{1}{2\ell}} + \max\{a, b\}.$$

Proof. The case $\ell = 2$ follows from a slightly stronger result of Reiman [34]. Now suppose $\ell \geq 3$ is odd. Let G be a $\mathcal{C}_{2\ell}$ -free bipartite graph with parts A and B of sizes a and b , and let α and β denote the average degrees of vertices in A and B . As in the proof of Proposition 3.8, there are at most $2ab$ non-returning walks of odd length at most ℓ . Using the lower bound from Proposition 3.5, we deduce that

$$2e(G) \sum_{i=0}^{(\ell-1)/2} (\alpha - 1)^i (\beta - 1)^i \leq 2ab. \tag{2}$$

Suppose for a contradiction that $e(G) > (ab)^c + \max\{a, b\}$, where $c = \frac{1}{2} + \frac{1}{2\ell}$. Then $\alpha - 1 = (e(G) - a)/a > a^{c-1}b^c$ and $\beta - 1 = (e(G) - b)/b > a^c b^{c-1}$, so $(\alpha - 1)(\beta - 1) > (ab)^{2c-1} = (ab)^{1/\ell}$. However, this gives

$$e(G)(\alpha - 1)^{(\ell-1)/2}(\beta - 1)^{(\ell-1)/2} > (ab)^c(ab)^{(\ell-1)/2\ell} = ab,$$

which contradicts (2). This gives the required bound on $e(G)$. ■

4. Degrees in nearly extremal graphs

In this section, we show that in nearly extremal $\mathcal{C}_{2\ell}$ -free bipartite graphs for $\ell \in \{2, 3, 5\}$, the number of edges containing a vertex of degree substantially more than the average degree is small. This is used in Step 1 of the stability result (Controlling the maximum degree) and Step 1 of the exact result (Pseudorandomness). We also show how to classify extremal graphs once extremal graphs of large minimum degree are classified. This is used in Step 2 of the exact result (Controlling the minimum degree).

4.1. Bounding the maximum degree of C_4 -free graphs

First we show that any C_4 -free bipartite graph on n vertices does not contain many edges on vertices of degree much more than roughly $(n/2)^{1/2}$, which is the average degree in extremal C_4 -free bipartite graphs by Proposition 3.8. Recall that we write $d_B(v)$ for the number of neighbors of a vertex v in a set B and $e(S, B)$ for the number of ordered pairs (s, b) with $s \in S$, $b \in B$ and $sb \in E(G)$.

Lemma 4.1. *Suppose G is a C_4 -free graph on n vertices and $A, B \subset V(G)$. Suppose $0 < \varepsilon < \sqrt{3}$ and let*

$$S = \{v \in A : d_B(v) \geq (1 + \varepsilon)|B|^{1/2}\}.$$

Then $e(S, B) \leq 2|B|/\varepsilon$. In particular, we have the following:

- (i) *If G is a C_4 -free graph then at most $2n/\varepsilon$ edges contain vertices of degree at least $(1 + \varepsilon)\sqrt{n}$.*
- (ii) *If G is a C_4 -free bipartite graph with parts X and Y , then $e(T, X \cup Y) \leq 2n/\varepsilon$, where*

$$T = \{v \in X : d_Y(v) \geq (1 + \varepsilon)|Y|^{1/2}\} \cup \{v \in Y : d_X(v) \geq (1 + \varepsilon)|X|^{1/2}\}.$$

Proof. Since G is C_4 -free, every choice of a vertex v in B and two neighbors s, s' of v in S gives rise to a different pair $\{s, s'\}$ in S , so

$$\sum_{v \in B} \binom{d_S(v)}{2} \leq \binom{|S|}{2}.$$

Since the function $f(x) = x(x - 1)$ is convex, Jensen's inequality gives

$$|S|(|S| - 1) \geq e(S, B)(e(S, B)/|B| - 1)$$

which implies

$$|B||S|^2 \geq e(S, B)(e(S, B) - |B|).$$

Suppose for a contradiction that $e(S, B) > 2|B|/\varepsilon$. Then

$$e(S, B) - |B| > (1 - \varepsilon/2)e(S, B).$$

Since $e(S, B) \geq |S| \cdot (1 + \varepsilon)|B|^{1/2}$ by definition of S , we have

$$\begin{aligned} |B||S|^2 &\geq e(S, B)(e(S, B) - |B|) \\ &\geq (1 - \varepsilon/2)e(S, B)^2 \geq (1 - \varepsilon/2)(1 + \varepsilon)^2|B||S|^2. \end{aligned}$$

However, $(1 - \varepsilon/2)(1 + \varepsilon)^2 = 1 + (3 - \varepsilon^2)\varepsilon/2 > 1$, contradiction. So we must instead have $e(S, B) \leq 2|B|/\varepsilon$. Now statement (i) follows by taking $A = B = V(G)$, and statement (ii) follows by first taking $A = X$ and $B = Y$ and then taking $A = Y$ and $B = X$. ■

4.2. Bounding the maximum degree in $\mathcal{C}_{2\ell}$ -free graphs

Next we prove an analogue of Lemma 4.1 for $\mathcal{C}_{2\ell}$ -free bipartite graphs. The accurate estimate given here is only needed for the pseudorandomness argument; a cruder ad hoc method will suffice for bounding the maximum degree in Theorem 1.1. In the following lemma we could give a quantitative description of how the $o(\cdot)$ estimate depends on d ; however the calculations are somewhat heavy, so we rather retain only the asymptotic statement as $n \rightarrow \infty$.

Lemma 4.2. *Let $\varepsilon > 0$ and suppose G' is a $\mathcal{C}_{2\ell}$ -free bipartite graph on n vertices with average degree $d \sim (n/2)^{1/\ell}$, where ℓ is odd. Then at most $o(e(G'))$ edges contain a vertex of degree at least $(1 + \varepsilon)d$.*

Proof. We start by repeatedly removing vertices of degree 0 or 1 to leave a graph G with minimum degree at least 2. This process removes at most $n = o(e(G'))$ edges, so Proposition 3.8 implies that we remove $o(n)$ vertices. Let A and B be the parts of G . As in the proof of Propositions 3.8 and 3.9, the girth assumption implies that the number of non-returning walks of odd length at most ℓ is at most $2|A||B|$. Here we will just use the estimate

$$\nu_\ell(G) \leq 2|A||B|/n \leq n/2. \tag{3}$$

In the lower bound for $\nu_\ell(G)$ we will use the full strength of Proposition 3.5 to get an improvement if there are many edges incident to vertices of large degree. First we show that the bipartition is roughly balanced. Proposition 3.9 gives $e(G) \leq (|A||B|)^{\frac{1}{2} + \frac{1}{2\ell}} + n$. Since $e(G) \sim e(G') = nd/2 \sim (n/2)^{1+1/\ell}$ we deduce that $|A| \sim |B| \sim n/2$.

Now let S be the set of vertices in A of degree at least $(1 + \varepsilon)d$. We will show that $e(S, B) = o(e(G))$. Proposition 3.5 gives

$$\nu_\ell(G) \geq d\pi(A)^{(\ell-1)/2}\pi(B)^{(\ell-1)/2}, \tag{4}$$

where for any $C \subseteq V(G)$ we write

$$\pi(C) = \prod_{v \in C} (d(v) - 1)^{d(v)/e(G)}.$$

For $\pi(B)$ we just use the simple bound

$$\pi(B) \geq e(G)/|B| - 1 \sim d.$$

To see this we apply Jensen’s inequality with the convex function $f(x) = x \log(x - 1)$ for $x \geq 2$, using the fact that G has minimum degree at least 2.

This shows that $\pi(B)$ is minimized when $d(v) = e(G)/|B|$ for all $v \in B$. Since $e(G) \sim (n/2)^{1+1/\ell}$ and $|B| \sim n/2$ we get the stated bound on $\pi(B)$. For $\pi(A)$ we estimate $\pi(S)$ and $\pi(A \setminus S)$ separately. We write

$$|S| = \sigma n/2 \quad \text{and} \quad e(S, B) = \rho e(G).$$

From Proposition 3.9, we have

$$e(S, B) \leq (|S||B|)^{\frac{1}{2} + \frac{1}{2\ell}} + n.$$

Also $e(S, B) \geq |S|(1 + \varepsilon)d$, so we conclude ρ and σ satisfy

$$(1 + \varepsilon)\sigma \leq \rho \leq \sigma^{\frac{1}{2} + \frac{1}{2\ell}} + o(1). \tag{5}$$

By Jensen’s inequality, $\pi(S)$ is minimized when $d(v) = \rho e(G)/|S|$ for all $v \in S$, and $\pi(A \setminus S)$ is minimized when $d(v) = (1 - \rho)e(G)/|A \setminus S|$ for all $v \in A \setminus S$. Therefore

$$\begin{aligned} \pi(A) &= \pi(S)\pi(A \setminus S) \\ &\geq \left(\frac{\rho e(G)}{|S|} - 1\right)^\rho \left(\frac{(1 - \rho)e(G)}{|A \setminus S|} - 1\right)^{1-\rho} \sim d \left(\frac{\rho}{\sigma}\right)^\rho \left(\frac{1 - \rho}{1 - \sigma}\right)^{1-\rho}. \end{aligned}$$

Applying (3) and (4) we deduce that

$$n/2 \geq \nu_\ell(G) \geq (1 + o(1))d \left(d^2 \left(\frac{\rho}{\sigma}\right)^\rho \left(\frac{1 - \rho}{1 - \sigma}\right)^{1-\rho} \right)^{(\ell-1)/2}.$$

Since $d \sim (n/2)^{1/\ell}$ we obtain

$$\left(\frac{\rho}{\sigma}\right)^\rho \left(\frac{1 - \rho}{1 - \sigma}\right)^{1-\rho} \leq 1 + o(1).$$

Now recall that our goal is to show that $e(S, B) = o(e(G))$, i.e. $\rho = o(1)$. Suppose for a contradiction that we can choose an infinite sequence of graphs G_n as above with analogous parameters ρ_n and σ_n such that $\rho_n \geq c$ for some constant $c > 0$. Then we can pass to a subsequence such that $\sigma_n \rightarrow s$ and $\rho_n \rightarrow r$, for some $r, s \in [0, 1]$ with $r \geq c$. Then r and s satisfy

$$f(r, s) := \left(\frac{r}{s}\right)^r \left(\frac{1 - r}{1 - s}\right)^{1-r} \leq 1. \tag{6}$$

We also have $(1+\varepsilon)s \leq r \leq s^{\frac{1}{2} + \frac{1}{2\ell}}$ by (5). Note that this implies $r < 1$. Consider the function $g(t) = f(r, t)$. Computation of derivatives gives

$$g'(t) = \frac{t - r}{t(1 - t)}g(t) \quad \text{and} \quad g''(t) = \frac{2(t - r)^2 + r(1 - r)}{t^2(1 - t)^2}g(t).$$

Thus $g'(r) = 0$ and $g''(t) > 0$ for all t , so $g(t)$ is convex, is decreasing for $t \leq r$ and is minimized at $t = r$. Since $s \leq r/(1 + \varepsilon)$ the minimum value possible for $f(r, s)$ is at $s = r/(1 + \varepsilon)$. Substituting in (6) and simplifying gives

$$1 + \varepsilon \leq \left(1 + \frac{\varepsilon}{1 - r}\right)^{1 - r}.$$

However this is a contradiction, by the standard inequality $(1 + y/x)^x < 1 + y$ for $x, y \in (0, 1)$. We deduce that $e(S, B) = o(e(G))$. The same argument shows that $o(e(G))$ edges contain a vertex in B of degree at least $(1 + \varepsilon)d$, so the proof is complete. ■

4.3. Bounding the minimum degree

Here we implement Step 2 in the proof of the exact result, by reducing the proof of Theorem 1.2 to the case when the minimum degree satisfies $\delta(G) > q/4$. Recall that G is a $\mathcal{C}_{2\ell}$ -free graph on n vertices with at least $\frac{1}{2}(q + 1)n$ edges, where n is large, $\ell \in \{2, 3, 5\}$ and $q \in \mathbb{R}^+$ is defined by $n = 2(q^\ell + q^{\ell-1} + \dots + 1)$. We use the vertex deletion argument as in Section 2. Consider a sequence of graphs $G = G_n, G_{n-1}, \dots, G_t$ for some $0 \leq t \leq n$, where G_{i-1} is obtained from G_i by deleting a vertex of degree at most $q/4$, while possible. Note that, by definition, the minimum degree of G_t is at least $q/4$. We claim that $t > n/2$. For suppose otherwise and consider $G_{n/2}$. We have $e(G_{n/2}) \leq \frac{1}{2}(n/2)^{1+1/\ell} + O(n)$ by (1). Also, the number of edges deleted is at most $(n/2)(q/4)$. Since we assume that $e(G) \geq (q + 1)n/2$ we get $e(G_{n/2}) > \frac{3}{4}qn/2$. But $qn/2 \sim (n/2)^{1+1/\ell}$, so this contradicts the upper bound. Thus we do have $t > n/2$. Next we claim that $e(G_t) \geq (r + 1)t/2$, where r is defined by $t = 2 \sum_{i=0}^{\ell} r^i$. To see this note that $e(G_t) \geq e(G) - (n - t)q/4 \geq (q + 1)n/2 - (n - t)q/4$. Then, using $q \geq r$, we calculate

$$\begin{aligned} & (q + 1)n/2 - (r + 1)t/2 - (n - t)q/4 \\ &= (q + 1) \sum_{i=0}^{\ell} q^i - (r + 1) \sum_{i=0}^{\ell} r^i - (q/2) \left(\sum_{i=0}^{\ell} q^i - \sum_{i=0}^{\ell} r^i \right) \end{aligned}$$

$$\begin{aligned}
 &= (q/2 + 1) \sum_{i=0}^{\ell} q^i - (r + 1 - q/2) \sum_{i=0}^{\ell} r^i \\
 &\geq (q/2 + 1) \left(\sum_{i=0}^{\ell} q^i - \sum_{i=0}^{\ell} r^i \right) \geq 0.
 \end{aligned}$$

We deduce that $e(G_t) \geq (r + 1)t/2$, with strict inequality unless $t = n$. Now suppose that we know Theorem 1.2 holds under the additional assumption $\delta(G) > q/4$. Applying this to G_t gives $e(G_t) \leq (r + 1)t/2$. Thus we must have $t = n$, so $\delta(G) > q/4$, and we are justified in assuming this when proving Theorem 1.2.

5. Pseudorandomness

A key ingredient in the proofs of Theorem 1.1 and 1.2 is the notion of *pseudorandomness*. There are many equivalent notions of pseudorandomness in graphs: we refer the reader to [25] for a survey. In this section we will present spectral properties of graphs that imply pseudorandomness. We will prove the following result, which expresses the pseudorandomness property of a $C_{2\ell}$ -free bipartite graph of close to maximum size: for any two large sets the number of edges between them is roughly the same as in a random bipartite graph of the same density.

Theorem 5.1. *For every $\delta > 0$, there exists $\varepsilon > 0$ such that if G is a $C_{2\ell}$ -free bipartite graph on n vertices with parts X and Y and average degree $d \geq (1 - \varepsilon)(n/2)^{1/\ell}$, then for any $S \subseteq X$ and $T \subseteq Y$ we have*

$$\left| e(S, T) - \frac{2d}{n} |S||T| \right| \leq \delta n^{1+1/\ell}.$$

This theorem for $\ell = 2$ is the pseudorandomness part of the sketch proof given in Section 2.2 and a purely combinatorial proof of this can be given, as pointed out by a referee. The proof of Theorem 5.1 is linear-algebraic, however, as pointed out by a referee a combinatorial proof is available when $\ell = 2$. In the first subsection we briefly discuss the fact, first proved by N. Alon, that a regular graph with a large spectral gap is pseudorandom (this is sometimes known as the ‘expander mixing lemma’). We also prove a bipartite version of this result which seems not have been explicitly presented in the previous literature. In the second subsection we adapt this result to the case of nearly regular bipartite graphs. Then we prove Theorem 5.1 in the final subsection.

5.1. Pseudorandomness of regular graphs

Suppose G is a graph on n vertices and let $A = (a_{uv})_{u,v \in V(G)}$ be its adjacency matrix, i.e. a_{uv} is 1 if uv is an edge or 0 otherwise. (We fix some ordering v_1, v_2, \dots, v_n of the vertices and identify v_i with i .) Then A is a real symmetric matrix, so has an orthonormal basis x_1, x_2, \dots, x_n of eigenvectors with real eigenvalues, which we order so that $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. Note that $\sum_{i=1}^n \lambda_i = \text{Tr}(A) = 0$, so $\lambda_1 \geq 0$ and $\lambda_n \leq 0$. The Perron-Frobenius theorem implies that $|\lambda_i| \leq \lambda_1$ for all i and all entries of x_1 are non-negative. If G is d -regular then we have $x_1 = e := n^{-1/2}(1, 1, \dots, 1)$ and $\lambda_1 = d$. In this case it is easy to verify that $|\lambda_i| \leq d$ for all i , since if $x_{i,j}$ has the largest absolute value among the coordinates of x_i then

$$|\lambda_i x_{i,j}| = |(Ax_i)_j| = \left| \sum_{k \in N(j)} x_{i,k} \right| \leq d|x_{i,j}|.$$

Write

$$\lambda = \max_{i \neq 1} |\lambda_i|.$$

We have the following pseudorandomness property for regular graphs (see, e.g., [4,25]):

Lemma 5.2. *Suppose G is a d -regular graph on n vertices. Then for any $S, T \subseteq V(G)$ we have*

$$\left| e(S, T) - \frac{d}{n} |S||T| \right| \leq \lambda \sqrt{|S||T|}.$$

This lemma can be adapted to the bipartite setting as follows. Let G be a bipartite graph on n vertices with parts X and Y . We choose the ordering v_1, v_2, \dots, v_n of the vertices so that X precedes Y . When we consider vectors of length n this gives us a natural partition of its co-ordinates into two blocks corresponding to X and Y . Then the adjacency matrix A has block structure $\begin{pmatrix} 0 & M \\ M^\top & 0 \end{pmatrix}$ where M is the bipartite incidence matrix of G , i.e. M has rows indexed by X , columns indexed by Y , and entries m_{xy} equal to 1 if xy is an edge, otherwise 0.

In this subsection we just consider the case when G is d -regular, which implies that $|X| = |Y| = n/2$. Then we have $\lambda_1 = d$, with eigenvector $e = n^{-1/2}(1, \dots, 1)^\top$ as before, and $\lambda_n = -d$, with eigenvector $\bar{e} = n^{-1/2}(1, \dots, 1, -1, \dots, -1)^\top$ having $n^{-1/2}$ in the coordinates corresponding to elements of X , and $-n^{-1/2}$ in the coordinates corresponding to elements

of Y . In the bipartite setting we re-define λ (henceforth we will not use the previous definition) by

$$\lambda = \max_{i \neq 1, n} |\lambda_i|.$$

We have the following pseudorandomness property for regular bipartite graphs.

Lemma 5.3. *Suppose G is a d -regular bipartite graph on n vertices with parts X and Y . Then for any $S \subseteq X$ and $T \subseteq Y$ we have*

$$\left| e(S, T) - \frac{2d}{n} |S||T| \right| \leq \lambda \frac{|S| + |T|}{2}.$$

Proof. Let $\chi = \chi_{S \cup T}$ denote the characteristic vector of $S \cup T$. Then $\chi^\top A \chi = \chi_S^\top M \chi_T + \chi_T^\top M^\top \chi_S = 2e(S, T)$. Writing $\chi = \sum_{i=1}^n a_i x_i$ in the eigenvector basis we obtain $2e(S, T) = \sum_{i=1}^n \lambda_i a_i^2$. Since $\sum_{i=1}^n a_i^2 = \chi \cdot \chi = |S| + |T|$ we can estimate $\left| \sum_{i \neq 1, n} \lambda_i a_i^2 \right| \leq |\lambda|(|S| + |T|)$. We have

$$\begin{aligned} \lambda_1 &= d, & a_1 &= e \cdot \chi = n^{-1/2}(|S| + |T|), \\ \lambda_n &= -d & \text{and} & \quad a_n = \bar{e} \cdot \chi = n^{-1/2}(|S| - |T|), \end{aligned}$$

so $\lambda_1 a_1^2 + \lambda_n a_n^2 = \frac{4d}{n} |S||T|$. This gives the stated estimate for $e(S, T)$. ■

We remark that an alternative derivation of these estimates may be obtained from the singular value decomposition of the bipartite incidence matrix.

5.2. Nearly regular bipartite graphs

We want to show pseudorandomness for $\mathcal{C}_{2\ell} \cup \{C_k\}$ -free graphs which are nearly extremal. Such graphs are not necessarily regular – for instance they may contain isolated vertices – so to treat them we will prove an analogue of Lemma 5.3 for nearly regular bipartite graphs. The quantity we use to measure irregularity of a graph is its *variance*. If G is a graph of average degree d with n vertices, then the variance of G is defined by

$$\text{Var}(G) := \frac{1}{n} \sum_v (d(v) - d)^2 = \frac{1}{n} \sum_v d(v)^2 - d^2.$$

As before we set $\lambda = \lambda(G) = \max_{i \neq 1, n} |\lambda_i|$ in the bipartite setting. We have the following pseudorandomness property for nearly regular bipartite graphs (for a similar statement for non-bipartite graphs see [25]).

Lemma 5.4. *Let $\beta, \gamma \in (0, 1)$ and $\alpha = 4\beta^{1/2}\gamma^{-1} < 1/4$. Suppose $G = G(X, Y)$ is a bipartite graph on n vertices with average degree d and*

- (i) $\lambda(G) < (1 - \gamma)d$,
- (ii) $\text{Var}(G) < \beta d^2$.

Then for any $S \subseteq X$ and $T \subseteq Y$ we have

$$\left| e(S, T) - \frac{2d}{n}|S||T| \right| \leq (4\alpha d + \lambda/2)n.$$

Let $e = n^{-1/2}(1, \dots, 1)^\top$, and let $\bar{e} := n^{-1/2}(1, \dots, 1, -1, \dots, -1)^\top$, where co-ordinates labelled by the elements of X are $n^{-1/2}$, and co-ordinates labelled by the elements of Y are $-n^{-1/2}$. The following estimates will be used in the proof and later in the paper.

$$\lambda_1 = \max\{x^\top Ax : \|x\| = 1\} \geq e^\top Ae = n^{-1} \sum_v d(v) = d, \quad \text{and} \quad (7)$$

$$\lambda_n = \min\{x^\top Ax : \|x\| = 1\} \leq \bar{e}^\top A\bar{e} = n^{-1} \sum_v -d(v) = -d. \quad (8)$$

Proof. Write $|S| = s$ and $|T| = t$. Without loss of generality $s \geq t$. As in Lemma 5.3, we consider the characteristic vector $\chi = \chi_{S \cup T}$ and write $\chi = \sum_{i=1}^n a_i x_i$, where $\{x_i : 1 \leq i \leq n\}$ is an orthonormal basis of eigenvectors and x_i is the eigenvector for λ_i . Then $\sum_{i=1}^n a_i^2 = \chi^\top \cdot \chi = s + t$ $2e(S, T) = \chi^\top A\chi = \sum_{i=1}^n \lambda_i a_i^2$, and we estimate, with $\lambda = \lambda(G)$:

$$|2e(S, T) - \lambda_1 a_1^2 - \lambda_n a_n^2| = \left| \sum_{i \neq 1, n} \lambda_i a_i^2 \right| \leq \lambda(s + t) \leq \lambda n. \quad (9)$$

Next write $e = \sum_{i=1}^n c_i x_i$ and $\bar{e} = \sum_{i=1}^n \bar{c}_i x_i$, where $\sum c_i^2 = \sum \bar{c}_i^2 = 1$; since we can replace any eigenvector x_i by $-x_i$ we can assume that $c_1 > 0$ and $\bar{c}_n > 0$. We will show that x_1 is close to e and x_n is close to \bar{e} . Consider $z = Ae - de$, which has co-ordinates $z_i = n^{-1/2}(d(v_i) - d)$, and $\bar{z} = A\bar{e} + d\bar{e}$, which has co-ordinates $\bar{z}_i = \pm n^{-1/2}(d(v_i) - d)$, with positive sign when $v_i \in Y$ and negative sign when $v_i \in X$. Then by (ii),

$$z \cdot z = \bar{z} \cdot \bar{z} = n^{-1} \sum_v (d(v) - d)^2 = \text{Var}(G) \leq \beta d^2.$$

We also have $z = \sum(\lambda_i - d)c_i x_i$ and $\bar{z} = \sum(\lambda_i + d)\bar{c}_i x_i$, so

$$z \cdot z = \bar{z} \cdot \bar{z} = \sum c_i^2 (\lambda_i - d)^2 = \sum \bar{c}_i^2 (\lambda_i + d)^2.$$

Then $\sum_{i \neq 1} c_i^2 \leq (d - \lambda)^{-2} \beta d^2 \leq \beta \gamma^{-2}$ by (i), so $c_1 \geq c_1^2 \geq 1 - \beta \gamma^{-2}$, and

$$\|e - x_1\|^2 = (1 - c_1)^2 + \sum_{i \neq 1} c_i^2 = (1 - c_1)^2 + 1 - c_1^2 = 2(1 - c_1) \leq 2\beta \gamma^{-2}.$$

Similarly, one can prove that $\bar{c}_n \geq 1 - \beta \gamma^{-2}$ and $\|\bar{e} - x_n\|^2 \leq 2\beta \gamma^{-2}$. Next we estimate λ_1 and λ_n . Recall that $\lambda_1 = \|Ax_1\|$ and $d = \|de\|$, and so by the triangle inequality,

$$|\lambda_1 - d| = \left| \|Ax_1\| - \|de\| \right| \leq \|Ax_1 - de\| \leq \|Ae - de\| + \|Ae - Ax_1\|. \quad (10)$$

By (7) we have $\lambda_1 \geq d$, so we can write $\lambda_1 = (1 + q_1)d$ with $q_1 \geq 0$. Since $\|Ae - de\| = \|z\| < \beta^{1/2}d$ and $\|A(e - x_1)\| \leq \lambda_1 \|e - x_1\| \leq (2\beta)^{1/2} \gamma^{-1} \lambda_1$, from (10) we have $q_1 \leq \beta^{1/2} + (2\beta)^{1/2} \gamma^{-1} (1 + q_1)$. Thus, recalling that $\alpha = 4\beta^{1/2} \gamma^{-1} < 1/4$, we can estimate

$$q_1 \leq \frac{\beta^{1/2} + (2\beta)^{1/2} \gamma^{-1}}{1 - (2\beta)^{1/2} \gamma^{-1}} \leq \frac{\alpha/4 + \alpha/(2\sqrt{2})}{1 - 1/(8\sqrt{2})} < \alpha,$$

i.e.

$$d \leq \lambda_1 < (1 + \alpha)d.$$

Similarly, we have

$$-d \geq \lambda_n > -(1 + \alpha)d.$$

We also estimate $a_1 = \chi \cdot x_1$ by $\chi \cdot e = n^{-1/2}(s + t)$ and the inequality

$$\|\chi \cdot (e - x_1)\|^2 \leq \|\chi\|^2 \|e - x_1\|^2 \leq 2\beta \gamma^{-2}(s + t) < \alpha^2(s + t),$$

which gives

$$|a_1 - n^{-1/2}(s + t)| \leq \alpha(s + t)^{1/2}. \quad (11)$$

Similarly, we estimate $a_n = \chi \cdot x_n$ by $\chi \cdot \bar{e} = n^{-1/2}(s - t)$, and get

$$|a_n - n^{-1/2}(s - t)| \leq \alpha(s + t)^{1/2}. \quad (12)$$

Now we have the necessary ingredients to estimate $\lambda_1 a_1^2 + \lambda_n a_n^2$. Recall from (7) and (8) that $\lambda_1 \geq d$, $\lambda_n \geq -d(1 + \alpha)$ and $s \geq t$. By (11) and (12) we have the lower bounds

$$\begin{aligned} \lambda_1 a_1^2 &\geq d(n^{-1/2}(s + t) - \alpha(s + t)^{1/2})^2, \text{ and} \\ \lambda_n a_n^2 &\geq -d(1 + \alpha)(n^{-1/2}(s - t) + \alpha(s + t)^{1/2})^2. \end{aligned}$$

Thus we obtain

$$\begin{aligned} & \lambda_1 a_1^2 + \lambda_n a_n^2 \\ & \geq \frac{4d}{n}st - \frac{\alpha d}{n}(s-t)^2 - 2\alpha d \left(\frac{s+t}{n}\right)^{1/2} \left((s+t) + (1+\alpha)(s-t) \right) - \alpha^3 d(s+t). \end{aligned}$$

Since $s+t \leq n$ and $\alpha < 1/2$ we get

$$\lambda_1 a_1^2 + \lambda_n a_n^2 - \frac{4d}{n}st \geq -8\alpha dn.$$

The estimates for the upper bound are similar but slightly more technical. We use

$$\begin{aligned} \lambda_1 a_1^2 & \leq d(1+\alpha)(n^{-1/2}(s+t) + \alpha(s+t)^{1/2})^2, \text{ and} \\ \lambda_n a_n^2 & \leq \begin{cases} -d(n^{-1/2}(s-t) - \alpha(s+t)^{1/2})^2 & \text{if } \alpha(s+t)^{1/2} \leq n^{-1/2}(s-t), \\ 0 & \text{if } \alpha(s+t)^{1/2} > n^{-1/2}(s-t). \end{cases} \end{aligned}$$

In the case $\alpha(s+t)^{1/2} \leq n^{-1/2}(s-t)$ we have

$$\begin{aligned} & \lambda_1 a_1^2 + \lambda_n a_n^2 \\ & \leq \frac{4d}{n}st + \frac{\alpha d}{n}(s+t)^2 + 2\alpha d \left(\frac{s+t}{n}\right)^{1/2} \left((1+\alpha)(s+t) + (s-t) \right) + \alpha^3 d(s+t). \end{aligned}$$

In the case $\alpha(s+t)^{1/2} \geq n^{-1/2}(s-t)$ we have $(s+t)^2 - 4st = (s-t)^2 \leq \alpha^2 n(s+t) \leq \alpha^2 n^2$, so

$$\lambda_1 a_1^2 + \lambda_n a_n^2 \leq \lambda_1 a_1^2 \leq d(1+\alpha)(n^{-1}(4st + \alpha^2 n^2) + 2\alpha n^{-1/2}(s+t)^{3/2} + \alpha^2(s+t)).$$

Since $s+t \leq n$ and $\alpha < 1/4$, in both cases we obtain

$$\lambda_1 a_1^2 + \lambda_n a_n^2 - \frac{4d}{n}st \leq 8\alpha dn.$$

Combining this with (9) we obtain the stated estimate for $e(S, T)$. ■

5.3. Proof of Theorem 5.1

The idea of the proof is to use the connection between eigenvalues and closed walks. We can control the maximum degree by deleting few edges; then the main contribution to the upper bound on closed walks of length $2\ell + 2$ in Lemma 3.3 is $(1/2 + o(1))n^2\Delta^2$. This is very close to $\lambda_1^{2\ell+2} + \lambda_n^{2\ell+2} \sim 2d^{2\ell+2}$, so the other eigenvalues of this graph are small.

We now give the details. Suppose $\varepsilon > 0$. Consider any $\mathcal{C}_{2\ell}$ -free bipartite graph H on n vertices with parts X and Y and average degree $d \sim (n/2)^{1/\ell}$. Suppose $S \subseteq X$ and $T \subseteq Y$. By Lemma 4.2 there are $o(e(H))$ edges incident to vertices of degree at least $(1 + \varepsilon)d$. We remove these edges to obtain a graph G of maximum degree $\Delta \leq (1 + \varepsilon)d$ and average degree $d_G \sim (n/2)^{1/\ell}$. Also, as in the proof of Lemma 4.2, Proposition 3.9 gives $|X| \sim |Y| \sim \frac{n}{2}$. By Proposition 3.2 and Lemma 3.3

$$\frac{1}{n} \sum_{i=1}^n \lambda_i^{2\ell+2} = w_{2\ell+2}^\circ(G) < (1/2 + o(1))n\Delta^2 + (4\Delta)^{\ell+1}.$$

Now we use the estimates $\lambda_1 \geq d_G$ and $\lambda_n \leq -d_G$ from (7) and (8). Recalling that $\lambda = \max_{i \neq 1, n} |\lambda_i|$ we have

$$\lambda^{2\ell+2} \leq nw_{2\ell+2}^\circ(G) - 2d_G^{2\ell+2} < (1/2 + o(1))n^2\Delta^2 - 2d_G^{2\ell+2} + n(4\Delta)^{\ell+1}.$$

Since $d_G \sim (n/2)^{1/\ell}$ and $\Delta \leq (1 + \varepsilon)d$ this gives $\lambda^{2\ell+2} \leq ((1 + \varepsilon)^2 - 1 + o(1))2d_G^{2\ell+2}$. It follows that

$$\lambda \leq (6\varepsilon)^{1/(2\ell+2)}d_G + o(d_G).$$

Also, the variance is bounded as

$$\begin{aligned} \text{Var}(G) &= \frac{\sum_v (d(v) - d_G)^2}{n} \\ &= \frac{\sum_v d(v)^2}{n} - d_G^2 \leq \Delta \cdot \frac{\sum_v d(v)}{n} - d_G^2 \leq (1 + \varepsilon)d_G^2 - d_G^2 = \varepsilon d_G^2. \end{aligned}$$

We now apply Lemma 5.4 with $\beta = \varepsilon$ and $\gamma = 1 - (6\varepsilon)^{1/(2\ell+2)} + o(1)$. Recall that we need $\alpha = 4\beta^{1/2}\gamma^{-1} < 1/4$, which holds if ε is small. Thus

$$\left| e_G(S, T) - \frac{2d_G}{n} |S||T| \right| \leq (4\alpha d_G + \lambda/2)n,$$

where $\alpha = 4\beta^{1/2}\gamma^{-1} < 5\varepsilon^{1/2}$ for small ε . Recalling that G was obtained from H by deleting $o(n^{1+1/\ell})$ edges we have

$$\left| e_H(S, T) - \frac{2d_G}{n}|S||T| \right| < \left(20\varepsilon^{1/2} + (6\varepsilon)^{1/(2\ell+2)} + o(1) \right) n^{1+1/\ell}.$$

Since ε is arbitrary this proves Theorem 5.1. ■

6. Proof of Theorem 1.1

Suppose that G is a $\mathcal{C}_{2\ell} \cup \{C_k\}$ -free graph with n vertices with average degree $d = \Theta(n^{1/\ell})$, where $k > 2\ell$ is odd. For the first part of Theorem 1.1, we are required to find a bipartite graph $H \subset G$ such that $e(H) \geq d^{\ell+1} - o(n^{1+1/\ell})$. Similarly to the sketch given for 4-cycles, the idea is that we can take H to be the bipartite subgraph spanned by $N_\ell(v)$ and $N_{\ell+1}(v)$ for a suitable vertex v . The first step is to pass to a subgraph with low maximum degree.

Let $\Delta = n^{1/\ell+c}$, where $c := 1/2\ell^2$. Let S be the set of vertices of degree more than Δ , and let G_0 be the graph obtained by removing all edges of G containing at least one vertex of S , where G_0 has average degree d_0 . We will show that $d_0 \sim d$. To estimate the number of edges removed, recall from (1) that $\text{ex}(n, \mathcal{C}_{2\ell}) \leq \frac{1}{2}n^{1+1/\ell} + O(n)$, so

$$|S| \leq \frac{n^{1+1/\ell} + O(n)}{\Delta} < m := 2n^{1-c}.$$

It follows that

$$e(S) \leq \frac{1}{2}m^{1+1/\ell} + O(m) < n^{1+1/\ell-c}.$$

Also Proposition 3.9 gives

$$e(S, V(G) \setminus S) < (mn)^{1/2+1/2\ell} + n < n^{1+1/\ell-c/2}.$$

In particular, we have $e(G_0) > e(G) - o(n^{1+1/\ell})$, and therefore $d_0 \sim d$. For the remainder of the proof we work in the graph G_0 , which has maximum degree at most Δ .

Next, by Lemma 3.6, we can choose a vertex v that is the start of at least $d_0^{\ell+1} - (\ell + 1)^2 \Delta^\ell = d^{\ell+1} + o(n^{1+1/\ell})$ paths of length $\ell + 1$ in G_0 . We claim that all but $o(n^{1+1/\ell})$ of these paths reach $N_{\ell+1}(v)$. Consider a breadth-first search tree T rooted at v . Consider any path P of length $\ell + 1$ that does not reach $N_{\ell+1}(v)$. Then there is a smallest i such that the $(i + 1)$ st edge of P does not go from $N_i(v)$ to $N_{i+1}(v)$. By construction of T this edge

must either go from $N_i(v)$ to $N_{i-1}(v)$ or lie within $N_i(v)$. The first case is impossible, as any edge of $E(G_0) \setminus E(T)$ between $N_{i-1}(v)$ and $N_i(v)$ would create an even cycle of length at most 2ℓ . For the second case we recall that Lemma 3.7 implies that $G[N_i(v)]$ has average degree at most $2k$ for any $i \leq \ell$, since $k > 2\ell$. Also note that from maximum degree assumption we have that $|N_i(v)| \leq \Delta^i$. This gives at most $k|N_i(v)| \leq k\Delta^i$ choices for the $(i+1)$ st edge of P . Let w be the first vertex of P in $N_i(v)$. The subpath of P from v to w is uniquely determined (otherwise we would have an even cycle of length at most 2ℓ). Then we have at most Δ choices for each of the $\ell - i$ subsequent edges of P . In total, since there are ℓ choices of i , the number of choices for P is at most $k\ell\Delta^\ell = o(n^{1+1/\ell})$, as required.

Each edge between $N_\ell(v)$ and $N_{\ell+1}(v)$ is contained in at most one path of length $\ell + 1$ from v to $N_{\ell+1}(v)$, otherwise we would have an even cycle of length at most 2ℓ . Thus, taking H to be the bipartite graph of edges between $N_\ell(v)$ and $N_{\ell+1}(v)$, we have $e(H) = d^{\ell+1} + o(n^{1+1/\ell})$, as required.

To prove the second part of Theorem 1.1, suppose that $d \geq (1 + o(1)) \cdot (n/2)^{1/\ell}$. The number of edges in the bipartite graph H constructed above is at least $(1 + o(1))nd^{\ell+1}/2 \geq (1 + o(1))(n/2)^{1+1/\ell}$. On the other hand, from Proposition 3.8 we have $e(H) \leq (1 + o(1))(n/2)^{1+1/\ell}$. Therefore $e(H) \sim (n/2)^{1+1/\ell}$ and $d \leq (1 + o(1))(n/2)^{1/\ell}$. Since $d \geq (1 + o(1))(n/2)^{1/\ell}$, we also have $d \sim (n/2)^{1/\ell}$. So $e(H) \sim e(G)$ and by Proposition 3.8, this shows $e(G) \sim z(n, \mathcal{C}_{2\ell})$. This completes the proof. ■

7. Proof of Theorem 1.2

Suppose that n is large, and G is a graph on n vertices with $e(G) \geq (q+1)n/2$, where $q \in \mathbb{R}^+$ is defined by $n = 2(q^\ell + q^{\ell-1} + \dots + 1)$. Suppose also that G does not contain an even cycle of length at most 2ℓ , or C_k for some odd $k > 2\ell + 2$. We will show that $e(G) = (q+1)n/2$ and G is bipartite. Then Proposition 3.8 characterizes equality, namely, G must be the incidence graph of a generalized polygon. We start by considering the case $\ell \geq 2$ and $k > 2\ell + 2$.

Case 1: $\ell \geq 2$ and $k > 2\ell + 2$. Let H be a bipartite subgraph of G with maximum size. We show $G = H$. By Theorem 1.1, $e(H) \sim e(G) \sim (n/2)^{1+1/\ell}$. Furthermore, maximality of H implies that $\delta(H) > \delta(G)/2$, as if there were a vertex of degree less than $\delta(G)/2$ in H we could move it to the other part and increase the number of edges in H . By Section 4.3, we can assume $\delta(G) > q/4$, so $\delta(H) > q/8$. Label the parts of H as X_0 and X_1 , with $X_0 \cup X_1 = V(G)$. Suppose for a contradiction that $G[X_0]$ contains an edge xy . Let z be a

neighbor of y in X_1 . We greedily construct a sequence of mutually disjoint sets $\{y\}$ and S_x^i and S_z^i for $0 \leq i \leq \ell$, where $S_x^0 = \{x\}, S_z^0 = \{z\}$,

$$S_x^i \subseteq N(S_x^{i-1}) \cap X_{i \bmod 2} \quad \text{and} \quad S_z^i \subseteq N(S_z^{i-1}) \cap X_{i+1 \bmod 2}.$$

Here $N(S)$ denotes the set of vertices of $V(H) \setminus S$ with at least one neighbor in S . Note that by definition we have $S_x^i \subseteq N_i(x)$ and $S_z^i \subseteq N_i(z)$. By consideration of breadth first search trees as in the proof of Theorem 1.1, every vertex in $N_i(x)$ has exactly one neighbor in $N_{i-1}(x)$ and $e(N_i(x)) < k|N_i(x)|$ for every $1 \leq i \leq \ell - 1$. Moreover, two distinct vertices of $N_i(x)$ can not have a common neighbor in $N_{i+1}(x)$. Similar statements hold for z . By the minimum degree assumption in H , we have $|N_i(x)|, |N_i(z)| \geq (q/8)^i$ for all $0 \leq i \leq \ell$. Since $q \sim (n/2)^\ell$, this allows us greedily to choose disjoint sets S_x^i, S_z^i so that

$$|S_x^i| \sim |S_z^i| \sim (cn)^{i/\ell},$$

for all $1 \leq i \leq \ell$ and some constant $c > 0$. Now we apply Theorem 5.1 to H with $S = S_x^\ell$ and $T = S_z^\ell$. This gives

$$e(S, T) \geq \frac{(1 + o(1))2(n/2)^{1/\ell}}{n} |S||T| - o(n^{1+1/\ell}) > 2kn.$$

In particular, we can find a path of length $k - 2\ell - 2$ using only edges between S and T . By construction this can be completed to a cycle of length k in G , which is a contradiction. We deduce that X_0 is an independent set in G . Similarly X_1 is independent, so $G = H$. This completes the proof for $k > 2\ell + 2$.

For the remainder of the proof we consider the special case $\ell = 2$ and $k = 5$.

Case 2: $\ell = 2$ and $k = 5$. We use a similar vertex deletion argument to that in Section 4.3 to find a subgraph G_t of G with $t \geq 0.01n$ vertices and $\delta(G_t) > 0.51t^{1/2}$. Starting with $G = G_n$, we produce a graph G_i with i vertices by deleting a vertex of G_{i+1} of degree at most $0.51(i+1)^{1/2}$ for each $i < n$. After t steps, the total number of edges deleted is less than

$$\sum_{i=t+1}^n 0.51i^{1/2} < 0.51 \int_{t+1}^{n+1} x^{1/2} dx = \frac{1.02}{3} ((n+1)^{3/2} - (t+1)^{3/2}).$$

Suppose that we fail to find a subgraph of minimum degree more than $0.51t^{1/2}$ in $n - t \leq 0.99n$ steps. Then we have a graph G_t with $t \geq 0.01n$ vertices and

$$e(G_t) > (q+1)n/2 - \frac{1.02}{3} ((n+1)^{3/2} - (t+1)^{3/2})$$

$$\begin{aligned}
 &> \frac{1}{2\sqrt{2}}n^{3/2} - \frac{1.02}{3}((n+1)^{3/2} - (t+1)^{3/2}) \\
 &\geq \left(\frac{1}{2\sqrt{2}} - \frac{1.02}{3} - o(1)\right)n^{3/2} > 0.01n^{3/2} \geq 10t^{3/2}.
 \end{aligned}$$

This contradicts Theorem 1.1, provided $t \geq 0.01n$ is large enough. So G has a subgraph G_t with t vertices and $\delta(G_t) > 0.51t^{1/2}$ and where $t \geq 0.01n$. For $i \leq n$ let q_i be the unique positive real defined by $i = 2(q_i^2 + q_i + 1)$, so that $q_n = q$. Then $q_i = (\sqrt{2i-3} - 1)/2$, and if $t < n$ then

$$e(G_{n-1}) \geq e(G) - 0.51n^{1/2} \geq (q+1)n/2 - 0.51n^{1/2} > (q_{n-1} + 1)(n-1)/2$$

for large enough n , using

$$\begin{aligned}
 &(q+1)n/2 - (q_{n-1} + 1)(n-1)/2 \\
 &= (\sqrt{2n-3} - 1)n/4 - (\sqrt{2n-5} - 1)(n-1)/4 + 1/2 \\
 &= \sqrt{2n-5}/4 + (\sqrt{2n-3} - \sqrt{2n-5})n/4 + 1/4 \sim \frac{3}{4\sqrt{2}}n^{1/2} > 0.51n^{1/2}.
 \end{aligned}$$

Repeating this calculation, we get that

$$\text{if } t < n \text{ then } e(G_t) > (q_t + 1)t/2, \tag{13}$$

provided $t \geq 0.01n$ is large enough. We will show that G_t is bipartite.

First we pass to a maximum bipartite subgraph H of G_t as in Case 1, with parts X_0 and X_1 . By Theorem 1.1, $e(H) \sim e(G_t) \sim (t/2)^{3/2}$. We claim that no vertex $x \in X_0$ has more than $0.009t^{1/2}$ neighbors in X_0 , and similarly for X_1 . To see this, note that such an $x \in X_0$ also has more than $0.009t^{1/2}$ neighbors in X_1 by maximality of H . Then $N_2(x)$ contains $\Theta(n)$ vertices of X_0 and $\Theta(n)$ vertices of X_1 . Choose a set S of $\Theta(n)$ vertices of $N_2(x) \cap X_0$ and a set T of $\Theta(n)$ vertices of $N_2(x) \cap X_1$ such that for each $w \in S$ and $z \in T$, there exist paths of length two from x to w and from x to z which share only the vertex x . Then in H we apply Theorem 5.1 (pseudorandomness) to conclude

$$e(S, T) \geq \frac{(1 + o(1))2(t/2)^{1/2}}{t} |S||T| - o(t^{3/2}).$$

In particular, $e(S, T) \neq 0$ and evidently there is a cycle of length five through x and any edge between S and T , a contradiction. Therefore no vertex has more than $0.009t^{1/2}$ vertices in its own part. It follows that every vertex has degree at least $0.501t^{1/2}$ in H .

We next claim that $|X_0| \sim |X_1| \sim t/2$. First note from Proposition 3.9 that

$$e(H) \leq z(t, C_4) \leq (|X_0||X_1|)^{3/4} + \max\{|X_0|, |X_1|\}.$$

On the other hand, $e(H) \sim e(G) \sim (t/2)^{3/2}$, and so we see

$$(|X_0||X_1|)^{3/4} \geq (1 + o(1))(t/2)^{3/2}.$$

Since $|X_0| + |X_1| = t$, and we just observed $|X_0||X_1| \geq (1 + o(1))(t/2)^2$, we conclude $|X_0| \sim |X_1| \sim t/2$.

Now we show $G_t[X_0]$ and $G_t[X_1]$ have no edges, so that G_t is bipartite. Suppose for a contradiction that G_t has an edge xy with $x, y \in X_0$. Note that x and y have at most one common neighbour, since G is C_4 -free. Let z be this common neighbour if it exists, or an arbitrary vertex otherwise. Let S be the set of ends of paths of length two in H that start at x and avoid $\{y, z\}$. Let T be the set of ends of paths of length two in H that start at y and avoid $\{x, z\}$. Since H has minimum degree more than $0.501t^{1/2}$, each of S and T have size at least $(0.501t^{1/2} - 1)^2 > |X_0|/2$, provided t is large enough. Then since $S, T \subset X_0$ there is a vertex $w \in S \cap T$. Thus we have paths xaw and ybw , where $a \neq b$ since our paths avoid z . However, $xawby$ forms a 5-cycle, so we have a contradiction. We conclude $G_t[X_0]$ is empty, and similarly, $G_t[X_1]$ is empty, so G_t is bipartite.

To complete the proof, recall that $e(G_t) \leq (q_t + 1)t/2$ by Proposition 3.8. However by (13), $e(G) > (q_t + 1)t/2$ for $t < n$. Thus we must have $t = n$ and $e(G_t) = e(G) = (q + 1)n/2$, so $G_t = G$. Therefore G itself is bipartite, and by Proposition 3.8, G is the bipartite incidence graph of a projective plane. ■

8. Proof of Theorem 1.3

Let $\ell \geq 2$ and $k \geq 4\ell + 1$ be odd and $c > 0$. Suppose that G is a $\mathcal{C}_{2\ell} \cup \{C_k\}$ -free graph on n vertices with minimum degree at least $cn^{1/\ell}$. We need to show that $\chi(G) < (4k)^{\ell+1}/c^\ell$. We use the approach of Thomassen [37]. For $r \geq 0$ write $N_{\leq r}(v) = N_0(v) \cup N_1(v) \cup \dots \cup N_r(v)$. Consider a maximal sequence of vertices v_1, v_2, \dots, v_s such that $N_{\leq \ell}(v_1), \dots, N_{\leq \ell}(v_s)$ are pairwise disjoint. By Lemma 3.7, since $k \geq 2\ell + 1$, for any $v \in V(G)$ and $r \leq \ell$ we have $e(N_r(v)) \leq k|N_r(v)|$. Since G is $\mathcal{C}_{2\ell}$ -free, no vertex of $N_r(v)$ has more than one neighbor in $N_{r-1}(v)$, so

$$e(N_{\leq r}(v)) \leq (k + 1)|N_{\leq r}(v)| < 2k|N_{\leq r}(v)|.$$

On the other hand, since G has minimum degree at least $cn^{1/\ell}$,

$$e(N_{\leq r}(v)) \geq \frac{1}{2}cn^{1/\ell}|N_{\leq r-1}(v)|.$$

We conclude that $|N_{\leq r}(v)| > \frac{1}{4k}cn^{1/\ell}|N_{\leq r-1}(v)|$ for all $r \leq \ell$, which implies

$$|N_{\leq \ell}(v)| > \frac{c^\ell}{(4k)^\ell}n$$

for every vertex $v \in V(G)$. In particular, this holds for v_1, v_2, \dots, v_s , so $s < (4k)^\ell/c^\ell$. The maximality of s implies that any vertex v is within distance 2ℓ of some v_i . Now we use the assumption that $k \geq 4\ell + 1$. The proof of Lemma 3.7 shows that for $r \leq 2\ell$, every subset of $N_r(v)$ induces a subgraph of average degree less than $2k$. It follows that $G[N_r(v)]$ has chromatic number at most $2k$. Furthermore, $G[N_1(v) \cup N_3(v) \cup \dots \cup N_{2\ell-1}(v)]$ has chromatic number at most $2k$, since there are no edges between $N_i(v)$ and $N_{i+2}(v)$ for any i , and similarly $G[N_0(v) \cup N_2(v) \cup \dots \cup N_{2\ell}(v)]$ also has chromatic number at most $2k$. Therefore $G[N_{\leq 2\ell}(v)]$ has chromatic number at most $4k$, so we can cover $N_{\leq 2\ell}(v_i)$ by at most $4k$ independent sets for $1 \leq i \leq s$. It follows that

$$\chi(G) \leq 4ks < \frac{(4k)^{\ell+1}}{c^\ell}.$$

This completes the proof of Theorem 1.3. ▀

9. Concluding remarks

- Our stability approach not only gives extremal results but describes the approximate structure of nearly extremal graphs. We only needed these results in the bipartite (Zarankiewicz) setting, but we note that very similar arguments give analogous results in the non-bipartite (Turán) setting. For example, we have the following result. Suppose G is a C_4 -free graph on n vertices with average degree $d \sim \sqrt{n}$. Then for any $S, T \subseteq V(G)$ we have $e(S, T) = \frac{d}{n}|S||T| + o(n^{3/2})$. The proof is very similar to that of Theorem 5.1. First we control the maximum degree as $\Delta < (1 + \varepsilon)d$ by deleting $O(n) = o(n^{3/2})$ edges. Then the argument of Lemma 3.3 shows that $w_6^\circ(G) < (1 + o(1))n\Delta^2$; the only difference is that there are $n - 1$ choices for u rather than $n/2 + o(n)$. On the other hand, $w_6^\circ(G) = \frac{1}{n} \sum \lambda_i^6$ has a contribution of $d^6/n \sim n^2$ from the first eigenvalue, so the other eigenvalues are $o(d)$ as $\varepsilon \rightarrow 0$. The pseudorandomness property now follows from the non-bipartite version of Lemma 5.4, which is given in [25, Section 2.4].

- No result similar to Lemma 4.1 can hold for C_6 -free graphs: in fact by the results of [20], there exist $\delta, \varepsilon > 0$ such that any extremal C_6 -free graph G with average degree d has at least $\delta e(G)$ edges containing a vertex of degree more than $(1 + \varepsilon)d$.
- We proved the even girth result Theorem 1.2 under the assumption that the forbidden odd cycle length satisfies $k \geq 2\ell + 3$. The polarity graphs show that no such result holds for $3 \leq k \leq \ell$, but some values of k remain open. In particular, one might think that the case $k = 2\ell + 1$ should be approachable by the methods we used to handle $\{C_4, C_5\}$ -free graphs. However, the vertex deletion method does not give sufficient minimum degree for a straightforward adaptation of this argument, so other ideas are needed.
- The polarity graphs have arbitrarily large chromatic numbers, as shown by estimates on their independence numbers by Godsil and Newman [22]. Thus Theorem 1.3 does not hold when $(k, \ell) \in \{(3, 2), (3, 3), (3, 5), (5, 5)\}$, but it seems likely that it should hold when $k \geq 2\ell + 1$. We also remark that the bound $(4k)^{\ell+1}/c^\ell$ is unlikely to be the correct dependence of $\chi(G)$ on c , and it would be interesting to determine this even for $\ell = 2$.
- We remarked earlier that there is little known about the approximate structure of nearly extremal graphs. In the case of 4-cycles, Füredi [17,18] showed that any C_4 -free graph on $q^2 + q + 1$ vertices has at most $\frac{1}{2}q(q+1)^2$ edges, and for large q equality can only hold for polarity graphs. However, we do not know whether all nearly extremal graphs are structurally close to polarity graphs. Such an understanding would probably have implications for the conjecture of Erdős that $\text{ex}(n, \{C_3, C_4\}) \sim z(n, C_4)$, and for other seemingly more basic questions, such as whether a graph on n vertices with at least $\text{ex}(n, C_4) + 1$ edges must contain many 4-cycles.
- We also considered more generally the problem of determining bipartite graphs F and odd integers k for which $\text{ex}(n, \{F, C_k\}) \sim z(n, F)$ and developed different methods to attack this problem when F is not an even cycle. This will be the subject of a second paper, co-authored with P. Allen.

Acknowledgement. We thank the anonymous referees for helpful comments.

References

- [1] P. ALLEN: Dense H -free graphs are almost $(\chi(H) - 1)$ -partite, *Electronic Journal of Combinatorics* **17** (2010), R21.
- [2] P. ALLEN, J. BÖTTCHER, S. GRIFFITHS, Y. KOHAYAKAWA and R. MORRIS: The chromatic thresholds of graphs, [arXiv:1108.1746](https://arxiv.org/abs/1108.1746).
- [3] N. ALON, S. HOORY and N. LINIAL: The Moore bound for irregular graphs, *Graphs Combin.* **18** (2002), 53–57.

- [4] N. ALON and J. SPENCER: *The Probabilistic Method*, 3rd edition, Wiley, New York, 2008.
- [5] N. ALON and B. SUDAKOV: H -free graphs of large minimum degree, *Electron. J. Combin.* **13** (2006), Research Paper 19, 9.
- [6] B. ANDRÁSFAI, P. ERDŐS and V. SÓS: On the connection between chromatic number, maximal clique and minimal degree of a graph, *Disc. Math.* **8** (1974), 205–218.
- [7] C.T. BENSON: Minimal regular graphs of girths eight and twelve, *Canad. J. Math.* **18** (1966), 1091–1094.
- [8] G. R. BLAKLEY and P. ROY: A Hölder type inequality for symmetric matrices with nonnegative entries, *Proc. Amer. Math. Soc.* **16** (1965), 1244–1245.
- [9] J. BONDY and M. SIMONOVITS: Cycles of even length in graphs, *J. Combin. Theory Ser. B* **16** (1974), 75–105.
- [10] P. ERDŐS: On sequences of integers no one of which divides the product of two others and on some related problems, *Mitt. Forsch.-Inst. Math. Mech. Univ. Tomsk* **2** (1938), 74–82.
- [11] P. ERDŐS, P.: Some recent progress on extremal problems in graph theory, *Congr. Numer.* **14** (1975), 3–14.
- [12] P. ERDŐS and A. RÉNYI: On a problem in the theory of graphs, *Publ. Math. Inst. Hungar. Acad. Sci.* **7A** (1962), 623–641.
- [13] P. ERDŐS and M. SIMONOVITS: Compactness results in extremal graph theory, *Combinatorica* **2** (1982), 275–288.
- [14] P. ERDŐS and M. SIMONOVITS: A limit theorem in graph theory, *Studia Sci. Math. Hungar.* **1** (1966), 51–57.
- [15] P. ERDŐS and A. H. STONE: On the structure of linear graphs, *Bull. Amer. Math. Soc.* **52** (1946), 1087–1091.
- [16] W. FEIT and G. HIGMAN: The nonexistence of certain generalized polygons, *J. Algebra* **1** (1964), 114–131.
- [17] Z. FÜREDI: Quadrilateral-free graphs with maximum number of edges, unpublished manuscript (1988).
- [18] Z. FÜREDI: On the number of edges of quadrilateral-free graphs, *J. Combin. Theory Ser. B* **68** (1996), 1–6.
- [19] Z. FÜREDI: New asymptotics for bipartite Turán numbers, *J. Combin. Theory Ser. A* **75** (1996), 141–144.
- [20] Z. FÜREDI, A. NAOR and J. VERSTRAËTE: On the Turán number for the hexagon, *Adv. Math.* **203** (2006), 476–496.
- [21] W. GODDARD and J. LYLE: Dense graphs with small clique number, *J. Graph Theory* **66** (2011), 319–331.
- [22] C. D. GODSIL and M. W. NEWMAN: Eigenvalue bounds for independent sets, *J. Combin. Theory Ser. B* **98** (2008), 721–734.
- [23] S. HOORY: The size of bipartite graphs with a given girth, *J. Combin. Theory Ser. B* **86** (2002), 215–220.
- [24] T. KÖVÁRI, V. SÓS and P. TURÁN: On a problem of K. Zarankiewicz, *Colloquium Math.* **3** (1954), 50–57.
- [25] M. KRIVELEVICH and B. SUDAKOV: Pseudo-random graphs, in: *More Sets, Graphs and Numbers*, *Bolyai Soc. Math. Studies* **15**, Springer (2006), 199–262.
- [26] T. LAM and J. VERSTRAËTE: A note on graphs without short even cycles, *Electron. J. Combin.* **12** (2005), N5.

- [27] F. LAZEBNIK, V. A. USTIMENKO and A. J. WOLDAR: A new series of dense graphs of high girth, *Bulletin of the AMS* **32** (1995), 73–79.
- [28] F. LAZEBNIK, V. A. USTIMENKO and A. J. WOLDAR: Polarities and $2k$ -cycle-free graphs, *Disc. Math.* **197/198** (1999), 503–513.
- [29] A. LUBOTZKY, R. PHILLIPS and P. SARNAK: Ramanujan graphs, *Combinatorica* **8** (1988), 261–277.
- [30] T. ŁUCZAK and S. THOMASSÉ: Coloring dense graphs via VC-dimension, preprint.
- [31] G. MARGULIS: Explicit constructions of graphs without short cycles and low density codes, *Combinatorica* **2** (1982), 71–78.
- [32] A. NAOR and J. VERSTRAËTE: A note on bipartite graphs without $2k$ -cycles, *Combin. Probab. Comput.* **14** (2005), 845–849.
- [33] T. D. PARSONS: Graphs from projective planes, *Aequationes Math.* **14** (1976), 167–189.
- [34] I. REIMAN: Über ein Problem von K. Zarankiewicz, *Acta Math. Acad. Sci. Hungar.* **9** (1958), 269–278.
- [35] A. SIDORENKO: Cycles in graphs and functional inequalities, *Mat. Zametki* **46** (1989), 72–79; translation in *Math. Notes* **46** (1989), 877–882.
- [36] R. SINGLETON: On minimal graphs of maximum even girth, *J. Combin. Theory* **1** (1966), 306–332.
- [37] C. THOMASSEN: On the chromatic number of pentagon-free graphs of large minimum degree, *Combinatorica* **27** (2007), 241–243.
- [38] J. TITS: Sur la trichotomie et certains groupes qui s'en déduisent, *Inst. Hautes Études Sci. Publ. Math.* **2** (1959), 13–60.
- [39] P. TURÁN: Eine Extremalaufgabe aus der Graphentheorie, *Mat. Fiz. Lapok* **48** (1941), 436–452.
- [40] J. VERSTRAËTE: On arithmetic progressions of cycle lengths in graphs, *Combin. Probab. Comput.* **9** (2000), 369–373.

Peter Keevash

School of Mathematical Sciences
Queen Mary University of London, UK
 p.keevash@qmul.ac.uk

Benny Sudakov

Department of Mathematics
University of California, Los Angeles
 CA, USA
 bsudakov@math.ucla.edu

Jacques Verstraëte

Department of Mathematics
University of California, San Diego, CA,
 USA
 jverstra@math.ucsd.edu