

Getting a Directed Hamilton Cycle Two Times Faster

CHOONGBUM LEE,[†] BENNY SUDAKOV[‡] and DAN VILENCHIK

Department of Mathematics, UCLA, Los Angeles, CA 90095, USA
(e-mail: choongbum.lee@gmail.com, {bsudakov,vilenchik}@math.ucla.edu)

Received 28 March 2011; revised 31 March 2012; first published online 4 May 2012

Consider the random graph process where we start with an empty graph on n vertices and, at time t , are given an edge e_t chosen uniformly at random among the edges which have not appeared so far. A classical result in random graph theory asserts that w.h.p. the graph becomes Hamiltonian at time $(1/2 + o(1))n \log n$. On the contrary, if all the edges were directed randomly, then the graph would have a directed Hamilton cycle w.h.p. only at time $(1 + o(1))n \log n$. In this paper we further study the directed case, and ask whether it is essential to have twice as many edges compared to the undirected case. More precisely, we ask if, at time t , instead of a random direction one is allowed to choose the orientation of e_t , then whether or not it is possible to make the resulting directed graph Hamiltonian at time earlier than $n \log n$. The main result of our paper answers this question in the strongest possible way, by asserting that one can orient the edges on-line so that w.h.p. the resulting graph has a directed Hamilton cycle exactly at the time at which the underlying graph is Hamiltonian.

AMS 2010 *Mathematics subject classification*: Primary 05C20, 05C45, 05C80

1. Introduction

The celebrated *random graph process*, introduced by Erdős and Rényi [12] in the 1960s, begins with an empty graph on n vertices, and at every round $t = 1, \dots, m$ adds to the current graph a single new edge chosen uniformly at random out of all missing edges. This distribution is commonly denoted as $G_{n,m}$. An equivalent ‘static’ way of defining $G_{n,m}$ would be to choose m edges uniformly at random out of all $\binom{n}{2}$ possible ones. One advantage in studying the random graph process, rather than the static model, is that it allows for a higher-resolution analysis of the appearance of monotone graph properties (a graph property is monotone if it is closed under edge addition).

[†] Research supported in part by a Samsung Scholarship.

[‡] Research supported in part by NSF grant DMS-1101185, NSF CAREER award DMS-0812005 and by a USA–Israeli BSF grant.

A *Hamilton cycle* of a graph is a simple cycle that passes through every vertex of the graph, and a graph containing a Hamilton cycle is called *Hamiltonian*. Hamiltonicity is one of the most fundamental notions in graph theory, and has been extensively studied in various contexts, including random graphs. The earlier results on Hamiltonicity of random graphs were obtained by Pósa [21], and Korshunov [18]. Improving on these results, Komlós and Szemerédi [17] proved that if $m' = \frac{1}{2}n \log n + \frac{1}{2}n \log \log n + c_n n$, then

$$\lim_{n \rightarrow \infty} \mathbb{P}(G_{n,m'} \text{ is Hamiltonian}) = \begin{cases} 0 & \text{if } c_n \rightarrow -\infty, \\ e^{-e^{-2c}} & \text{if } c_n \rightarrow c, \\ 1 & \text{if } c_n \rightarrow \infty. \end{cases}$$

One obvious necessary condition for the graph to be Hamiltonian is for the minimum degree to be at least 2, and surprisingly, the probability of $G_{n,m'}$ having minimum degree two at time m' has the same asymptotic behaviour as the probability of it being Hamiltonian. Bollobás [7] strengthened this observation by proving that w.h.p. the random graph process becomes Hamiltonian when the last vertex of degree one disappears. Moreover, Bollobás, Fenner and Frieze [8] described a polynomial time algorithm which w.h.p. finds a Hamilton cycle in random graphs.

Hamiltonicity has been studied for directed graphs as well. Consider a *random directed graph process* where, at time t , a random directed edge is chosen uniformly at random among all missing edges, and let $D_{n,m}$ be the graph consisting of the first m edges. Frieze [15] proved that for $m' = n \log n + c_n n$, the probability of $D_{n,m'}$ containing a (directed) Hamilton cycle is

$$\lim_{n \rightarrow \infty} \mathbb{P}(D_{n,m'} \text{ is Hamiltonian}) = \begin{cases} 0 & \text{if } c_n \rightarrow -\infty, \\ e^{-2e^{-c}} & \text{if } c_n \rightarrow c, \\ 1 & \text{if } c_n \rightarrow \infty. \end{cases}$$

As for the undirected case, this probability has the same asymptotic behaviour as the probability of the directed graph having minimum in-degree and out-degree 1. In fact, Frieze proved [15] that when the last vertex to have in- or out-degree less than one disappears, the graph has a Hamilton cycle w.h.p.

Hamiltonicity of various other random graph models has also been studied [22, 3]. One model which will be of particular interest to us is the k -in k -out model, in which every vertex chooses k in-neighbours and k out-neighbours uniformly at random and independently of the others. Improving on several previous results, Cooper and Frieze [10] proved that a random graph in this model is Hamiltonian w.h.p. when $k = 2$ (which is best possible since it is easy to see that a 1-in 1-out random graph is w.h.p. not Hamiltonian).

1.1. Our contribution

The results of Bollobás [7] and Frieze [15] introduced above suggest that the main obstacle to Hamiltonicity of random graphs lies in ‘reaching’ certain minimum degree conditions. It is therefore natural to ask how the thresholds change if we modify the random graph process so that we can somehow bypass this obstacle.

We consider the following process suggested by Frieze [16], which was designed for this purpose. Starting from the empty graph, at time t , an undirected edge (u, v) is given uniformly at random out of all missing edges, and a choice of its orientation ($u \rightarrow v$ or $v \rightarrow u$) is to be made at the time of its arrival. In this process, we can attempt to accelerate the appearance of monotone directed graph properties, or delay them, by applying an appropriate on-line algorithm. It is important to stress that the process is *on-line* in nature, namely, we cannot see any future edges at the current round and are forced to make the choice based only on the edges seen so far. In this paper, we investigate the property of containing a directed Hamilton cycle by asking the question: Can one speed up the appearance of a directed Hamilton cycle? The best we can hope for is to obtain a directed Hamilton cycle at the time when the underlying graph has minimum degree 2. The following result asserts that directed Hamiltonicity is in fact achievable exactly at that time, and this answers the above question positively in the strongest possible way.

Theorem 1.1. *Let \mathcal{G} be a random (undirected) graph process that terminates when the last vertex of degree one disappears. There exists an on-line algorithm **Orient** that orients the edges of \mathcal{G} so that with high probability the resulting directed graph is Hamiltonian.*

Let us note that \mathcal{G} w.h.p. contains $(1 + o(1))n \log n / 2$ edges, in contrast with $(1 + o(1))n \log n$ edges in the random directed graph model. Thus the required number of random edges is reduced by half.

Our model is similar in spirit to the so-called Achlioptas process. It is well known that a giant connected component (*i.e.*, a component of linear size) appears in the random graph $G_{n,m}$ when $m = (1 + o(1))n/2$. Inspired by the celebrated ‘power of two choices’ result [2], Achlioptas posed the following question. Suppose that edges arrive in pairs, that is, at round t the pair of edges (e_t, e'_t) chosen uniformly at random is given, and one is allowed to pick an edge out of it for the graph (the other edge will be discarded). Can one delay the appearance of the giant component? Bohman and Frieze answered this question positively [4] by describing an algorithm whose choice rule allows for the ratio $m/n \geq 0.53$, and this ratio has since been improved [5]. Quite a few papers have thereafter studied various related problems that arise in the above model [6, 14, 19, 23, 24]. As an example, Krivelevich, Loh and Sudakov [19] studied the question: How long can one delay the appearance of a certain fixed subgraph?

One such paper, closely related to our work, is the recent paper by Krivelevich, Lubetzky and Sudakov [20]. They studied the Achlioptas process for Hamiltonicity, and proved that by exploiting the ‘power of two choices’, one can construct a Hamilton cycle at time $(1 + o(1))n \log n / 4$, which is twice as fast as in the random case. Both our result and their result suggest that the ‘bottleneck’ to Hamiltonicity of random graphs indeed lies in the minimum degree, and thus these results can be understood in the context of complementing the results of Bollobás [7] and Frieze [15].

1.2. Preliminaries

The paper is rather involved technically. One factor that contributes to this is that we are establishing the ‘hitting time’ version of the problem. That is, we determine the exact

threshold for the appearance of a Hamilton cycle. The analysis can be simplified if one only wishes to estimate this threshold asymptotically (see Section 7). To make the current analysis more approachable without risking any significant change to the random model, we consider the following variant of the graph process, which we call the *random edge process*: at time t , an edge is given as an ordered pair of vertices $e_t = (v_t, w_t)$ chosen uniformly at random, with repetition, from the set of all possible n^2 ordered pairs (note that this model allows loops and repeated edges). In what follows, we use G_t to denote the graph induced by the first t edges, and given the orientation of each edge, use D_t to denote the directed graph induced by the first t edges. By m_* we denote the time t when the last vertex of degree one in G_t becomes a degree two vertex.

We will first prove that there exists an on-line algorithm **Orient** which w.h.p. orients the edges of the graph G_{m_*} so that the resulting directed graph D_{m_*} is Hamiltonian, and then in Section 6 we show how Theorem 1.1 can be recovered from this result.

1.3. Organization of the paper

In the next section we describe the algorithm **Orient** that is used to prove Theorem 1.1 (in the modified model). Then in Section 3 we outline the proof of Theorem 1.1. Section 4 describes several properties that a typical random edge process possesses. Using these properties we prove Theorem 1.1 in Section 5. Then in Section 6, we show how to modify the algorithm **Orient**, in order to make it work for the original random graph process.

Notation. A *directed 1-factor* is a directed graph in which every vertex has in-degree and out-degree exactly 1, and a *1-factor* of a directed graph is a spanning subgraph which is a directed 1-factor. The function $\exp(x) := e^x$ is the exponential function. Throughout the paper $\log(\cdot)$ denotes the natural logarithm. For the sake of clarity, we often omit floor and ceiling signs whenever these are not crucial, and we make no attempt to optimize our absolute constants. We also assume that the order n of all graphs tends to infinity and is therefore sufficiently large whenever necessary.

2. The orientation rule

Here we describe the algorithm **Orient**. Its input is the edge process $\mathbf{e} = (e_1, e_2, \dots, e_{m_*})$, and output is an on-line orientation of each edge e_t . The algorithm proceeds in two steps. In the first step, which consists of the first $2n \log \log n$ edges, the algorithm builds a ‘core’ which contains almost all the vertices, and whose edges are distributed (almost) like a 6-in 6-out random graph. In the second step, which contains all edges that follow, the remaining $o(n)$ non-core vertices are taken care of, by being connected to the core in a way that will guarantee w.h.p. the existence of a directed Hamilton cycle.

2.1. Step I

Recall that each edge is given as an ordered pair (v, w) . For every vertex v we keep a count of the number of times v appears as the first vertex. We update the set of *saturated* vertices, which consists of the vertices that have appeared at least 12 times as the first vertex. Given the edge (v, w) at time t , if v is still not saturated, direct the edge (v, w)

alternatingly with respect to v starting from an out-edge. (By ‘alternatingly’ we mean that if the last edge having v as the first vertex was directed as an out-edge of v , then we direct the current one as an in-edge of v , and *vice versa*. For the first edge we choose the out direction.) Otherwise, if v is saturated, then count the number of times w has appeared as a second vertex when the first vertex is already saturated, and direct the edges alternatingly according to this count with respect to w starting from an in-edge. This alternation process is independent of the previous one. That is, even if w has previously appeared somewhere as a first vertex, the count should be kept track of separately from it.

For a vertex $v \in V$, let the *first vertex degree* of v be the number of times v has appeared as a first vertex in Step I, and denote it by $d_1(v)$. Let the *second vertex degree* of v be the number of times v has appeared in Step I as a second vertex of an edge whose first vertex is already saturated, and denote it by $d_2(v)$. Note that the sum of the first vertex degree and second vertex degree of v is not necessarily equal to the degree of v in Step I, as v might appear as a second vertex of an edge whose first vertex is not yet saturated. We will call such an edge a *neglected edge* of v .

2.2. Step II

Let A be the set of saturated vertices at the end of Step I, and $B = V \setminus A$. Call an edge an A – B edge if one end-point lies in A and the other end-point lies in B , and define A – A edges and B – B edges similarly. Given an edge $e = (v, w)$ at time t , if e is an A – B edge, and without loss of generality assume that $v \in B$ and $w \in A$, then direct e alternatingly with respect to v , where the alternation process of Step II continues the one from Step I as follows.

- (1) If v has appeared as a first vertex in Step I at least once, then pick up where the alternation process of v as a first vertex in Step I stopped, and continue the alternation.
- (2) If v did not appear as a first vertex in Step I but did appear as a second vertex of an already saturated vertex, then pick up where the alternation process of v as a second vertex of a saturated vertex stopped in Step I, and continue the alternation.
- (3) If v appeared in Step I but does not belong to the above two cases, then consider the first neglected edge connected to v , and start the alternation process from the opposite direction of this edge.
- (4) If none of the above, then start from an out-edge.

Otherwise, if e is an A – A edge or a B – B edge, orient it uniformly at random. Note that unlike Step I, the order of vertices of the given edge does not affect the orientation of the edge in Step II.

For a vertex $v \in B$, let the A – B degree of v be the number of A – B edges incident to v in Step II, and denote it by $d_{AB}(v)$. For $v \in A$, let $d_{AB}(v) = 0$.

3. Proof outline

Our approach builds on Frieze’s proof of Hamiltonicity of the random directed graph process [15] with some additional ideas. His proof consists of two phases (the original

proof consists of three phases, but for simplicity we describe it as two phases). We shall first describe these two phases of Frieze's proof, and then point out the modifications that are necessary to accommodate our different setting. Let $m = (1 + o(1))n \log n$ be the time at which the random directed graph process has minimum in-degree and out-degree 1, and let $D_{n,m}$ be the directed graph at time m (throughout this section we say that random directed graph has certain property if the property holds w.h.p.).

3.1. Phase 1: Find a small 1-factor

In Phase 1, a 1-factor of $D_{n,m}$ consisting of at most $O(\log n)$ cycles is constructed. To this end, a subgraph $D_{5\text{-in}, 5\text{-out}}$ of $D_{n,m}$ is constructed which uses only a small number of the edges. Roughly speaking, for each vertex, use its first 5 out-neighbours and 5 in-neighbours (if possible) to construct $D_{5\text{-in}, 5\text{-out}}$. Note that the resulting graph will be similar to a random 5-in 5-out directed graph, but still different, as some vertices will only have 1 in-neighbour and 1 out-neighbour even at time m . Finally, viewing $D_{5\text{-in}, 5\text{-out}}$ as a bipartite graph $G'(V \cup V^*, E')$, where V^* is a copy of V , and $\{u, v^*\} \in E'$ if and only if $u \rightarrow v$ belongs to $D_{5\text{-in}, 5\text{-out}}$, one proves that G' has a perfect matching. It turns out that this matching can be viewed as a uniform random permutation of the set of vertices V . A well-known fact about such permutations is that w.h.p. they consist of at most $O(\log n)$ cycles.

3.2. Phase 2: Combining the cycles into a Hamilton cycle

In Phase 2, the cycles of the 1-factor are combined into a Hamilton cycle. The technical issue to overcome in this step is the fact that in order to construct $D_{5\text{-in}, 5\text{-out}}$ all of the edges were scanned, and now supposedly we have no remaining random edges in the process to combine the cycles of the 1-factor. However, note that since $D_{5\text{-in}, 5\text{-out}}$ consists of at most $10n$ edges, the majority of edges need not be exposed. More rigorously, let LARGE be the vertices whose degree is $\Omega(\log n / \log \log n)$ at time $t_0 = 2n \log n / 3$ in the directed graph process. For the LARGE vertices, its 5 neighbours in $D_{5\text{-in}, 5\text{-out}}$ will be determined solely by the edges up to time t_0 , leaving the remaining edges (edges after time t_0) of the process unexposed. Two key properties used in Phase 2 are that w.h.p. (a) $|\text{LARGE}| = n - o(n^{1/2})$, and (b) every cycle of the 1-factor contains many LARGE vertices. Note that by (a), out of the remaining $n \log n / 3$ edges, all but $o(1)$ -fraction will connect two LARGE vertices. Phase 2 can now be summarized by the following theorem, which can be read out from the proof in [15].

Theorem 3.1. *Let V be a set of n vertices and let $L \subset V$ be a subset of size at least $n - o(n^{1/2})$. Assume that D is a directed 1-factor over V consisting of at most $O(\log n)$ cycles, and the vertices $V \setminus L$ are at distance at least 10 away from each other in this graph.*

If $(1/3 - o(1))n \log n$ L - L edges are given uniformly at random, then w.h.p. the union of these edges and the graph D contains a directed Hamilton cycle.

3.3. Comparing with our setting

The main technical issue in this paper is to re-prove Phase 1, namely, the existence of a 1-factor with small number of cycles. In [15], the fact that all vertices have the same

degree distribution in $D_{5\text{-in},5\text{-out}}$, led to an argument showing the existence of a matching that translates into a uniform random permutation. Our case is different because of the orientation rule. We have different types of vertices, each being oriented in a different way, breaking the nice symmetry. The bulk of our technical work is spent in resolving this technical issue.

Once this is done, that is, after achieving the 1-factor, we come up with an analogue of LARGE, which we call ‘saturated’. As in Phase 2 described above, we prove that w.h.p. (a’) most of the vertices are saturated, and (b’) every cycle in the 1-factor contains many saturated vertices. However, the naive approach results in a situation where one cannot apply Theorem 3.1 ((a’) and (b’) are quantitatively weaker than (a) and (b)). Thus we develop the argument of ‘compressing’ vertices of a given cycle. This idea allows us to get rid of all the non-saturated vertices, leading to another graph which only has saturated vertices. Details will be given in Section 5.2. Once we apply the compression argument, we can use Theorem 3.1 to finish the proof. Let us mention that the compression argument can be applied after Phase 1 in [15] as well to simplify the proof.

4. A typical random process

The following well-known concentration result (see, e.g., [1, Corollary A.1.14]) will be used several times in the proof. We denote by $\text{Bi}(n, p)$ the binomial random variable with parameters n and p .

Theorem 4.1 (Chernoff’s inequality). *If $X \sim \text{Bi}(n, p)$ and $\varepsilon > 0$, then*

$$\mathbb{P}(|X - \mathbb{E}[X]| \geq \varepsilon \mathbb{E}[X]) \leq e^{-\Omega_\varepsilon(\mathbb{E}[X])}.$$

4.1. Classifying vertices

To analyse the algorithm it will be convenient to work with three sets of vertices. The first is the set of *saturated* vertices at Step I. Throughout we will use A to denote this set. Let us now consider the non-saturated vertices $B = V \setminus A$. Here we distinguish between two types. We say that $v \in B$ *blossoms* if there are at least 12 edges of the form $\{v, A\}$ in Step II (by A we mean an arbitrary vertex from A), and let B_1 be the collection of vertices which blossom. All the remaining vertices are *restricted*, and are denoted by B_2 . Thus every vertex is either saturated (A), blossoms (B_1), or restricted (B_2).

Furthermore, the set of restricted vertices has two important subclasses which are determined by the first vertex degree $d_1(v)$, second vertex degree $d_2(v)$, and A – B degree $d_{AB}(v)$ defined in the previous section. We say that a restricted vertex v *partially blossoms* if the sum of its first vertex degree, second vertex degree, and A – B degree is at least 2. Note that since we stopped the process when the graph has minimum degree 2, every vertex v has degree at least 2. Thus, if the above mentioned sum is at most 1, then v has either a neglected edge or a B – B edge connected to it. A useful fact that we prove in Lemma 4.5 says that w.h.p. all such vertices v have one A – B edge (thus $d_{AB}(v) = 1$), and at least one neglected edge. Thus, a restricted vertex v that is not partially blossomed, and has one A – B edge and at least one neglected edge, is called a *bud*.

4.2. Properties of a typical random process

In this section we list several properties that hold w.h.p. for random edge processes. We will call an edge process *typical* if the properties indeed hold. Let

$$m_1 = \frac{1}{2}n \log n + \frac{1}{2}n \log \log n - n \log \log \log n,$$

$$m_2 = \frac{1}{2}n \log n + \frac{1}{2}n \log \log n + n \log \log \log n.$$

Note that for a fixed vertex v , the probability of an edge being incident to v is $\frac{2n-1}{n^2} = \frac{2}{n} - \frac{1}{n^2}$ (this is because, in our process, each edge is given by an ordered pair of vertices). However, as it turns out, the $\frac{1}{n^2}$ term is always negligible for our purpose, so we will use the probability $\frac{2}{n}$ for this event, and remind the reader that the term $\frac{1}{n^2}$ is omitted. Recall that the stopping time m_* is the time at which the last vertex of degree one becomes a degree two vertex and the process stops.

Claim 4.2. *Let m_* be the stopping time of the random process. Then w.h.p.*

$$m_1 \leq m_* \leq m_2.$$

Proof. For a fixed vertex v , the probability of an edge being incident to v is about $\frac{2}{n}$. Hence the probability of v having degree at most 1 at time m_2 is

$$\left(1 - \frac{2}{n}\right)^{m_2} + \binom{m_2}{1} \frac{2}{n} \cdot \left(1 - \frac{2}{n}\right)^{m_2-1} \leq 3 \log n \cdot e^{-\log n - \log \log n - 2 \log \log \log n}$$

$$= O\left(\frac{1}{n(\log \log n)^2}\right).$$

Thus, by Markov’s inequality, w.h.p. there is no vertex of degree at most 1 after m_2 edges. This shows that $m_* \leq m_2$. Similarly, the expected number of vertices having degree at most 1 after seeing m_1 edges is $\Omega((\log \log n)^2)$, and by computing the second moment of the number of vertices having degree at most 1, we can show that after m_1 edges w.h.p. at least one such vertex exists. This shows that $m_* \geq m_1$. The rest of the details are fairly standard, and are omitted. □

Next we are going to list some properties of the different types of vertices.

Claim 4.3. *The number of saturated vertices satisfies w.h.p.*

$$|A| \geq n \left(1 - \frac{(\log \log n)^{12}}{\log^2 n}\right).$$

Proof. For a fixed vertex v , the probability of v occurring as the first vertex of an edge is (exactly) $\frac{1}{n}$, and thus the probability of v ending up non-saturated at Step I is at most

$$\sum_{k=0}^{11} \binom{2n \log \log n}{k} \left(\frac{1}{n}\right)^k \cdot \left(1 - \frac{1}{n}\right)^{2n \log \log n - k} \leq \sum_{k=0}^{11} (2 \log \log n)^k \frac{1}{\log^2 n} = O\left(\frac{(\log \log n)^{11}}{\log^2 n}\right).$$

The claim follows from Markov’s inequality. □

Our next goal is to prove that the restricted vertices consist only of partially blossomed and bud vertices. For that we need the following auxiliary lemma.

Claim 4.4. *Let E_{BB} be the collection of all B – B edges (in Step II). The graph $G_m \setminus E_{BB}$ has w.h.p. minimum degree 2.*

Proof. If the graph $G_m \setminus E_{BB}$ has minimum degree less than 2 for some edge process \mathbf{e} , then there exists a vertex v which gets at most one edge other than a B – B edge, and at least one B – B edge. By Claim 4.2, it suffices to prove that the graph w.h.p. does not contain a vertex which has at most one edge other than a B – B edge at time m_1 , and at least one B – B edge at time m_2 . Let \mathcal{A}_v be the event that v is such a vertex. Let \mathcal{BS} be the event that

$$|B| \leq \frac{(\log \log n)^{12}}{\log^2 n} n \quad (B \text{ is small}),$$

and note that $\mathbb{P}(\mathcal{BS}) = 1 - o(1)$ by Claim 4.3. Then we have

$$\mathbb{P}(G_m \setminus E_{BB} \text{ has minimum degree less than } 2) = \mathbb{P}\left(\bigcup_{v \in V} \mathcal{A}_v\right) \leq n \cdot \mathbb{P}(\mathcal{A}_v \cap \mathcal{BS}) + o(1). \tag{4.1}$$

The event \mathcal{A}_v is equivalent to the vertex v receiving k B – B edges, for some $k > 0$, and at most one edge other than a B – B edge at appropriate times. This event is contained in the event $\mathcal{C}_v \cap \mathcal{D}_{v,k}$, where \mathcal{C}_v is the event ‘ v appears at most once in Step I’, and $\mathcal{D}_{v,k}$ is the event ‘ $d_{AB}(v) \leq 1$ by time m_1 and v receives k B – B edges by time m_2 ’. Therefore our next goal is to bound

$$\mathbb{P}(\mathcal{C}_v \cap \mathcal{D}_{v,k} \cap \mathcal{BS}) = \mathbb{P}(\mathcal{C}_v \cap \mathcal{BS}) \cdot \mathbb{P}(\mathcal{D}_{v,k} | \mathcal{C}_v \cap \mathcal{BS}) \leq \mathbb{P}(\mathcal{C}_v) \cdot \mathbb{P}(\mathcal{D}_{v,k} | \mathcal{C}_v \cap \mathcal{BS}). \tag{4.2}$$

We can bound the probability of the event \mathcal{C}_v by

$$\left(1 - \frac{2}{n}\right)^{2n \log \log n} + \binom{2n \log \log n}{1} \left(\frac{2}{n}\right) \cdot \left(1 - \frac{2}{n}\right)^{2n \log \log n - 1} = O\left(\frac{\log \log n}{\log^4 n}\right). \tag{4.3}$$

To bound the event $\mathcal{D}_{v,k}$ which is ‘ $d_{AB}(v) \leq 1$ at time m_1 and v receives k B – B edges by time m_2 ’, note that \mathcal{C}_v and \mathcal{BS} are events which depend only on the first $2 \log \log n$ edges (Step I edges). Therefore conditioning on this event does not affect the distribution of edges in Step II (each edge is chosen uniformly at random among all possible n^2 pairs).

We only consider the case $d_{AB}(v) = 1$ (the case $d_{AB}(v) = 0$ can be handled similarly, and turns out to be dominated by the case $d_{AB}(v) = 1$). Thus, to bound the probability, we choose $k + 1$ edges among the $m_2 - 2n \log \log n$ edges, let one of them be an A - B edge, and let k of them be B - B edges incident to v . Moreover, since $d_{AB}(v) \leq 1$ at time m_1 , we know that at least $m_1 - 2n \log \log n - k - 1$ edges are not incident to v . Thus,

$$\begin{aligned} & \mathbb{P}(\mathcal{D}_{v,k} \mid \mathcal{C}_v \cap \mathcal{BS}) \\ & \leq \binom{m_2 - 2n \log \log n}{k + 1} \left(\frac{2}{n}\right)^{k+1} \binom{k + 1}{1} \frac{|A|}{n} \left(\frac{|B|}{n}\right)^k \left(1 - \frac{2}{n}\right)^{m_1 - 2n \log \log n - k - 1}. \end{aligned}$$

By using the inequalities $1 - x \leq e^{-x}$, $|A| \leq n$, and $\binom{m_2 - 2n \log \log n}{k+1} \leq m_2^{k+1}$, the probability above is bounded by

$$(k + 1)m_2^{k+1} \left(\frac{2}{n}\right)^{k+1} \left(\frac{|B|}{n}\right)^k \exp\left(-\frac{2}{n}(m_1 - 2n \log \log n - k - 1)\right). \tag{4.4}$$

Therefore, by (4.2), (4.3), and (4.4),

$$\begin{aligned} & \mathbb{P}(\mathcal{C}_v \cap \mathcal{D}_{v,k} \cap \mathcal{BS}) \\ & \leq O\left(\frac{\log \log n}{\log^4 n}\right) (k + 1)m_2^{k+1} \left(\frac{2}{n}\right)^{k+1} \left(\frac{|B|}{n}\right)^k \exp\left(-\frac{2}{n}(m_1 - 2n \log \log n - k - 1)\right). \end{aligned}$$

Plugging the bound $|B| \leq \frac{n \log \log^2 n}{\log^2 n}$ and $m_2 \leq n \log n$ in the latter, we obtain

$$O(k) \left(\frac{\log \log n}{\log^3 n}\right) \left(\frac{2(\log \log n)^{12}}{\log n}\right)^k \exp\left(-\frac{2}{n}(m_1 - 2n \log \log n - k - 1)\right).$$

By the definition $m_1 = \frac{1}{2}n \log n + \frac{1}{2}n \log \log n - n \log \log \log n$, this simplifies further to

$$O(k) \left(\frac{(\log \log n)^3}{n}\right) \left(\frac{2e^{2/n}(\log \log n)^{12}}{\log n}\right)^k.$$

Summing over all possible values of k ,

$$\sum_{k=1}^{\infty} \mathbb{P}(\mathcal{C}_v \cap \mathcal{D}_{v,k} \cap \mathcal{BS}) \leq \sum_{k=1}^{\infty} \frac{O(k)(\log \log n)^3}{n} \left(\frac{4(\log \log n)^{12}}{\log n}\right)^k = o(n^{-1}).$$

Going back to (4.1), we get that

$$\mathbb{P}(G_{m_*} \setminus E_{BB} \text{ has minimum degree less than } 2) = n \cdot o(n^{-1}) + o(1) = o(1).$$

Note that, as mentioned at the beginning of this section, we used $\frac{2}{n}$ to estimate the probability of an edge being incident to a fixed vertex. This probability is in fact $\frac{2}{n} - \frac{1}{n^2}$, but the term $\frac{1}{n^2}$ will only affect the lower-order estimates. \square

Claim 4.5. *Every restricted vertex is w.h.p. either partially blossomed or a bud.*

Proof. Assume there exists a restricted vertex v which is not partially blossomed or a bud. Then, by definition, the sum $d_1(v) + d_2(v) + d_{AB}(v) \leq 1$. The possible values of the degrees $(d_1(v), d_2(v), d_{AB}(v))$ are $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$, or $(0, 0, 0)$. Vertices which correspond to

$(0, 0, 1)$ will all be bud vertices w.h.p. by Claim 4.4. It suffices to show then that w.h.p. there do not exist vertices corresponding to $(1, 0, 0)$, $(0, 1, 0)$, or $(0, 0, 0)$. Let T be the collection of vertices which have $d_1(v) + d_2(v) \leq 1$ and $d_{AB}(v) = 0$ at time m_1 . By Claim 4.2 it suffices to prove that T is empty. Let \mathcal{BS} be the event

$$|\mathcal{B}| \leq \frac{(\log \log n)^{12}}{\log^2 n} n,$$

and note that by Claim 4.3, $\mathbb{P}(\mathcal{BS}) = 1 - o(1)$. The event $\{T \neq \emptyset\}$ is the same as $\cup_{v \in V} \{v \in T\}$, and thus by the union bound

$$\begin{aligned} \mathbb{P}(T \neq \emptyset) &\leq o(1) + \sum_{v \in V} \mathbb{P}(\{v \in T\} \cap \mathcal{BS}) \\ &= o(1) + \sum_{v \in V} \mathbb{P}(\{d_1(v) + d_2(v) \leq 1\} \cap \{d_{AB}(v) = 0\} \cap \mathcal{BS}). \end{aligned}$$

By Bayes' equation, the second term of the right-hand side splits into

$$\begin{aligned} &\sum_{v \in V} \mathbb{P}(\{d_1(v) + d_2(v) \leq 1\} \cap \mathcal{BS}) \cdot \mathbb{P}(d_{AB}(v) = 0 \mid \{d_1(v) + d_2(v) \leq 1\} \cap \mathcal{BS}) \\ &\leq \sum_{v \in V} \mathbb{P}(d_1(v) + d_2(v) \leq 1) \cdot \mathbb{P}(d_{AB}(v) = 0 \mid \{d_1(v) + d_2(v) \leq 1\} \cap \mathcal{BS}). \end{aligned} \tag{4.5}$$

The probability $\mathbb{P}(d_1(v) + d_2(v) \leq 1)$ can be bounded by $\mathbb{P}(\{d_1(v) \leq 1\} \cap \{d_2(v) \leq 1\})$, which satisfies

$$\mathbb{P}(\{d_1(v) \leq 1\} \cap \{d_2(v) \leq 1\}) = \mathbb{P}(d_1(v) \leq 1) \cdot \mathbb{P}(d_2(v) \leq 1 \mid d_1(v) \leq 1).$$

The term $\mathbb{P}(d_1(v) \leq 1)$ can be easily calculated by

$$\left(1 - \frac{1}{n}\right)^{2n \log \log n} + \binom{2n \log \log n}{1} \left(\frac{1}{n}\right) \cdot \left(1 - \frac{1}{n}\right)^{2n \log \log n - 1} = O\left(\frac{\log \log n}{\log^2 n}\right).$$

To estimate $\mathbb{P}(d_2(v) \leq 1 \mid d_1(v) \leq 1)$, expose the edges of Step I as follows. First expose all the first vertices. Then expose the second vertices whose first vertex is saturated ($d_2(v)$ is now determined for every $v \in V$). The number of second vertex spots that are considered is at least $2n \log \log n - 12n$, and thus $\mathbb{P}(d_2(v) \leq 1 \mid d_1(v) \leq 1)$ is at most

$$\left(1 - \frac{1}{n}\right)^{2n \log \log n - 12n} + \binom{2n \log \log n}{1} \left(\frac{1}{n}\right) \cdot \left(1 - \frac{1}{n}\right)^{2n \log \log n - 12n - 1} = O\left(\frac{\log \log n}{\log^2 n}\right).$$

Thus, as a crude bound we have

$$\mathbb{P}(d_1(v) + d_2(v) \leq 1) \leq \mathbb{P}(d_1(v) \leq 1) \cdot \mathbb{P}(d_2(v) \leq 1 \mid d_1(v) \leq 1) = O\left(\frac{(\log \log n)^2}{\log^4 n}\right).$$

Since $d_1(v) + d_2(v) \leq 1$ implies that $v \in \mathcal{B}$, and $d_{AB}(v)$ depends only on the Step II edges (which are independent from $d_1(v)$, $d_2(v)$, and \mathcal{BS}), the second term of the right-hand side of equation (4.5), the probability $\mathbb{P}(d_{AB}(v) = 0 \mid \{d_1(v) + d_2(v) \leq 1\} \cap \mathcal{BS})$ can be bounded

by

$$\begin{aligned} \left(1 - 2\frac{1}{n}\frac{|A|}{n}\right)^{m_1 - 2n \log \log n} &\leq \exp\left(-2(m_1 - 2n \log \log n)|A|/n^2\right) \\ &\leq \exp\left(-(\log n - 3 \log \log n - 2 \log \log \log n)\left(1 - \frac{(\log \log n)^{12}}{\log^2 n}\right)\right) \\ &\leq \exp\left(-\log n + 3 \log \log n + 2 \log \log \log n + o(1)\right) = O\left(\frac{(\log n)^3(\log \log n)^2}{n}\right). \end{aligned}$$

Therefore in (4.5),

$$\begin{aligned} \mathbb{P}(T \neq \emptyset) &\leq o(1) + \sum_{v \in V} O\left(\frac{(\log \log n)^2}{\log^4 n}\right) O\left(\frac{(\log n)^3(\log \log n)^2}{n}\right) \\ &= o(1) + O\left(\frac{(\log \log n)^4}{\log n}\right) = o(1). \end{aligned} \quad \square$$

Claim 4.6. *The following properties hold w.h.p. for restricted vertices:*

- (i) *there are at most $\log^{13} n$ such vertices,*
- (ii) *every two such vertices are at distance at least 3 in G_m , from each other.*

Proof. Since being a restricted vertex is a monotone decreasing property, by Claim 4.2 it suffices to prove (i) at time m_1 . Recall that B_2 is the collection of restricted vertices (a vertex is restricted if it is not saturated or blossomed).

First, condition on the whole outcome of Step I edges (first $2n \log \log n$ edges) and the event that

$$|B| \leq \frac{(\log \log n)^{12}}{\log^2 n} n.$$

Then the set B is determined, and for a vertex $v \in B$ we can bound the probability of the event $v \in B_2$ as follows:

$$\mathbb{P}(v \in B_2) \leq \sum_{\ell=0}^{11} \binom{m_2}{\ell} \left(\frac{2}{n}\right)^\ell \left(1 - \frac{2|A|}{n^2}\right)^{m_1 - 2n \log \log n - \ell}. \tag{4.6}$$

Use the inequalities $m_1 = \frac{1}{2}n \log n + \frac{1}{2}n \log \log n - \log \log \log n \leq n \log n$, $m_2 \leq n \log n$, $1 - x \leq e^{-x}$, and

$$|A| = n - |B| \geq n \left(1 - \frac{(\log \log n)^{12}}{\log^2 n}\right)$$

to bound the above by

$$\sum_{\ell=0}^{11} (2 \log n)^\ell \exp\left(-\left(\log n - 3 \log \log n - 2 \log \log \log n - \frac{2\ell}{n}\right)\left(1 - \frac{(\log \log n)^{12}}{\log^2 n}\right)\right).$$

The sum is dominated by $\ell = 11$, and this gives

$$O(\log^{11} n) \exp(-\log n + 3 \log \log n + 2 \log \log \log n + o(1)) \leq O\left(\frac{(\log \log n)^2 \log^{14} n}{n}\right).$$

Thus the expected size of B_2 given the Step I edges is

$$\mathbb{E}[|B_2| \mid \text{Step I edges}] \leq |B| \cdot O\left(\frac{(\log \log n)^2 \log^{14} n}{n}\right) \leq O((\log \log n)^{14} \log^{12} n).$$

Since the assumptions on A and B hold w.h.p. by Claim 4.3, we can use Markov's inequality to conclude that w.h.p. there are at most $\log^{13} n$ vertices in B_2 . Let us now prove (ii).

For three distinct vertices v_1, v_2 and w in V , let $\mathcal{A}(v_1, v_2, w)$ be the event that w is a common neighbour of v_1 and v_2 . The probability of there being edges (v_1, w) (or (w, v_1)) and (v_2, w) (or (w, v_2)) and $v_1, v_2 \in B_2$ can be bounded by first choosing two time slots where (v_1, w) (or (w, v_1)) and (v_2, w) (or (w, v_2)) will be placed, and then filling in the remaining edges so that $v_1, v_2 \in B_2$. We will only bound the event of there being edges (v_1, w) and (w, v_2) in the edge process (other cases can be handled in a similar manner). The probability we would like to bound is

$$\mathbb{P}(\exists v_1, v_2, w, \exists 1 \leq t_1, t_2 \leq m_2, e_{t_1} = (v_1, w), e_{t_2} = (w, v_2), v_1, v_2 \in B_2).$$

By the union bound this probability is at most

$$\sum_{v_1, v_2, w \in V} \sum_{t_1, t_2=1}^{m_2} \mathbb{P}(e_{t_1} = (v_1, w), e_{t_2} = (w, v_2), v_1, v_2 \in B_2) \tag{4.7}$$

$$\begin{aligned} &= \sum_{v_1, v_2, w \in V} \sum_{t_1, t_2=1}^{m_2} \mathbb{P}(e_{t_1} = (v_1, w), e_{t_2} = (w, v_2)) \mathbb{P}(v_1, v_2 \in B_2 \mid e_{t_1} = (v_1, w), e_{t_2} = (w, v_2)) \\ &\leq \frac{1}{n} \sum_{t_1, t_2=1}^{m_2} \mathbb{P}(v_1, v_2 \in B_2 \mid e_{t_1} = (v_1, w), e_{t_2} = (w, v_2)). \end{aligned} \tag{4.8}$$

To simplify the notation we abbreviate

$$\mathbb{P}(v_1, v_2 \in B_2 \mid e_{t_1} = (v_1, w), e_{t_2} = (w, v_2))$$

by

$$\mathbb{P}(v_1, v_2 \in B_2 \mid e_{t_1}, e_{t_2}).$$

By using the independence of Step I and Step II edges, we have

$$\mathbb{P}(v_1, v_2 \in B_2 \mid e_{t_1}, e_{t_2}) = \mathbb{P}(v_1, v_2 \in B \mid e_{t_1}, e_{t_2}) \mathbb{P}(v_1, v_2 \notin B_1 \mid v_1, v_2 \in B, e_{t_1}, e_{t_2}).$$

For fixed t_1 and t_2 , we can bound $\mathbb{P}(v_1, v_2 \in B \mid e_{t_1}, e_{t_2})$ by the probability of ' v_1 and v_2 appear at most 22 times combined in Step I as a first vertex other than at time t_1 and t_2 ',

whose probability can be bounded as follows, regardless of the value of t_1 and t_2 :

$$\begin{aligned} \sum_{k=0}^{22} \binom{2n \log \log n}{k} \left(\frac{2}{n}\right)^k \cdot \left(1 - \frac{2}{n}\right)^{2n \log \log n - 2 - k} &\leq \sum_{k=0}^{22} (4 \log \log n)^k \frac{O(1)}{\log^4 n} \\ &= O\left(\frac{(\log \log n)^{22}}{\log^4 n}\right). \end{aligned}$$

To bound $\mathbb{P}(v_1, v_2 \notin B_1 | v_1, v_2 \in B, e_{t_1}, e_{t_2})$, it suffices to bound $\mathbb{P}(v_1, v_2 \notin B_1 | v_1, v_2 \in B, e_{t_1}, e_{t_2}, \mathcal{BS})$, which can be bounded by the probability of ‘ v_1 and v_2 receive in total at most 22 A - B edges in Step II other than at time t_1 and t_2 ’. Regardless of the value of t_1 and t_2 , this satisfies the bound

$$\sum_{\ell=0}^{22} \binom{m_2}{\ell} \left(\frac{4}{n}\right)^\ell \left(1 - \frac{4|A|}{n}\right)^{m_1 - 2 - 2n \log \log n - \ell}.$$

Note that the $\frac{4}{n}$ and $\frac{2}{n}$ terms throughout the proof should in fact involve some terms of order $\frac{1}{n^2}$, but these are omitted for simplicity since they do not affect the final asymptotic outcome. By a similar calculation to (4.6), this can eventually be bounded by $O\left(\frac{\log^{29} n}{n^2}\right)$. Thus we have

$$\mathbb{P}(v_1, v_2 \in B_2 | e_{t_1}, e_{t_2}) = O\left(\frac{\log^{26} n}{n^2}\right),$$

which by (4.8) and $m_2 \leq n \log n$ gives

$$\mathbb{P}(\exists v_1, v_2, w, \exists 1 \leq t_1, t_2 \leq m_2, e_{t_1} = (v_1, w), e_{t_2} = (w, v_2), v_1, v_2 \in B_2) \leq O\left(\frac{\log^{28} n}{n}\right).$$

Therefore, by Markov’s inequality, w.h.p. no such three vertices exist, which implies that two vertices $v_1, v_2 \in B_2$ cannot be at distance two from each other in G_m . Similarly, we can prove that w.h.p. every two vertices $v_1, v_2 \in B_2$ are not adjacent to each other, and hence w.h.p. every $v_1, v_2 \in B_2$ are at a distance at least two away from each other. \square

4.3. Configuration of the edge process

To prove that our algorithm succeeds with high probability, we first reveal some pieces of information on the edge process, which we call the ‘configuration’ of the process. This information will allow us to determine whether the underlying edge process is typical. Then, in the next section, using the remaining randomness, we will construct a Hamilton cycle.

At the beginning, rather than thinking of edges coming one by one, we regard our edge process $\mathbf{e} = (e_1, e_2, \dots, e_{m_*})$ as a collection of edges e_i for $i = 1, \dots, m_*$, neither of whose end-points are known. We can decide to reveal certain information as necessary. Let us first reveal the following.

- (1) For $t \leq 2n \log \log n$, reveal the first vertex of the t th edge e_t . If this vertex has already appeared as the first vertex at least 12 times among the edges e_1, \dots, e_{t-1} , then also reveal the second vertex.

Given this information, we can determine the saturated vertices, and hence we know the sets A and B . Therefore, it is possible to reveal the following information.

(2) For $t > 2n \log \log n$, reveal all the vertices that belong to B .

The information we have revealed determines the blossomed (B_1), and restricted (B_2) vertices. Thus we can reveal the following information.

(3) For $t \leq 2n \log \log n$, further reveal all the non-revealed vertices that belong to B_2 .

(4) For every edge $e_t = (v_t, w_t)$ in which we already know that either $v_t \in B_2$ or $w_t \in B_2$, reveal the other vertex too.

We define the *configuration* of an edge process as the above four pieces of information.

We want to say that all the non-revealed vertices are uniformly distributed over certain sets. But in order for this to be true, we must make sure that the distribution of the non-revealed vertices is not affected by the fact that we know the value of m_* (some vertex has degree exactly 2 at time m_* , and maybe a non-revealed vertex will make this vertex have degree 2 earlier than m_*). This is indeed the case, since the last vertex to have degree 2 is necessarily a restricted vertex, and all the locations of the restricted vertices are revealed. Thus the non-revealed vertices cannot change the value of m_* . Therefore, once we condition on the configuration of an edge process, the remaining vertices are distributed in the following way.

(a) For $t \leq 2n \log \log n$, if the first vertex of the edge e_t has appeared at most 12 times among e_1, \dots, e_{t-1} , then its second vertex is either a known vertex in B_2 or is a random vertex in $V \setminus B_2$.

(b) For $t > 2n \log \log n$, if both vertices of e_t are not revealed, then e_t consists of two random vertices of A . If only one of the vertices of e_t is not revealed, then the revealed vertex is in B , and the non-revealed vertex is a random vertex of A .

Definition 4.7. A configuration of an edge process is *typical* if it satisfies the following.

(i) The number of saturated and blossomed vertices satisfies

$$|A| \geq n - \frac{(\log \log n)^{12}}{\log^2 n} n, \quad |B_1| \leq \frac{(\log \log n)^{12}}{\log^2 n} n,$$

respectively.

(ii) The number of restricted vertices satisfies $|B_2| \leq \log^{13} n$.

(iii) Every vertex appears at least twice in the configuration even without considering the B - B edges.

(iv) All the restricted vertices are either partially blossomed or buds.

(v) In the non-directed graph induced by the edges both of whose end-points are revealed, every two restricted vertices v_1, v_2 are at distance at least 3 away from each other.

(vi) There are at least $\frac{1}{3}n \log n$ edges e_t for $t > 2n \log \log n$ neither of whose end-points are revealed.

Lemma 4.8. *The random edge process has a typical configuration w.h.p.*

Proof. The fact that the random edge process has w.h.p. a configuration satisfying (i), (iii) and (iv) follows from Claims 4.3, 4.4, and 4.5 respectively; (ii) and (v) follow from Claim 4.6. To verify (vi), note that by Claim 4.2 and 4.3, w.h.p. there are at least $\frac{1}{2}n \log n - 2n \log \log n$ edges of Step II, and $|A| = (1 - o(1))n$. Therefore the probability of a Step II edge being an A - A edge is $1 - o(1)$, and the expected number of A - A edges is $(1/2 - o(1))n \log n$. Then, by Chernoff's inequality, w.h.p. there are at least $\frac{1}{3}n \log n$ A - A edges. These edges are the edges we are looking for in (vi). \square

5. Finding a Hamilton cycle

In the previous section we established several useful properties of the underlying graph G_{m_n} . In this section, we study the algorithm **Orient** using these properties, and prove that, conditioned on the edge process having a typical configuration, the graph D_{m_n} w.h.p. contains a Hamilton cycle (recall that the graph D_{m_n} is the set of random edges of the edge process, oriented according to **Orient**). As described in Section 3, the proof is a constructive proof, in the sense that we describe how to find such a cycle. The algorithm is similar to that used in [15], which we described in some detail in Section 3. Let us briefly recall that it proceeds in two stages.

- (1) Find a 1-factor of G . If it contains more than $O(\log n)$ cycles, fail.
- (2) Join the cycles into a Hamilton cycle.

The main challenge in our case is to prove that the first step of the algorithm does not fail. Afterwards, we argue why we can apply Frieze's results for the remaining step.

5.1. Almost 5-in 5-out subgraph

Let $D_{5\text{-in}, 5\text{-out}}$ be the following subgraph of D_{m_n} . For each vertex v , assign a set of neighbours $\text{OUT}(v)$ and $\text{IN}(v)$, where $\text{OUT}(v)$ are out-neighbours of v and $\text{IN}(v)$ are in-neighbours of v . For saturated and blossomed vertices, $\text{OUT}(v)$ and $\text{IN}(v)$ will be of size 5, and for restricted vertices, they will be of size 1 (thus $D_{5\text{-in}, 5\text{-out}}$ is not a 5-in 5-out directed graph under the strict definition).

Let E_1 be the edges of Step I (first $2n \log \log n$ edges), and let E_2 be the edges of Step II (remaining edges).

- If v is saturated, then consider the first 12 appearances in E_1 of v as a first vertex. Some of these edges might later be used as OUT or IN for other vertices. Hence, among these 12 appearances, consider only those whose second vertex is not in B_2 . By property (v) of Definition 4.7, there will be at least 11 such second vertices for a typical configuration. Define $\text{OUT}(v)$ as the first 5 vertices among them which were directed out from v , and $\text{IN}(v)$ as the first 5 vertices among them which were directed in to v in **Orient**.
- If v blossoms, then consider the first 10 A - B edges in E_2 connected to v , and look at the other end-points. Let $\text{OUT}(v)$ be the first 5 vertices which are an out-neighbour of v and let $\text{IN}(v)$ be the first 5 vertices which are an in-neighbour of v .

A partially blossomed vertex, by definition, has $d_1(v) + d_2(v) + d_{AB}(v) \geq 2$, and must fall into one of the following categories: (i) $d_1(v) \geq 2$, (ii) $d_2(v) \geq 2$, (iii) $d_{AB}(v) \geq 2$,

(iv) $d_1(v) = 1, d_2(v) = 1$, (v) $d_1(v) = 1, d_{AB}(v) = 1$, and (vi) $d_1(v) = 0, d_2(v) = 1, d_{AB}(v) = 1$. If it falls into several categories, then pick the first one among them.

- If v partially blossoms and $d_1(v) \geq 2$, consider the first two appearances of v in E_1 as a first vertex. The first is an out-edge and the second is an in-edge (see Section 2.1).
- If v partially blossoms and $d_2(v) \geq 2$, consider the first two appearances of v in E_1 as a second vertex whose first vertex is saturated. The first is an in-edge and the second is an out-edge (see Section 2.1).
- If v partially blossoms and $d_{AB}(v) \geq 2$, consider the first two A – B edges in E_2 incident to v . One is an out-edge and the other is an in-edge. Note that, unlike other cases, the actual order of in-edge and out-edge will depend on the configuration. But since the configuration contains all the positions at which v appeared in the process, the choice of in-edge or out-edge only depends on the configuration and not on the non-revealed vertices (note that this is slightly different from the blossomed vertices).
- If v partially blossoms and $d_1(v) = 1, d_2(v) = 1$, consider the first appearance of v in E_1 as a first vertex, and the first appearance of v in E_1 as a second vertex whose first vertex is saturated. The former is an out-edge and the latter is an in-edge.
- If v partially blossoms and $d_1(v) = 1, d_{AB}(v) = 1$, consider the first appearance of v in E_1 as a first vertex, and the first A – B edge connected to v in E_2 . The former is an out-edge and the latter is an in-edge (see rule (1) in Section 2.2).
- If v partially blossoms and $d_1(v) = 0, d_2(v) = 1, d_{AB}(v) = 1$, consider the first appearance of v in E_1 as a second vertex whose first vertex is saturated, and the first A – B edge connected to v in E_2 . The former is an in-edge and the latter is an out-edge (see rule (2) in Section 2.2). Thus we can construct $\text{OUT}(v)$ and $\text{IN}(v)$ of size 1 each, for all partially blossomed vertices.
- If v is a bud, then consider the first (and only) A – B edge connected to v . Let this edge be e_s . For a typical configuration, by property (iii) of Definition 4.7, we know that v has a neglected edge connected to it. Let e_t be the first neglected edge of v . By property (v) of Definition 4.7, we know that the first vertex of the neglected edge is either in A or B_1 . According to the direction of this edge, the direction of e_s will be chosen as the opposite direction (see rule (3) in Section 2.2). As in the partially blossomed case with $d_{AB}(v) \geq 2$, the direction is solely determined by the configuration. Thus we can construct $\text{OUT}(v)$ and $\text{IN}(v)$ of size 1 each (which is already fixed once we have fixed the configuration).

This in particular shows that D_m has minimum in-degree and out-degree at least 1, which is clearly a necessary condition for the graph to be Hamiltonian. A crucial observation is that, once we condition on the random edge process having a fixed typical configuration, we can determine exactly which edges are going to be used to construct the graph $D_{5\text{-in}, 5\text{-out}}$ just by looking at the configuration.

For a set X , let $RV(X)$ be an element chosen independently and uniformly at random in the set (consider each appearance of $RV(X)$ as a new independent copy).

Proposition 5.1. *Let $V' = V \setminus B_2$. Conditioned on the edge process having a typical configuration, $D_{5\text{-in}, 5\text{-out}}$ has the following distribution.*

- (i) If v is saturated, then $OUT(v)$ and $IN(v)$ are a union of 5 copies of $RV(V')$.
- (ii) If v blossoms, then $OUT(v)$ and $IN(v)$ are a union of 5 copies of $RV(A)$.

Proof. For a vertex $v \in V$, the configuration contains the information on the time of arrival of the edges that will be used to construct the set $OUT(v)$ and $IN(v)$.

If v is a saturated vertex, then we even know which edges belong to $OUT(v)$ and $IN(v)$ (if there are no B_2 vertices connected to the first 12 appearances of v as a first vertex, then the first five odd appearances of v as a first vertex will be used to construct $OUT(v)$, and the first five even appearances of v as a first vertex will be used to construct $IN(v)$). Since the non-revealed vertices are independent random vertices in V' , we know that $OUT(v)$ and $IN(v)$ of these vertices consist of 5 independent copies of $RV(V')$.

If v blossoms, then the analysis is similar to that of the saturated vertices. However, even though the configuration contains the information as to which 10 edges will be used to construct $OUT(v)$ and $IN(v)$, the decision of whether the odd edges or the even edges will be used to construct $OUT(v)$ depends on the particular edge process (this is determined by the orientation rule at Step I). However, since the other end-points are independent identically distributed random vertices in A , the distribution of $OUT(v)$ and $IN(v)$ is not affected by the previous edges, and is always $RV(A)$ (this is analogous to the fact that the distribution of the outcome of a coin flip does not depend on whether the initial position was head or tail). \square

5.2. A small 1-factor

The main result that we are going to prove in this section is summarized in the following proposition.

Proposition 5.2. *Conditioned on the random edge process having a typical configuration, there exists w.h.p. a 1-factor of $D_{5\text{-in}, 5\text{-out}}$ containing at most $2 \log n$ cycles, and in which at least $9/10$ of each cycle are saturated vertices.*

Throughout this section, rather than vaguely conditioning on the process having a typical configuration, we will consider a fixed typical configuration \mathbf{c} and condition on the event that the edge process has configuration \mathbf{c} . Proposition 5.2 easily follows once we prove that there exists a Hamilton cycle w.h.p. under this assumption. The reason we do this more precise conditioning is to fix the sets A, B, B_1, B_2 and the edges incident to vertices of B_2 (note that these are determined solely by the configuration). In our later analysis, it is crucial to have these fixed.

To prove Proposition 5.2, we represent the graph $D_{5\text{-in}, 5\text{-out}}$ as a certain bipartite graph in which a perfect matching corresponds to the desired 1-factor of the original graph D_m . Then, using the edge distribution of $D_{5\text{-in}, 5\text{-out}}$ given in the previous section, we will show that the bipartite graph w.h.p. contains a perfect matching. The proof of Proposition 5.2 will be given at the end after a series of lemmas.

Define a new vertex set $V^* = \{v^* | v \in V\}$ as a copy of V , and for sets $X \subset V$ use X^* to denote the set of vertices in V^* corresponding to X . Then, in order to find a 1-factor in $D_{5\text{-in}, 5\text{-out}}$, define an auxiliary bipartite graph $\mathbf{BIP}(V, V^*)$ over the vertex set

$V \cup V^*$ whose edges are given as follows. For every (directed) edge (u, v) of $D_{5\text{-in}, 5\text{-out}}$, add the (undirected) edge (u, v^*) to **BIP**. Note that perfect matchings of **BIP** have a natural one-to-one correspondence with 1-factors of $D_{5\text{-in}, 5\text{-out}}$. Moreover, the edge distribution of **BIP** easily follows from the edge distribution of $D_{5\text{-in}, 5\text{-out}}$. We will say that $D_{5\text{-in}, 5\text{-out}}$ is the *underlying directed graph* of **BIP**. A permutation σ of V^* acts on **BIP** to construct another bipartite graph which has edges $(v, \sigma(w^*))$ for all edges (v, w^*) in **BIP**.

Our plan is to find a perfect matching which is (almost) a uniform random permutation, and show that this permutation has at most $O(\log n)$ cycles (if it were a uniform random permutation, then this would be a well-known result: see, e.g., [13]). Since our distribution is not a uniform distribution, we will rely on the following lemma. Its proof is rather technical, and to avoid distraction, it will be given at the end of this subsection.

Lemma 5.3. *Let X be subset of V . Assume that w.h.p. (i) **BIP** contains a perfect matching, (ii) every cycle of the underlying directed graph $D_{5\text{-in}, 5\text{-out}}$ contains at least one element from X , and (iii) the edge distribution of **BIP** is invariant under arbitrary permutations of X^* . Then w.h.p. there exists a perfect matching which, when considered as a permutation, contains at most $2 \log n$ cycles.*

The next set of lemmas establishes the fact that **BIP** satisfies all the conditions we need in order to apply Lemma 5.3. First we prove that **BIP** contains a perfect matching. We use the following version of the well-known Hall’s theorem (see, e.g., [11]).

Theorem 5.4. *Let Γ be a bipartite graph with vertex set $X \cup Y$ and $|X| = |Y| = n$. If, for all $X' \subset X$ of size $|X'| \leq n/2$, $|N(X')| \geq |X'|$ and for all $Y' \subset Y$ of size $|Y'| \leq n/2$, $|N(Y')| \geq |Y'|$, then G contains a perfect matching.*

Lemma 5.5. *The graph **BIP** contains a perfect matching w.h.p.*

Proof. We will verify Hall’s condition for the graph **BIP** to prove the existence of a perfect matching. Recall that **BIP** is a bipartite graph over the vertex set $V \cup V^*$.

Let us show that every set $D \subset V$ of size $|D| \leq n/2$ satisfies $|N(D)| \geq |D|$. This will be done in two steps. First, if $D \subset B_2$, then this follows from the fact that $\text{OUT}(v)$ are distinct sets for all $v \in B_2$ (if they were not distinct, then there would be two restricted vertices at distance two away, and this violates property (v) of Definition 4.7). Second, we prove that for $D \subset V \setminus B_2$ and $|D| \leq n/2$,

$$|N(D) \cap (V^* \setminus N(B_2))| \geq |D|.$$

It is easy to see that the above two facts prove our claim.

Let $D \subset V \setminus B_2$ be a set of size at most $k \leq n/2$. The inequality $|N(D) \cap (V^* \setminus N(B_2))| < |D|$ can happen only if there exists a set $N^* \subset V^* \setminus N(B_2)$ such that $|N^*| < k$, and for all $v \in D$ all the vertices of $\text{OUT}(v)$ belong to $N^* \cup N(B_2)$. Since $D \subset V \setminus B_2$, every vertex in D has 5 random neighbours distributed uniformly over some set of size $(1 - o(1))n$, and

thus the probability of the above event happening is at most

$$k \binom{n}{k}^2 \left(\frac{|N(B_2)| + |N^*|}{(1 - o(1))n} \right)^{5k} \leq \left(\frac{e^2 n^2 (\log^{13} n + k)^5}{k^2 \cdot (1 - o(1))n^5} \right)^k \leq \left(\frac{9(\log^{13} n + k)^5}{k^2 n^3} \right)^k.$$

For the range $9n/20 \leq k \leq n/2$, we will use the following bound:

$$k \binom{n}{k}^2 \left(\frac{\log^{13} n + k}{(1 - o(1))n} \right)^{5k} \leq 2^{2n} \left(\frac{1 + o(1)}{2} \right)^{9n/4} \leq 2^{-n/5}.$$

Summing over all choices of k , we obtain

$$\begin{aligned} & \sum_{k=1}^{n/2} k \binom{n}{k}^2 \left(\frac{\log^{13} n + k}{(1 - o(1))n} \right)^{5k} \\ & \leq \sum_{k=1}^{\log^{14} n} \left(\frac{9(\log^{13} n + k)^5}{k^2 n^3} \right)^k + \sum_{k=\log^{14} n}^{9n/20} \left(\frac{9(\log^{13} n + k)^5}{k^2 n^3} \right)^k + \sum_{k=9n/20}^{n/2} 2^{-n/5} \\ & \leq \sum_{k=1}^{\log^{14} n} \left(\frac{10 \log^{70} n}{n^3} \right)^k + \sum_{k=\log^{14} n}^{9n/20} \left(\frac{10k^3}{n^3} \right)^k + o(1) = o(1). \end{aligned}$$

This finishes the proof that w.h.p. $|N(D)| \geq |D|$ for all $D \subset V$ of size at most $n/2$. Similarly, for sets $D^* \subset V^*$ of size $|D^*| \leq n/2$, using the sets $\text{IN}(v)$ instead of $\text{OUT}(v)$ we can show that w.h.p. $|N(D^*)| \geq |D^*|$ in **BIP**. \square

For restricted vertices v , the sets $\text{OUT}(v)$ and $\text{IN}(v)$ are of size 1 and are already fixed, since we have fixed the configuration. Thus the edge corresponding to these vertices will be in **BIP**. Let

$$\hat{A} = A \setminus (\cup_{v \in B_2} \text{OUT}(v)),$$

and let \hat{A}^* be the corresponding set inside V^* (note that \hat{A} and \hat{A}^* are fixed sets). This set will be our set X when applying Lemma 5.3. We next prove that every cycle of $D_{5\text{-in}, 5\text{-out}}$ contains vertices of \hat{A} .

Lemma 5.6. *With high probability, every cycle C of $D_{5\text{-in}, 5\text{-out}}$ contains at least $\lceil \frac{9}{10} |C| \rceil$ vertices of \hat{A} .*

Proof. Recall that by Proposition 5.1, for vertices $v \in V \setminus B_2$, the set $\text{OUT}(v)$ and $\text{IN}(v)$ are uniformly distributed over $V \setminus B_2$, or A . Therefore, for a vertex $w \in B_2$, the only out-neighbour of w is $\text{OUT}(w)$, and the only in-neighbour is $\text{IN}(w)$ (note that they are both fixed since we have fixed the configuration). Also, note that

$$|V \setminus \hat{A}| \leq |V \setminus A| + |B_2| \leq |B_1| + 2|B_2| \leq \frac{(\log \log n)^{12}}{\log^2 n} n + 2 \log^{13} n \leq \frac{n}{\log n}.$$

We want to show that in the graph $D_{5\text{-in}, 5\text{-out}}$, w.h.p. every cycle of length k has at most $k/10$ points from $V \setminus \hat{A}$, for all $k = 1, \dots, n$. Let us compute the expected number of cycles for which this condition fails and show that it is $o(1)$. First choose k vertices v_1, v_2, \dots, v_k

(with order) and assume that a of them are in B_2 . Then, since we already know the (unique) out-neighbour and in-neighbour for vertices in B_2 , for the vertices v_1, \dots, v_k to form a cycle in that order, we must fix $3a$ positions (a for the vertices in B_2 , and $2a$ for their in- and out-neighbours by property (v) of Definition 4.7). Assume that among the remaining $k - 3a$ vertices, ℓ vertices belong to $V \setminus (\hat{A} \cup B_2)$. Then, for there to be at least $\lceil k/10 \rceil$ vertices among v_1, \dots, v_k not in \hat{A} , we must have $3a + \ell \geq \lceil k/10 \rceil$. There are at most 3^k ways to assign one of the three types \hat{A}, B_2 , and $V \setminus (\hat{A} \cup B_2)$ to each of v_1, \dots, v_k . Therefore the number of ways to choose k vertices as above is at most

$$3^k \cdot n^{k-\ell-3a} |V \setminus \hat{A}|^\ell |B_2|^a \leq 3^k \cdot n^{k-\ell-3a} \left(\frac{n}{\log n}\right)^\ell (\log^{13} n)^a.$$

There are $k - 2a$ random edges that have to be present in order to make the above k vertices into a cycle. For all $i \leq k - 1$, the pair (v_i, v_{i+1}) can become an edge either by $v_{i+1} \in \text{OUT}(v_i)$ or $v_i \in \text{IN}(v_{i+1})$ (and also for the pair (v_1, v_k)). There are two ways to choose where the edge $\{v_i, v_{i+1}\}$ comes from, and if both v_i and v_{i+1} are not in B_2 , then $\{v_i, v_{i+1}\}$ will become an edge with probability at most $\frac{5}{(1-o(1))n}$. Therefore the probability of a fixed v_1, \dots, v_k chosen as above being a cycle is at most

$$2^{k-2a} \left(\frac{5}{(1-o(1))n}\right)^{k-2a},$$

and the expected number of such cycles is at most

$$\begin{aligned} & 2^{k-2a} \left(\frac{5}{(1-o(1))n}\right)^{k-2a} \cdot 3^k \cdot n^{k-\ell-3a} \left(\frac{n}{\log n}\right)^\ell (\log^{13} n)^a \\ & \leq \left(\frac{\log^{13} n}{n}\right)^a \cdot \left(\frac{1}{\log n}\right)^\ell \cdot (30 + o(1))^k \\ & \leq \left(\frac{\log^{13} n}{n}\right)^a \cdot \left(\frac{1}{\log n}\right)^{\lceil k/10 \rceil - 3a} \cdot (30 + o(1))^k \leq \left(\frac{\log^{16} n}{n}\right)^a \cdot \left(\frac{40}{(\log n)^{1/10}}\right)^k, \end{aligned}$$

where we used $3a + \ell \geq \lceil k/10 \rceil$ for the second inequality. Sum this over $0 \leq \ell \leq k$ and $0 \leq a \leq k$ and we get

$$\sum_{k=1}^n \sum_{\ell=0}^k \sum_{a=0}^k \left(\frac{\log^{16} n}{n}\right)^a \cdot \left(\frac{40}{(\log n)^{1/10}}\right)^k = O\left(\sum_{k=1}^n (k+1) \left(\frac{40}{(\log n)^{1/10}}\right)^k\right) = o(1),$$

which proves our lemma. □

The following simple observation is the last ingredient of our proof.

Lemma 5.7. *The distribution of **BIP** is invariant under the action of an arbitrary permutation of \hat{A}^* .*

Proof. This lemma follows from the following three facts about the distribution of $D_{5\text{-in}, 5\text{-out}}$. First, all the saturated vertices have the same distribution of IN. Second, for the vertices $v \in V \setminus B_2$, the distribution of OUT and IN is uniform over a set which contains

all the saturated vertices (for some vertices it is $V \setminus B_2$, and for others it is A). Third, for the vertices $v \in B_2$, the set $\text{OUT}(v)$ lies outside \hat{A} by definition. Therefore, the action of an arbitrary permutation of \hat{A}^* does not affect the distribution of **BIP**. \square

Note that here it is important to fix the configuration beforehand, as otherwise the set \hat{A}^* will vary and a statement such as Lemma 5.7 will not make sense.

By combining Lemmas 5.3, 5.5, 5.6 and 5.7, we obtain Proposition 5.2.

Proof of Proposition 5.2. Lemmas 5.5, 5.6, and 5.7 show that the graph **BIP** has all the properties required for the application of Lemma 5.3 (we use $X = \hat{A}$). Thus we know that w.h.p., $D_{5\text{-in}, 5\text{-out}}$ has a 1-factor containing at most $2 \log n$ cycles, and in which at least $9/10$ of each cycle are saturated vertices (the second property by Lemma 5.6). \square

We conclude this subsection with the proof of Lemma 5.3.

Proof of Lemma 5.3. For simplicity of notation, we use the notation \mathcal{B} for the random bipartite graph **BIP**. Note that both a 1-factor over the vertex set V and a perfect matching of (V, V^*) can be considered as a permutation of V . Throughout this proof we will not distinguish between these interpretations, and treat both 1-factors and perfect matchings as permutations.

First, let f be an arbitrary function which, for every bipartite graph, outputs one fixed perfect matching in it. Then, given a bipartite graph Γ over the vertex set $V \cup V^*$, let Φ be the random variable $\Phi(\Gamma) := \tau^{-1}f(\tau\Gamma)$, where τ is a permutation of the vertices \hat{A}^* chosen uniformly at random. Since the distribution of \mathcal{B} and the distribution of $\tau\mathcal{B}$ are the same by condition (iii), for an arbitrary permutation σ of \hat{A}^* , Φ has the following property:

$$\begin{aligned} \mathbb{P}(\Phi(\mathcal{B}) = \phi) &= \mathbb{P}(\tau^{-1}f(\tau\mathcal{B}) = \phi) \stackrel{(*)}{=} \mathbb{P}((\tau\sigma)^{-1}f(\tau\sigma\mathcal{B}) = \phi) \\ &= \mathbb{P}(\tau^{-1}f(\tau\sigma\mathcal{B}) = \sigma\phi) \stackrel{(*)}{=} \mathbb{P}(\tau^{-1}f(\tau\mathcal{B}) = \sigma\phi) = \mathbb{P}(\Phi(\mathcal{B}) = \sigma\phi). \end{aligned} \tag{5.1}$$

In the (*) steps, we used (iii), and the fact that if τ is a uniform random permutation of \hat{A}^* then so is $\tau\sigma$, and therefore \mathcal{B} , $\tau\mathcal{B}$ and $\tau\sigma\mathcal{B}$ all have identical distribution.

Define a map Π from the 1-factors over the vertex set V to the 1-factors over the vertex set \hat{A} obtained by removing all the vertices that belong to $V \setminus \hat{A}$ from every cycle. For example, a cycle of the form $(x_1x_2y_1y_2x_3y_3x_4)$ will become the cycle $(x_1x_2x_3x_4)$ when mapped by Π (where $x_1, \dots, x_4 \in \hat{A}$, and $y_1, y_2, y_3 \in V \setminus \hat{A}$). Note that if all the original 1-factors contained at least one element from \hat{A} , then the total number of cycles does not change after applying the map Π . This observation combined with condition (ii) implies that it suffices to obtain a bound on the number of cycles after applying Π .

Let σ, ρ be permutations of the vertex set \hat{A}^* . We claim that for every 1-factor ϕ of the vertex set V , the equality $\sigma \cdot \Pi(\phi) = \Pi(\sigma \cdot \phi)$ holds. This claim together with (5.1) gives us

$$\begin{aligned} \mathbb{P}(\Pi(\Phi(\mathcal{B})) = \rho) &= \mathbb{P}(\Phi(\mathcal{B}) \in \Pi^{-1}(\rho)) \stackrel{(5.1)}{=} \mathbb{P}(\sigma\Phi(\mathcal{B}) \in \Pi^{-1}(\rho)) = \mathbb{P}(\Pi(\sigma\Phi(\mathcal{B})) = \rho) \\ &= \mathbb{P}(\sigma \cdot \Pi(\Phi(\mathcal{B})) = \rho) = \mathbb{P}(\Pi(\Phi(\mathcal{B})) = \sigma^{-1}\rho). \end{aligned}$$

Since σ and ρ were arbitrary permutations of the vertex set \hat{A} , we can conclude that, conditioned on there existing a perfect matching, $\Pi(\Phi(\mathcal{B}))$ has a uniform distribution over the permutations of \hat{A} . It is a well-known fact (see, e.g., [13]) that a uniformly random permutation over a set of size n has w.h.p. at most $2 \log n$ cycles. Since \mathcal{B} w.h.p. contains a perfect matching by condition (i), it remains to verify the equality $\sigma \cdot \Pi(\phi) = \Pi(\sigma \cdot \phi)$. Thus we conclude the proof by proving this claim.

For a vertex $x \in \hat{A}$, assume the cycle of ϕ containing x is of the form $(\cdots xy_1y_2 \cdots y_kx_+ \cdots)$ ($k \geq 0$) for $y_1, \dots, y_k \in V \setminus \hat{A}$. Then, by definition $\Pi(\phi)(x) = x_+$, and thus $(\sigma \cdot \Pi(\phi))(x) = \sigma(x_+)$. On the other hand, since σ only permutes \hat{A} and fixes every other element of V , we have $(\sigma \cdot \phi)(x) = \sigma(y_1) = y_1$, and $(\sigma \cdot \phi)(y_i) = y_{i+1}$ for all $i \leq k - 1$, and $(\sigma \cdot \phi)(y_k) = \sigma(x_+)$. Therefore the cycle in $\sigma \cdot \phi$ which contains x will be of the form $(\cdots xy_1y_2 \cdots y_k\sigma(x_+) \cdots)$, and then by definition we have $(\Pi(\sigma \cdot \phi))(x) = \sigma(x_+)$. \square

5.3. Combining the cycles into a Hamilton cycle

Assume that, as in the previous subsection, we started with a fixed typical configuration \mathbf{c} , conditioned on the edge process having configuration \mathbf{c} , and found a 1-factor of $D_{5\text{-in}, 5\text{-out}}$ by using Proposition 5.2. Since this 1-factor only uses the edges which have been used to construct the graph $D_{5\text{-in}, 5\text{-out}}$, it is independent of the A - A edges in Step II that we did not reveal. Moreover, by the definition of a typical configuration, there are at least $\frac{1}{3}n \log n$ such edges. Note that the algorithm gives a random direction to these edges. So interpret this as receiving $\frac{1}{3}n \log n$ randomly directed A - A edges with repeated edges allowed. Then the problem of finding a directed Hamilton cycle in D_m can be reduced to the following problem.

Let V be a given set and let A be a subset of size $(1 - o(1))n$. Assume that we are given a 1-factor over this vertex set, where at least 9/10 of each cycle lies in the set A . If we are given $\frac{1}{3}n \log n$ additional A - A edges chosen uniformly at random, can we find a directed Hamilton cycle?

To further simplify the problem, we remove the vertices $V \setminus A$ from the picture. Given a 1-factor over the vertex set V , mark in red all the vertices not in A . Pick any red vertex v , and assume that $v_-, v, v_+ \in V$ appear in this order in some cycle of the given 1-factor. If $v_- \neq v_+$, replace the three vertices v_-, v, v_+ by a new vertex v' , where v' takes as in-neighbours the in-neighbours of v_- , and as out-neighbours the out-neighbours of v_+ . We call the above process a *compression* of the three vertices v_-, v, v_+ . A crucial property of compression is that every 1-factor of the compressed graph corresponds to a 1-factor in the original graph (with the same number of cycles). Since a directed Hamilton cycle is also a 1-factor, if we can find a Hamilton cycle in the compressed graph, then we can also find one in the original graph.

Now, for each $v \in V \setminus A$ compress the three vertices v_-, v, v_+ into a vertex v' and mark it red if and only if either v_- or v_+ is a red vertex. This process always decreases the number of red vertices. Repeat it until there are no red vertices remaining, or $v_- = v_+$ for all red vertices v . As long as there is no red vertex in a cycle of length 2 at any point of the process, the latter will not happen. Consider a cycle whose length was k at the beginning. Since at least 9/10 of each cycle comes from A and every compression

decreases the number of vertices by 2, at any time there will be at least $(8/10)k$ non-red vertices, and at most $(1/10)k$ red vertices remaining in the cycle. Thus, if a cycle has a red vertex, then its length will be at least 9, and this prevents length 2 red cycles. So the compressing procedure will be over when all the red vertices disappear. Note that since $|V \setminus A| = |B| = o(n)$, the number of remaining vertices after the compression procedure is over is at least $n - 2|B| = (1 - o(1))n$. As mentioned above, it suffices to find a Hamilton cycle in the graph after the compression process is over.

Another important property of this procedure is related to the additional A - A edges that we are given. Assume that v is the first red vertex that we have compressed where the vertices v_-, v, v_+ appeared in this order in some 1-factor. Further assume that v_- and v_+ are not red vertices. Then, since the new vertex v' obtained from the compression will take as out-neighbours the out-neighbours of v_+ , and as in-neighbours the in-neighbours of v_- , we may assume that this vertex v' is a vertex in A from the perspective of the new $\frac{1}{3}n \log n$ edges that will be given.

This observation shows that every pair of vertices of the compressed graph has the same probability of being one of the new $\frac{1}{3}n \log n$ edges. Since the number of vertices reduced by $o(n)$, only $o(n \log n)$ of the new edges will be lost because of the compression. Thus w.h.p. we will be given $(\frac{1}{3} - o(1))n \log n$ new uniform random edges of the compressed graph.

Theorem 5.8. *For a typical configuration \mathbf{c} , conditioned on the random edge process having configuration \mathbf{c} , the directed graph D_m w.h.p. contains a Hamilton cycle.*

Proof. By Proposition 5.2, there exists w.h.p. a perfect matching of **BIP** which corresponds to a 1-factor in D_m , consisting of at most $2 \log n$ cycles. Also, at least $9/10$ of the vertices in each cycle lies in A . After using the compression argument discussed above, we may assume that we are given a 1-factor over some vertex set of size $(1 - o(1))n$. Moreover, the random edge process contains at least $(\frac{1}{3} - o(1))n \log n$ additional random directed edges (distributed uniformly over that set). By Theorem 3.1, with L being the whole vertex set, we can conclude that w.h.p. the compressed graph contains a directed Hamilton cycle, and this in turn implies that D_m contains a directed Hamilton cycle. \square

Corollary 5.9. *The directed graph D_m w.h.p. contains a Hamilton cycle.*

Proof. Let \mathbf{e} be a random edge process. Let $D = D_m(\mathbf{e})$ and let \mathcal{Ham} be the collection of directed graphs that contain a directed Hamilton cycle. For a configuration \mathbf{c} , denote by $\mathbf{e} \in \mathbf{c}$ the event that \mathbf{e} has configuration \mathbf{c} . If $\mathbf{e} \in \mathbf{c}$ for some typical configuration \mathbf{c} , then we say that \mathbf{e} is typical.

By Theorem 5.8, we know that for any typical configuration \mathbf{c} , $\mathbb{P}(D \notin \mathcal{Ham} | \mathbf{e} \in \mathbf{c}) = o(1)$, from which we know that $\mathbb{P}(\{D \notin \mathcal{Ham}\} \cap \{\mathbf{e} \text{ is typical}\}) = o(1)$. On the other hand, by Lemma 4.8 we know that the probability of an edge process having a non-typical configuration is $o(1)$. Therefore w.h.p. the directed graph D_m is Hamiltonian. \square

6. Going back to the original process

Recall that the distribution of the random edge process is slightly different from that of the random graph process since it allows repeated edges and loops. In fact, one can show that at time m_* , the edge process w.h.p. contains at least $\Omega(\log^2 n)$ repeated edges. Therefore, to obtain our main theorem for random graph processes, we cannot simply condition on the event that the edge process does not contain any repeated edges or loops. Our next theorem shows that there exists an on-line algorithm **OrientPrime** which successfully orients the edges of the random graph process.

Theorem 6.1. *There exists a randomized on-line algorithm **OrientPrime** which orients the edges of the random graph process, so that the resulting directed graph is Hamiltonian w.h.p. at the time at which the underlying graph has minimum degree 2.*

The algorithm **OrientPrime** will mainly follow **Orient** but with a slight modification. Assume that we are given a random graph process (call it the underlying process). Using this random graph process, we want to construct an auxiliary process whose distribution is identical to the random edge process. Let $t = 1$ at the beginning and let a_t be the number of distinct edges up to time t in our auxiliary process (disregarding loops). Thus $a_1 = 0$. At time t , with probability $(2a_t + n)/n^2$ we will produce a redundant edge, and with probability $1 - (2a_t + n)/n^2$ we will receive an edge from the underlying random graph process. Once we have decided to produce a redundant edge, with probability $2a_t/(2a_t + n)$ we choose uniformly at random an edge out of the a_t edges that have already appeared, and with probability $n/(2a_t + n)$ choose uniformly at random a loop. Let e_t be the edge produced at time t (it is either a redundant edge or an edge from the underlying process), and choose its first and second vertex uniformly at random. One can easily check that the process (e_1, e_2, \dots) has the same distribution as the random edge process.

In the algorithm **OrientPrime** we feed this new auxiliary process into the algorithm **Orient** and orient the edges accordingly. Since the distribution of the auxiliary process is the same as that of the random edge process, **Orient** will give an orientation which w.h.p. contains a directed Hamilton cycle. However, what we are seeking is a Hamilton cycle with no redundant edge. Thus, in the edge process, whenever we see a redundant edge that is a repeated edge (not a loop), we colour it blue. In order to show that **OrientPrime** gives a Hamiltonian graph with high probability, it suffices to show that we can find a Hamilton cycle in D_m , which does not contain a blue edge (note that loops cannot be used in constructing a Hamilton cycle). We first state two useful facts.

Claim 6.2. *With high probability, there are no blue edges incident to B used in constructing $D_{5\text{-in}, 5\text{-out}}$.*

Proof. The expected number of blue edges incident to B in Step I used in constructing $D_{5\text{-in}, 5\text{-out}}$ can be computed by choosing two vertices v and w , and then computing the probability that $v \in B$, and (v, w) or (w, v) together appears twice among Step I edges. The probability that v appears as a first vertex exactly i times is $\binom{n \log \log n}{i} \left(\frac{1}{n}\right)^i \left(1 - \frac{1}{n}\right)^{n \log \log n - i}$.

Condition on the event that v has appeared i times as a first vertex for some $i < 12$ (and also reveal the i positions in which v appeared). We then compute the probability that some two Step I edges are (v, w) or (w, v) . There are three events that we need to consider: first the event that (v, w) appears twice, whose probability is $\binom{i}{2} \left(\frac{1}{n}\right)^2$; second the event that (v, w) appears once and (w, v) appears once, whose probability is at most $\binom{n \log \log n}{1} \frac{1}{n(n-1)} \cdot \binom{i}{1} \frac{1}{n}$; third the event that (w, v) appears twice, whose probability is at most $\binom{n \log \log n}{2} \left(\frac{1}{n(n-1)}\right)^2$. Combining everything, we see that the expected number of Step I blue edges incident to B is at most

$$n^2 \cdot \sum_{i=0}^{11} \binom{n \log \log n}{i} \left(\frac{1}{n}\right)^i \left(1 - \frac{1}{n}\right)^{n \log \log n - i} \\ \times \left(\binom{i}{2} \left(\frac{1}{n}\right)^2 + \binom{n \log \log n}{1} \left(\frac{1}{n(n-1)}\right) \binom{i}{1} \frac{1}{n} + \binom{n \log \log n}{2} \left(\frac{1}{n(n-1)}\right)^2 \right).$$

The main term comes from $i = 11$, and the third term in the final bracket. Consequently, we can bound the expectation by

$$(1 + o(1)) \cdot n^2 \cdot \binom{n \log \log n}{11} \left(\frac{1}{n}\right)^{11} \left(1 - \frac{1}{n}\right)^{n \log \log n - 11} \cdot \left(\binom{n \log \log n}{2} \left(\frac{1}{n(n-1)}\right)^2\right) \\ = o(1).$$

We would then like to compute the expected number of blue edges incident to B in Step 2 used in constructing $D_{5\text{-in}, 5\text{-out}}$. Condition on the first vertices of the Step I edges so that we can determine the sets A and B . By Claim 4.3, we may condition on the event

$$|B| = O\left(\frac{(\log \log n)^{12}}{\log^2 n}\right).$$

Fix a vertex $v \in B$, and expose all appearances of v in Step II, and note that only the first 10 appearances are relevant. By Claim 4.4, it suffices to bound the probability of the event that there exists a vertex $w \in A$ such that (v, w) or (w, v) appears twice among the at most 24 Step I edges where v or w are the first vertices, and the at most 10 Step II edges which we know are going to be used to construct the OUT and IN of the vertex v . Therefore the expectation is

$$|B| \cdot n \cdot \left(\frac{34}{n}\right)^2 = O\left(\frac{(\log \log n)^{12}}{\log^2 n} n^2\right) \cdot \left(\frac{34}{n}\right)^2 = o(1). \quad \square$$

Claim 6.3. *With high probability, there are at most $(\log \log n)^3$ blue edges used in constructing $D_{5\text{-in}, 5\text{-out}}$.*

Proof. By Claim 6.2, we know that w.h.p. all the blue edges used in constructing $D_{5\text{-in}, 5\text{-out}}$ are incident to A . Therefore it suffices to show that there are at most $\log n$ blue edges among the Step I edges. The expected number of such edges can be computed by choosing two vertices v, w , and computing the probability that (v, w) or (w, v) appears twice. This is

at most

$$n^2 \cdot \binom{n \log \log n}{2} \left(\frac{2}{n^2}\right)^2 = o((\log \log n)^3).$$

Consequently, by Markov's inequality we can derive the conclusion. \square

Claim 6.4. *With high probability, each vertex is incident to at most one blue edge.*

Proof. It suffices to show that there do not exist three distinct vertices v, w_1, w_2 such that both $\{v, w_1\}$ and $\{v, w_2\}$ appear at least twice. The probability of this event is at most

$$\binom{n}{3} \binom{m_2}{4} \cdot \binom{4}{2} \left(\frac{2}{n^2}\right)^4 = o(1). \quad \square$$

Now assume that we have found a 1-factor as in Section 5.2. After performing the compression process given in the beginning of Section 5.3, by Claim 6.2, no blue edges will 'disappear' during the process. Therefore, if we can find a Hamilton cycle in the compressed graph with no blue edge, then the original graph will also have a Hamilton cycle with no blue edge. Since the compressed graph can be considered as a graph given over a subset of A , we will use the additional non-revealed A - A edges to find a directed Hamilton cycle in the compressed graph that uses no blue edge. By Claims 6.3 and 6.4, it suffices to prove the following theorem in order to conclude that **OrientPrime** succeeds with high probability.

Theorem 6.5. *Suppose we are given a 1-factor over a vertex set of size $(1 - o(1))n$ consisting of $O(\log n)$ cycles, and let G be the graph obtained by adding to this 1-factor $\Omega(n \log n)$ independent uniformly chosen random edges. Then w.h.p. for all matchings H that intersect the 1-factor in at most $(\log \log n)^3$ edges, the graph $G - H$ contains a directed Hamilton cycle.*

One can prove this theorem by taking various approaches, such as those used in [9] or [15]. However, since we want to avoid using blue (repeated) edges, none of these proofs directly apply, and it is indeed necessary to make at least some degree of modification. Since these adjustments are rather straightforward, we omit the details here, and will provide them in the arXiv version of our paper for interested readers.

7. Concluding remarks

In this paper we have considered the following natural question. Consider a random edge process where, at each time t , a random edge (u, v) arrives. We are to give an on-line orientation to each edge at the time of its arrival. At what time t^* can one make the resulting directed graph Hamiltonian? The best that one can hope for is to have a Hamilton cycle when the last vertex of degree one disappears, and we prove that this is indeed achievable with high probability.

The main technical difficulty in the proof arose from the existence of bud vertices. These were degree two vertices that were adjacent to a saturated vertex in the auxiliary graph $D_{5\text{-in}, 5\text{-out}}$. Note that for our proof we used the method of deferred decisions, not exposing the end-points of certain edges and leaving them as random variables. Bud vertices precluded us from doing this naively and forced us to expose the end-point of some of the edges which we wanted to keep unexposed (it is not difficult to show that without exposing these end-points we cannot guarantee that the bud vertices have degree at least 2). If one is willing to settle for an asymptotically tight upper bound on t^* , then one can choose $t^* = (1 + \varepsilon)n \log n/2$, and then for $n = n(\varepsilon)$ sufficiently large there are no bud vertices. Moreover, since for this range of t^* the vertices will have significantly larger degree, the orienting rule can also be simplified. While not making the analysis ‘trivial’ (i.e., an immediate consequence of the work in [15]), this will considerably simplify the proof.

Acknowledgement

We are grateful to Alan Frieze for generously sharing this problem with us, and we thank Igor Pak for reference [13]. We would also like to thank the two referees for their valuable comments.

References

- [1] Alon, N. and Spencer, J. (2000) *The Probabilistic Method*, second edition, Wiley.
- [2] Azar, Y., Broder, A., Karlin, A. and Upfal, E. (1999) Balanced allocations. *SIAM J. Comput.* **29** 180–200.
- [3] Balogh, J., Bollobás, B., Krivelevich, M., Muller, T. and Walters, M. (2011) Hamilton cycles in random geometric graphs. *Ann. Appl. Probab.* **21** 1053–1072.
- [4] Bohman, T. and Frieze, A. (2001) Avoiding a giant component. *Random Struct. Alg.* **19** 75–85.
- [5] Bohman, T., Frieze, A. and Wormald, N. (2004) Avoidance of a giant component in half the edge set of a random graph. *Random Struct. Alg.* **25** 432–449.
- [6] Bohman, T. and Kravitz, D. (2006) Creating a giant component. *Combin. Probab. Comput.* **15** 489–511.
- [7] Bollobás, B. (1984) The evolution of sparse graphs. In *Graph Theory and Combinatorics: Proc. Cambridge Combinatorial Conference in Honour of Paul Erdős, 1984* (B. Bollobás, ed.), pp. 335–357.
- [8] Bollobás, B., Fenner, T. and Frieze, A. (1985) An algorithm for finding Hamilton cycles in random graphs. In *Proc. 17th Annual ACM Symposium on Theory of Computing, 1985*, pp. 430–439.
- [9] Cooper, C. and Frieze, A. (1994) Hamilton cycles in a class of random directed graphs. *J. Combin. Theory Ser. B* **62** 151–163.
- [10] Cooper, C. and Frieze, A. (2000) Hamilton cycles in random graphs and directed graphs. *Random Struct. Alg.* **16** 369–401.
- [11] Diestel, R. (2005) *Graph Theory*, Vol. 173 of *Graduate Texts in Mathematics*, third edition, Springer.
- [12] Erdős, P. and Rényi, A. (1960) On the evolution of random graphs. *Publ. Math. Inst. Hung. Acad. Sci.* **5A** 17–61.
- [13] Erdős, P. and Turán, P. (1965) On some problems of a statistical group-theory I. *Z. Wahrsch. Verw. Gebiete* **4** 175–186.

- [14] Flaxman, A., Gamarnik, D. and Sorkin, G. (2005) Embracing the giant component. *Random Struct. Alg.* **27** 277–289.
- [15] Frieze, A. (1988) An algorithm for finding Hamilton cycles in random directed graphs. *J. Algorithms* **9** 181–204.
- [16] Frieze, A. Personal communication.
- [17] Komlós, J. and Szemerédi, E. (1983) Limit distribution for the existence of Hamilton cycles in random graphs. *Discrete Math.* **43** 55–63.
- [18] Korshunov, A. (1976) Solution of a problem of Erdős and Rényi on Hamilton cycles in non-oriented graphs. *Soviet Math. Dokl.* **17** 760–764.
- [19] Krivelevich, M., Loh, P. and Sudakov, B. (2009) Avoiding small subgraphs in Achlioptas processes. *Random Struct. Alg.* **34** 165–195.
- [20] Krivelevich, M., Lubetzky, E. and Sudakov, B. (2010) Hamiltonicity thresholds in Achlioptas processes. *Random Struct. Alg.* **37** 1–24.
- [21] Pósa, L. (1976) Hamiltonian circuits in random graphs. *Discrete Math.* **14** 359–364.
- [22] Robinson, R. and Wormald, N. C. (1994) Almost all regular graphs are Hamiltonian. *Random Struct. Alg.* **5** 363–374.
- [23] Sinclair, A. and Vilenchik, D. (2010) Delaying satisfiability for random 2SAT. In *APPROX-RANDOM*, pp. 710–723.
- [24] Spencer, J. and Wormald, N. (2007) Birth control for giants. *Combinatorica* **27** 587–628.